



ANNALES
HENRI LEBESGUE

CLIFTON CUNNINGHAM

DAVID ROE

COMMUTATIVE CHARACTER SHEAVES AND GEOMETRIC TYPES FOR SUPERCUSPIDAL REPRESENTATIONS

FAISCEAUX CARACTÈRES COMMUTATIFS ET
TYPES GÉOMÉTRIQUES POUR LES
REPRÉSENTATIONS SUPERCUSPIDALES

ABSTRACT. — We show that some types for supercuspidal representations of tamely ramified p -adic groups that appear in Jiu-Kang Yu's work are geometrizable. To do so, we define a function-sheaf dictionary for one-dimensional characters of arbitrary smooth group schemes over finite fields. In previous work we considered the case of commutative smooth group schemes and found that the standard definition of character sheaves produced a dictionary with a nontrivial kernel. In this paper we give a modification of the category of character sheaves that remedies this defect, and is also extensible to non-commutative groups. We then use these *commutative character sheaves* to geometrize the linear characters that appear in the types introduced by Jiu-Kang Yu, assuming that the character vanishes on a certain derived subgroup. To define *geometric types*, we combine commutative character sheaves with Gurevich and Hadani's geometrization of the Weil representation and Lusztig's character sheaves.

Keywords: function-sheaf dictionary, commutative character sheaves, types for supercuspidal representations.

2020 *Mathematics Subject Classification*: 14F06, 14L15, 22E50.

DOI: <https://doi.org/10.5802/ahl.105>

(*) Roe was supported by Simons Foundation grant 550033.

RÉSUMÉ. — Nous montrons que certains types associés à des représentations supercuspidales de groupes p -adiques modérément ramifiés qui apparaissent dans les travaux de Jiu-Kang Yu sont géométrisables. Pour cela, nous définissons un dictionnaire entre fonctions et faisceaux pour les caractères de dimension un de schémas en groupes arbitraires sur les corps finis. Dans un travail précédent, nous avons considéré le cas des schémas en groupes lisses commutatifs et nous avons montré que la définition standard des faisceaux caractères donne lieu à un dictionnaire avec un noyau non trivial. Dans cet article, nous proposons une modification de la catégorie des faisceaux caractères qui corrige ce défaut et s'étend aux groupes non commutatifs. Nous utilisons ensuite ces *faisceaux caractères commutatifs* pour géométriser les caractères linéaires qui apparaissent dans les types introduits par Jiu-Kang Yu, en supposant que le caractère s'annule sur un certain sous-groupe dérivé. Pour définir les *types géométriques*, nous combinons les faisceaux caractères commutatifs avec la géométrisation par Gurevich et Hadani de la représentation de Weil et les faisceaux caractères de Lusztig.

Introduction

The combined work of Ju-Lee Kim in [Kim07] and Jessica Fintzen in [Fin18, Fin19] establishes that all irreducible supercuspidal representations of a tamely ramified p -adic group G can be built from “data” introduced by Jiu-Kang Yu [Yu01, § 15], as long as p does not divide the order of the Weyl group of G . While the type, in the sense of Bushnell & Kutzko [BK98], of a supercuspidal representation built from Yu data can be constructed directly from the datum, it is convenient to consider an intermediate object, introduced in [Yu01, Remark 15.4], which we call a *Yu type datum*. Yu type data are studied in [Yu15], which concludes with the following observation.

Therefore, up to some linear characters, all the ingredient representations are on groups of the form $\underline{H}(\mathcal{O})$, where \underline{H} is a smooth group scheme over [a Henselian discrete valuation ring with finite residue field κ] \mathcal{O} , and the representations are inflated from $\underline{H}(\kappa)$. These results suggest that algebraic geometry and group schemes should play an important role in the representation theory of p -adic groups.

In this paper we follow the suggestion above by showing that certain Yu type data are geometrizable, in the following sense. A Yu type datum determines a sequence of representations ${}^\circ\rho_i$ of compact p -adic groups ${}^\circ K^i$, for $i = 0, \dots, d$, such that $({}^\circ K^d, {}^\circ\rho_d)$ is a type for a supercuspidal representation of a p -adic group. Let R be the ring of integers of a local field with finite residue field k . The main result of [Yu15] shows how to find, for each $i = 0, \dots, d$, a smooth group scheme \underline{G}^i over the ring R with $\underline{G}^i(R) = {}^\circ K^i$. Under certain assumptions on the Yu type datum, we show how each representation ${}^\circ\rho_i$ can be replaced by a pair $(\underline{G}^i, \mathcal{F}^i)$, where \mathcal{F}^i is a formal \mathbb{Q} -linear combination of conjugation-equivariant sheaf complexes on the Greenberg transform G^i of \underline{G}^i . Writing $t_{\mathcal{F}^i}$ for the function on $G^i(k) = \underline{G}^i(R) = {}^\circ K^i$ obtained by evaluating the trace of the action of Frobenius on \mathcal{F}^i , we show in Theorem 5.2 that

$$(0.1) \quad t_{\mathcal{F}^i} = \mathrm{Tr}({}^\circ\rho_i).$$

By this theorem, then, we obtain geometric avatars for each type in a Yu datum:

$$\begin{array}{ccc}
 & \text{geometrization} & \\
 \left({}^\circ K^i, {}^\circ \rho_i \right) & \xrightarrow{\quad \quad \quad} & \left(\underline{G}^i, \mathcal{F}^i \right) \\
 & \xleftarrow{\quad \quad \quad} & \\
 & \text{trace of Frob} &
 \end{array}$$

We refer to the pair $(\underline{G}^d, \mathcal{F}^d)$ as a *geometric type*.

To prove Theorem 5.2, we must find a way to geometrize linear characters of groups of the form $\underline{H}(R)$, where \underline{H} is a smooth group scheme over R . In order to do so in a systematic manner, we begin this paper by describing the function-sheaf dictionary for characters of arbitrary smooth group schemes over finite fields. When coupled with the Greenberg transform, this dictionary will allow for the geometrization of certain linear characters of $\underline{H}(R)$.

The function-sheaf dictionary over a finite field k [Del77, Sommes trig.] provides a way of encoding functions on the k -rational points of an algebraic group G as ℓ -adic local systems on G . More specifically, if G is a connected, commutative, algebraic group then there is a certain category $\mathcal{CS}(G)$ of rank-one local systems on G and an explicit isomorphism between isomorphism classes of objects in $\mathcal{CS}(G)$ and $G(k)^* := \text{Hom}(G(k), \bar{\mathbb{Q}}_\ell^\times)$; the isomorphism is given by mapping \mathcal{L} to the function $\text{Tr}_G : g \mapsto \text{Tr}(\text{Fr} | \mathcal{L}_g)$.

In previous work [CR18], we studied the function-sheaf dictionary for characters smooth commutative group schemes G , allowing for non-connected groups. We gave a description of the category $\mathcal{CS}(G)$ in this context, as well as an epimorphism $\text{Tr}_G : \mathcal{CS}(G)_{/\text{iso}} \rightarrow G(k)^*$. In contrast to the connected case, Tr_G may have nontrivial kernel; we gave an explicit description of its kernel as $H^2(\pi_0(\bar{G}), \bar{\mathbb{Q}}_\ell^\times)^{\text{Fr}}$ [CR18, Theorem 3.6].

In this paper we repair this defect in the function-sheaf dictionary by describing a full subcategory $\mathcal{CCS}(G)$ of $\mathcal{CS}(G)$ so that Tr_G restricts to an isomorphism $\mathcal{CCS}(G)_{/\text{iso}} \rightarrow G(k)^*$. We refer to objects of $\mathcal{CS}(G)$ as character sheaves and objects in $\mathcal{CCS}(G)$ as *commutative character sheaves*, since the passage from $\mathcal{CS}(G)$ to $\mathcal{CCS}(G)$ involves a condition that exchanges the inputs to the multiplication morphism on G (see Definition 2.1). When G is connected, all character sheaves on G are commutative.

Category $\mathcal{CCS}(G)$ clarifies several questions about $\mathcal{CS}(G)$. Invisible character sheaves [CR18, Definition 2.8] are precisely those \mathcal{L} with $\text{Tr}_G(\mathcal{L}) = 1$ that are not commutative. Moreover, $\text{Tr}_G^{-1} : G(k)^* \rightarrow \mathcal{CCS}(G)_{/\text{iso}}$ provides a canonical splitting of $\text{Tr}_G : \mathcal{CS}(G)_{/\text{iso}} \rightarrow G(k)^*$ [CR18, Remark 3.7].

Next, we broaden our scope further to encompass smooth group schemes G over k that are not necessarily commutative. We assume G is smooth, but not that it is connected, reductive or commutative; we write $G_{\text{ab}} := G/G_{\text{der}}$ for the abelianization. The category $\mathcal{CS}(G)$ has a straightforward generalization to this case. We then define category $\mathcal{CCS}(G)$ for such G and a forgetful functor to $\mathcal{CS}(G)$ so that $\text{Tr}_G : \mathcal{CCS}(G)_{/\text{iso}} \rightarrow G_{\text{ab}}(k)^*$ is an isomorphism. Since $G_{\text{ab}}(k)^*$ maps onto $G(k)^*$ with cokernel $G_{\text{der}}(k)^*$, it follows that for each character $\chi \in G(k)^*$ trivial on $G_{\text{der}}(k)$ there is a commutative character sheaf \mathcal{L} on G with $\text{Tr}_G(\mathcal{L}) = \chi$. Moreover, we find

that pullback along the quotient $q : G \rightarrow G_{\text{ab}}$ defines an equivalence of categories $\mathcal{CCS}(G_{\text{ab}}) \rightarrow \mathcal{CCS}(G)$. The functor $\mathcal{CCS}(G) \rightarrow \mathcal{CS}(G)$ is not essentially surjective, missing the kinds of linear character sheaves highlighted by Kamgarpour in [Kam09, (1.1)].

In order to provide further justification for referring to objects in $\mathcal{CCS}(G)$ as commutative character sheaves, suppose for the moment that G is a connected, reductive algebraic group over k . Let $\bar{\mathcal{L}}$ be the geometric part of an object in $\mathcal{CCS}(G)$; see Section 1. Let T be a maximal torus in the base change \bar{G} of G to \bar{k} and let $\bar{\mathcal{L}}_T$ be the restriction of $\bar{\mathcal{L}}$ to T . Then the perverse sheaf $\bar{\mathcal{L}}[\dim G]$ appears in the semisimple complex $\text{ind}_{B,T}^{\bar{G}}(\bar{\mathcal{L}}_T)$ produced by parabolic induction. It follows that every object in $\mathcal{CCS}(G)$ determines a Frobenius-stable character sheaf on G , in the sense of [Lus85, Definition 2.10]. Of course, the sheaves arising in this way represent a small part of Lusztig’s geometrization of characters of representations of connected, reductive groups over finite fields, but they are precisely those needed to describe one-dimensional characters of such groups.

Armed with the function-sheaf dictionary for smooth group schemes over finite fields, we return to the task of geometrizing Yu type data. The proof of Theorem 5.2 requires: Yu’s work on smooth integral models [Yu15]; the geometrization of the character of the Heisenberg–Weil representation over finite fields by Gurevich & Hadani [GH07]; Lusztig’s character sheaves on reductive groups over finite fields; and finally, the function-sheaf dictionary for characters of smooth group schemes over finite fields, now at our disposal in Theorem 3.12.

In order to use these tools we must restrict our attention to Yu type data that satisfy a technical condition, appearing in Section 5.4 as Hypothesis **(H1)**. Granting this hypothesis, these pieces are assembled in Section 5.4, where we prove Theorem 5.2. With this theorem, we provide all of the ingredients needed to geometrize a class of supercuspidal representations of arbitrary depth.

Yu’s construction has been revisited recently by several authors. Hakim [Hak18, § 2.1] gives an alternate construction beginning with a representation of a compact-mod-center subgroup of G . Unfortunately, this ingredient does not lend itself to geometrization using our techniques. In contrast, Kaletha [Kal19] has given a construction in terms of a tame elliptic maximal torus $S \subset G$ and a character θ of S . The pair (S, θ) can already be geometrized using [CR18] since S is commutative. However, Kaletha’s construction only produces a subset of Yu’s supercuspidal representations [Kal19, Proposition 3.7.14], so we phrase our geometrization in terms of Yu’s original construction.

We now summarize the sections of the paper in more detail. In Section 1, we recall the category $\mathcal{CS}(G)$ from [CR18] and note that it still makes sense when G is not commutative. We focus on the case of commutative G in Section 2, giving the definition of a commutative character sheaf and proving our first main theorem, that $\text{Tr}_G : \mathcal{CS}(G)_{/\text{iso}} \rightarrow G(k)^*$ induces an isomorphism on $\mathcal{CCS}(G)_{/\text{iso}}$. Passing to the case that G is non-commutative, we give the definition of and main results about commutative character sheaves in Section 3. We note that we should only consider character sheaves that arise via pullback from G_{ab} in order to eliminate those that have nontrivial restriction to the derived subgroup. This observation underlies

the definition of commutative character sheaves for non-commutative G . We state our second main result, Theorem 3.12, that pullback along the abelianization map defines an equivalence of categories $\mathcal{CCS}(G) \rightarrow \mathcal{CCS}(G_{\text{ab}})$. In Section 3.3, we use Galois cohomology to describe the relationship between $G(k)^*$ and $G_{\text{ab}}(k)^*$. We also compute the automorphism groups in $\mathcal{CCS}(G)$. In Section 5 we use Theorem 3.12 to geometrize types for certain supercuspidal representations of p -adic groups, in a sense made precise in Theorem 5.2. As preparation for the proof, we review some facts about the Heisenberg–Weil representation and its geometrization, in Section 5.2. Then, in Section 5.3, we review Yu’s theory of types and his study of smooth integral models. These elements are pulled together in Section 5.4, where the proof Theorem 5.2 is given.

Acknowledgements

We are extremely grateful to Loren Spice for explaining Yu’s types for supercuspidal representations and Masoud Kamgarpour for helpful conversations. We also thank Will Sawin and the anonymous referee, who pointed out errors in earlier versions of this paper and helped us clarify several points.

1. Recollections and definitions

Let G be a smooth group scheme over a finite field k ; that is, let G be a group scheme over k for which the structure morphism $G \rightarrow \text{Spec}(k)$ is smooth in the sense of [Gro67, Definition 17.3.1]. This implies $G \rightarrow \text{Spec}(k)$ is locally of finite type, but not that it is of finite type. We remark that the identity component G^0 of G is of finite type over k , while the component group scheme $\pi_0(G)$ of G is an étale group scheme over k , and both are smooth over k .

Let \bar{k} be a fixed algebraic closure of k . In general, we use overline to denote a separable closure of a field or the base change a geometric object to a scheme over a separable closure of a field. In this paper we use a common formalism for Weil sheaves, writing \mathcal{L} for the pair $(\bar{\mathcal{L}}, \phi)$, where $\bar{\mathcal{L}}$ is an ℓ -adic sheaf on $\bar{G} = G \otimes_k \bar{k}$ and where $\phi : \text{Fr}^* \bar{\mathcal{L}} \rightarrow \bar{\mathcal{L}}$ is an isomorphism of ℓ -adic sheaves. We also follow convention by referring to \mathcal{L} as a Weil sheaf on G , as in [Del80, Définition 1.1.10]. If \mathcal{L} and $\mathcal{L}' := (\bar{\mathcal{L}}', \phi')$ are Weil sheaves, we write $\alpha : \mathcal{L} \rightarrow \mathcal{L}'$ for a morphism $\alpha : \bar{\mathcal{L}} \rightarrow \bar{\mathcal{L}}'$ such that

$$\begin{array}{ccc} \text{Fr}^* \bar{\mathcal{L}} & \xrightarrow{\text{Fr}^* \alpha} & \text{Fr}^* \bar{\mathcal{L}}' \\ \phi \downarrow & & \downarrow \phi' \\ \bar{\mathcal{L}} & \xrightarrow{\alpha} & \bar{\mathcal{L}}' \end{array}$$

commutes. These conventions simplify notation considerably, but they were not employed in [CR18].

We write $m : G \times G \rightarrow G$ for the multiplication morphism, and $G(k)^*$ for $\text{Hom}(G(k), \bar{\mathbb{Q}}_\ell^\times)$. Define $\theta : G \times G \rightarrow G \times G$ by $\theta(g, h) = (h, g)$.

When G is commutative, a *character sheaf* on G is a triple $(\bar{\mathcal{L}}, \mu, \phi)$, where $\bar{\mathcal{L}}$ is a rank-one ℓ -adic local system on \bar{G} , $\mu : \bar{m}^* \bar{\mathcal{L}} \rightarrow \bar{\mathcal{L}} \boxtimes \bar{\mathcal{L}}$ is an isomorphism of sheaves on $\bar{G} \times \bar{G}$, and $\phi : \mathrm{Fr}_G^* \bar{\mathcal{L}} \rightarrow \bar{\mathcal{L}}$ is an isomorphism of sheaves on \bar{G} ; the triple $(\bar{\mathcal{L}}, \mu, \phi)$ is required to satisfy certain conditions [CR18, Definition 1.1]. Write $\mathcal{CS}(G)$ for the category of character sheaves on G .

Even when G is not commutative, the category $\mathcal{CS}(G)$, defined as in [CR18, Definition 1.1], still makes sense. In order to distinguish the resulting objects from the character sheaves of Lusztig, we will refer to the former as *linear character sheaves* (to evoke the one-dimensional character sheaves of [Kam09]).

2. Commutative character sheaves on commutative groups

We consider first the case that G is commutative, which we will later apply to the case of general smooth G . Let \mathcal{L} be a character sheaf on G . Since $m = m \circ \theta$ in this case, there is a canonical isomorphism $\xi : m^* \mathcal{L} \rightarrow \theta^* m^* \mathcal{L}$. There is also an isomorphism $\vartheta : \mathcal{L} \boxtimes \mathcal{L} \rightarrow \theta^*(\mathcal{L} \boxtimes \mathcal{L})$ given on stalks by the canonical map $\bar{\mathcal{L}}_g \otimes \bar{\mathcal{L}}_h \rightarrow \bar{\mathcal{L}}_h \otimes \bar{\mathcal{L}}_g$.

DEFINITION 2.1. — *A character sheaf (\mathcal{L}, μ) on a smooth commutative group scheme G is commutative if the following diagram of Weil sheaves on $G \times G$ commutes.*

$$\begin{array}{ccc} m^* \mathcal{L} & \xrightarrow{\mu} & \mathcal{L} \boxtimes \mathcal{L} \\ \xi \downarrow m = m \circ \theta & & \downarrow \vartheta \\ \theta^*(m^* \mathcal{L}) & \xrightarrow{\theta^* \mu} & \theta^*(\mathcal{L} \boxtimes \mathcal{L}) \end{array}$$

We write $\mathcal{CCS}(G)$ for the full subcategory of $\mathcal{CS}(G)$ consisting of commutative character sheaves.

In [CR18, Theorem 3.6], we showed that $\mathrm{Tr}_G : \mathcal{CS}(G)_{/\mathrm{iso}} \rightarrow G(k)^*$ is surjective and explicitly computed its kernel. In this section, we show that the corresponding map $\mathrm{Tr}_G : \mathcal{CCS}(G)_{/\mathrm{iso}} \rightarrow G(k)^*$ for commutative character sheaves is an isomorphism. We begin by reinterpreting Definition 2.1 in terms of cocycles.

Let G be a commutative étale group scheme over k . For a character sheaf \mathcal{L} on G , recall [CR18, § 2.3] that $S_G : \mathcal{CS}(G)_{/\mathrm{iso}} \rightarrow H^2(E_G^\bullet)$ is an isomorphism mapping $[\mathcal{L}]$ to $[\alpha \oplus \beta]$, where E_G^\bullet is the total space of the zeroth page of the Hochschild–Serre spectral sequence, $\alpha \in {}^\circ K^0(\mathcal{W}, {}^\circ K^2(\bar{G}, \bar{\mathbb{Q}}_\ell^\times))$ is obtained from μ and $\beta \in {}^\circ K^1(\mathcal{W}, {}^\circ K^1(\bar{G}, \bar{\mathbb{Q}}_\ell^\times))$ is obtained from ϕ .

Let $a \in Z^2(\bar{G}, \bar{\mathbb{Q}}_\ell^\times)$ correspond to α . We say that $[\alpha \oplus \beta] \in H^2(E_G^\bullet)$ is *symmetric* if $a(x, y) = a(y, x)$ for all $x, y \in \bar{G}$. This condition is well defined, since every coboundary in $B^2(\bar{G}, \bar{\mathbb{Q}}_\ell^\times)$ is symmetric. The connection between commutative character sheaves and symmetric classes is given in the following lemma.

LEMMA 2.2. — *Suppose G is a smooth commutative group scheme, and let \mathcal{L} be a character sheaf on G . Then \mathcal{L} is commutative if and only if $S_G(\mathcal{L})$ is symmetric.*

Proof. — The symmetry of $S_G(\mathcal{L})$ is a direct consequence of the commutativity of the diagram in Definition 2.1 after choosing bases for each stalk. \square

We may similarly define a symmetric class in $H^2(\bar{G}, \bar{\mathbb{Q}}_\ell^\times)$ to be one represented by a symmetric 2-cocycle. The following lemma will allow us to show that there are no invisible commutative character sheaves.

LEMMA 2.3. — *Let \bar{G} be a commutative group. Then the only symmetric class in $H^2(\bar{G}, \bar{\mathbb{Q}}_\ell^\times)$ is the trivial class.*

Proof. — By the universal coefficient theorem,

$$0 \rightarrow \mathrm{Ext}_{\mathbb{Z}}^1(H_{n-1}(\bar{G}, \mathbb{Z}), \bar{\mathbb{Q}}_\ell^\times) \rightarrow H^n(\bar{G}, \bar{\mathbb{Q}}_\ell^\times) \rightarrow \mathrm{Hom}(H_n(\bar{G}, \mathbb{Z}), \bar{\mathbb{Q}}_\ell^\times) \rightarrow 0$$

is exact for all $n > 0$. When $n = 2$, using the fact that \bar{G} is commutative, we have that $H_1(\bar{G}, \mathbb{Z}) \cong \bar{G}$ and that $H_2(\bar{G}, \mathbb{Z}) \cong \wedge^2 \bar{G}$. We get

$$0 \rightarrow \mathrm{Ext}_{\mathbb{Z}}^1(\bar{G}, \bar{\mathbb{Q}}_\ell^\times) \rightarrow H^2(\bar{G}, \bar{\mathbb{Q}}_\ell^\times) \rightarrow \mathrm{Hom}(\wedge^2 \bar{G}, \bar{\mathbb{Q}}_\ell^\times) \rightarrow 0.$$

The map $H^2(\bar{G}, \bar{\mathbb{Q}}_\ell^\times) \rightarrow \mathrm{Hom}(\wedge^2 \bar{G}, \bar{\mathbb{Q}}_\ell^\times)$ maps a 2-cocycle f to the alternating function

$$(x, y) \mapsto \frac{f(x, y)}{f(y, x)}.$$

Thus the cohomology classes represented by symmetric cocycles are precisely those in the image of $\mathrm{Ext}_{\mathbb{Z}}^1(\bar{G}, \bar{\mathbb{Q}}_\ell^\times)$. But $\mathrm{Ext}_{\mathbb{Z}}^1(-, \bar{\mathbb{Q}}_\ell^\times)$ vanishes because $\bar{\mathbb{Q}}_\ell^\times$ is divisible. \square

LEMMA 2.4. — *If G is a connected commutative algebraic group over k then every character sheaf on G is commutative.*

Proof. — Suppose $S_G(\mathcal{L}) = [\alpha \oplus \beta] \in H^2(E_G^\bullet)$. We can use étale descent to see that pullback by the Lang isogeny defines an equivalence of categories between local systems on G and $G(k)$ -equivariant local systems on G . Thus every character sheaf \mathcal{L} on G arises through the Lang isogeny, together with a character $G(k) \rightarrow \bar{\mathbb{Q}}_\ell^\times$. Pushing forward the Lang isogeny along this character defines an extension of \bar{G} by $\bar{\mathbb{Q}}_\ell^\times$ whose class is fixed by Frobenius; let $a \in Z^2(\bar{G}, \bar{\mathbb{Q}}_\ell^\times)$ be a representative 2-cocycle. Then a corresponds to the $\alpha \in {}^\circ K^0(\mathcal{W}, {}^\circ K^2(\bar{G}, \bar{\mathbb{Q}}_\ell^\times))$, above. Since the covering group of the Lang isogeny is $G(k)$, which is commutative, the class of this extension satisfies $a(x, y) = a(y, x)$ for all $x, y \in \bar{G}$. This shows that $S_G(\mathcal{L})$ is symmetric. It follows from Lemma 2.2 that \mathcal{L} is a commutative character sheaf. \square

THEOREM 2.5. — *If G is a smooth commutative group scheme over k then $\mathrm{Tr}_G : \mathcal{CCS}(G)_{/iso} \rightarrow G(k)^*$ is an isomorphism.*

Proof. — Suppose first that G is étale. Consider the isomorphism of short exact sequences

$$\begin{array}{ccccccc} 0 & \longrightarrow & \ker \mathrm{Tr}_G & \longrightarrow & \mathcal{CS}(G)_{/iso} & \xrightarrow{\mathrm{Tr}_G} & G(k)^* \longrightarrow 0 \\ & & \downarrow & & \downarrow S_G & & \downarrow \\ 0 & \longrightarrow & H^0(\mathcal{W}, H^2(\bar{G}, \bar{\mathbb{Q}}_\ell^\times)) & \longrightarrow & H^2(E_G^\bullet) & \longrightarrow & H^1(\mathcal{W}, H^1(\bar{G}, \bar{\mathbb{Q}}_\ell^\times)) \longrightarrow 0 \end{array}$$

from [CR18, Proposition 2.7].

Suppose that \mathcal{L} is a commutative character sheaf with $t_{\mathcal{L}} = 1$, and set $[\alpha, \beta] = S_G([\mathcal{L}])$. Then $S_G([\mathcal{L}])$ is in the image of $H^2(\bar{G}, \bar{\mathbb{Q}}_{\ell}^{\times})^{\mathcal{W}}$, so is cohomologous to $[\alpha', 0]$. Since α is symmetric and coboundaries are symmetric, α' is symmetric as well. So by Lemma 2.3, α' is cohomologically trivial, and thus $[\mathcal{L}]$ is trivial as well.

To see that Tr_G is still surjective on $\mathcal{CCS}(G)_{/\mathrm{iso}}$, note that the character sheaf constructed in the proof of [CR18, Proposition 2.6] has trivial α , and is thus commutative.

For general smooth commutative group schemes, we use Lemma 2.4 and the snake lemma, as in the proof of [CR18, Theorem 3.6]. \square

Remark 2.6. — Since $H^0(\mathcal{W}, H^2(\bar{G}, \bar{\mathbb{Q}}_{\ell}^{\times}))$ is not necessarily trivial [CR18, Example 2.10], the functor $\mathcal{CCS}(G) \rightarrow \mathcal{CS}(G)$ is not necessarily essentially surjective. Indeed, the invisible character sheaves [CR18, Definition 2.8] defined in our previous paper are precisely those non-commutative character sheaves with trivial trace of Frobenius.

3. Commutative character sheaves on non-commutative groups

We now consider the case of a smooth group scheme without the commutativity assumption. Our approach is to relate linear character sheaves on G to character sheaves on its abelianization $G_{\mathrm{ab}} = G/G_{\mathrm{der}}$, where G_{der} is defined by [Ber70, Définition 7.2.2] and the quotient G_{ab} is an instance of [Gab70, Section 7.2.2]. While every character $\chi \in G(k)^*$ vanishes on $G(k)_{\mathrm{der}}$, it may not vanish on $G_{\mathrm{der}}(k)$. For example, if k has odd characteristic then there are nontrivial characters $\mathrm{PGL}_2(k) \rightarrow \bar{\mathbb{Q}}_{\ell}^{\times}$ vanishing on $\mathrm{PGL}_2(k)_{\mathrm{der}} = \mathrm{PSL}_2(k)$ (see Section 4). In passing to G_{ab} , we may only hope to geometrize characters that vanish on all of $G_{\mathrm{der}}(k)$.

We begin this section with the main definition in this paper – the category $\mathcal{CCS}(G)$ of commutative character sheaves, Definition 3.2. This definition is delicate and somewhat technical, but it is vindicated in Theorem 3.10 which shows that $\mathcal{CCS}(G)$ is equivalent to the category of commutative character sheaves on the abelianization G_{ab} of G . To prove Theorem 3.10 we use descent theory in Section 3.2, in the process giving insight into Definition 3.2. Section 3 concludes with Theorem 3.12, showing that the dictionary from $\mathcal{CCS}(G)$ to characters of $G(k)$ in fact encompasses every character vanishing on $G_{\mathrm{der}}(k)$.

3.1. Main definition

Recall from Section 1 that we refer to objects in category $\mathcal{CS}(G)$, defined as in [CR18, Definition 1.1], as linear character sheaves when G is smooth but not necessarily commutative. We define the following category to track the trivialization on the derived subgroup; commutative character sheaves will then be defined as a subcategory.

DEFINITION 3.1. — Let $\mathcal{CS}_{\text{ab}}(G)$ denote the category of triples $(\mathcal{L}, \mu, \beta)$ where $(\mathcal{L}, \mu) \in \mathcal{CS}(G)$ and $\beta : \mathcal{L}|_{G_{\text{der}}} \rightarrow (\bar{\mathbb{Q}}_{\ell})_{G_{\text{der}}}$ is an isomorphism of Weil local systems on G_{der} with the constant sheaf on G_{der} . A morphism $(\mathcal{L}, \mu, \beta) \rightarrow (\mathcal{L}', \mu', \beta')$ is a morphism $\alpha : (\mathcal{L}, \mu) \rightarrow (\mathcal{L}', \mu')$ in $\mathcal{CS}(G)$ such that $\beta = \beta' \circ \alpha|_{G_{\text{der}}}$.

The reason for tracking β is that it determines an isomorphism $\gamma : m^* \mathcal{L} \rightarrow \theta^* m^* \mathcal{L}$ which will replace the ξ of Definition 2.1, as follows. Let $i : G \rightarrow G$ be inversion and $c : G \times G \rightarrow G_{\text{der}}$ be the commutator map, defined by $c(x, y) = xyx^{-1}y^{-1}$. Both are morphisms of k -schemes. Set $m' = i \circ m \circ \theta$ and let $j_{\text{der}} : G_{\text{der}} \rightarrow G$ be inclusion; then $j_{\text{der}} \circ c = m \circ (m \times m')$. Then $\beta : \mathcal{L}|_{G_{\text{der}}} \rightarrow (\bar{\mathbb{Q}}_{\ell})_{G_{\text{der}}}$ determines the isomorphism $\gamma' : m^* \mathcal{L} \otimes \theta^* m^* i^* \mathcal{L} \rightarrow (\bar{\mathbb{Q}}_{\ell})_{G \times G}$ by the diagram of isomorphisms below.

$$(3.1) \quad \begin{array}{ccc} c^*(\mathcal{L}|_{G_{\text{der}}}) & \xrightarrow{c^*(\beta)} & c^*\left((\bar{\mathbb{Q}}_{\ell})_{G_{\text{der}}}\right) \\ \parallel & & \parallel \\ c^*j_{\text{der}}^* \mathcal{L} & & (\bar{\mathbb{Q}}_{\ell})_{G \times G} \\ \downarrow j_{\text{der}} \circ c = m \circ (m \times m') & & \uparrow \gamma' \\ (m \times m')^* m^* \mathcal{L} & & m^* \mathcal{L} \otimes \theta^* m^* i^* \mathcal{L} \\ \downarrow (m \times m')^*(\mu) & & \uparrow m' = i \circ m \circ \theta \\ (m \times m')^*(\mathcal{L} \boxtimes \mathcal{L}) & \equiv & m^* \mathcal{L} \otimes (m')^* \mathcal{L} \end{array}$$

In the diagram above, the arrows labeled with equations come from canonical isomorphisms of functors on Weil sheaves derived from the equations; so, for example, the middle left isomorphism comes from $(m \times m')^* m^* \cong c^* j_{\text{der}}^*$ since $j_{\text{der}} \circ c = m \circ (m \times m')$. Using the monoidal structure of the category of Weil local systems on $G \times G$, the isomorphism $\gamma' : m^* \mathcal{L} \otimes \theta^* m^* i^* \mathcal{L} \rightarrow (\bar{\mathbb{Q}}_{\ell})_{G \times G}$ defines an isomorphism

$$m^* \mathcal{L} \rightarrow (\theta^* m^* i^* \mathcal{L})^{\vee}.$$

Applying the canonical isomorphisms $(\theta^* m^* i^* \mathcal{L})^{\vee} \cong \theta^* m^* i^*(\mathcal{L}^{\vee})$ and $i^*(\mathcal{L}^{\vee}) \cong \mathcal{L}$, this map provides the promised isomorphism

$$\gamma : m^* \mathcal{L} \longrightarrow \theta^* m^* \mathcal{L}.$$

DEFINITION 3.2. — The category $\mathcal{CCS}(G)$ of commutative character sheaves on G is the full subcategory of $\mathcal{CS}_{\text{ab}}(G)$ consisting of triples $(\mathcal{L}, \mu, \beta)$ such that the following diagram of Weil sheaves on $G \times G$ commutes:

$$\begin{array}{ccc} m^* \mathcal{L} & \xrightarrow{\mu} & \mathcal{L} \boxtimes \mathcal{L} \\ \gamma \downarrow & & \downarrow \vartheta \\ \theta^*(m^* \mathcal{L}) & \xrightarrow{\theta^* \mu} & \theta^*(\mathcal{L} \boxtimes \mathcal{L}). \end{array}$$

Here $\gamma : m^* \mathcal{L} \rightarrow \theta^* m^* \mathcal{L}$ is the isomorphism built from $\beta : \mathcal{L}|_{G_{\text{der}}} \rightarrow (\bar{\mathbb{Q}}_{\ell})_{G_{\text{der}}}$ as above.

3.2. Descent

In this section we give an equivalence of categories between $\mathcal{CS}(G_{\text{ab}})$ and $\mathcal{CS}_{\text{ab}}(G)$ and use it to describe the pullback functor $q^* : \mathcal{CS}(G_{\text{ab}}) \rightarrow \mathcal{CS}(G)$ in terms of the forgetful functor $\mathcal{CS}_{\text{ab}}(G) \rightarrow \mathcal{CS}(G)$, where $q : G \rightarrow G_{\text{ab}}$ is the abelianization quotient with kernel G_{der} . But first, in order to study commutative character sheaves, we need some auxiliary categories.

3.2.1. Equivariant Weil local systems

Let $\text{Loc}(G)$ be the category of Weil local systems on G . Let $\text{Loc}_{\text{der}}(G)$ be the category of G_{der} -equivariant Weil local systems on G , whose definition we now recall. Let $n : G_{\text{der}} \times G \rightarrow G$ be the restriction of $m : G \times G \rightarrow G$ to $G_{\text{der}} \times G$, let $p : G_{\text{der}} \times G \rightarrow G$ be projection to the second component, and let $s : G \rightarrow G_{\text{der}} \times G$ be given by $s(g) = (1, g)$. Then the quotient $q : G \rightarrow G_{\text{ab}}$ is a regular epimorphism of smooth group schemes with kernel pair (n, p) .

$$G_{\text{der}} \times G \begin{array}{c} \xrightarrow{n} \\ \xrightarrow{p} \end{array} G \xrightarrow{q} G_{\text{ab}}$$

Consider the morphisms

$$G_{\text{der}} \times G_{\text{der}} \times G \begin{array}{c} \xrightarrow{b_1, b_2, b_3} \\ \xrightarrow{\quad \quad \quad} \end{array} G_{\text{der}} \times G \begin{array}{c} \xrightarrow{n} \\ \xrightarrow{p} \end{array} G$$

defined by

$$\begin{aligned} b_1(h_1, h_2, g) &= (h_1 h_2, g) \\ b_2(h_1, h_2, g) &= (h_1, h_2 g) \\ b_3(h_1, h_2, g) &= (h_2, g). \end{aligned}$$

Note that

$$\begin{aligned} (3.2) \quad n \circ b_1 &= n \circ b_2 \\ n \circ b_3 &= p \circ b_2 \\ p \circ b_1 &= p \circ b_3. \end{aligned}$$

A G_{der} -equivariant Weil local system on G is a Weil local system \mathcal{L} on G together with an isomorphism

$$\nu : n^* \mathcal{L} \rightarrow p^* \mathcal{L}$$

of Weil local systems on $G_{\text{der}} \times G$ such that

$$(3.3) \quad s^*(\nu) = \text{id}_{\mathcal{L}}$$

and such that the following diagram of isomorphisms of Weil local systems on $G_{\text{der}} \times G_{\text{der}} \times G$ commutes.

$$(3.4) \quad \begin{array}{ccccc} & b_2^* n^* \mathcal{L} & \xrightarrow{b_2^*(\nu)} & b_2^* p^* \mathcal{L} & \\ & \swarrow n \circ b_1 = n \circ b_2 & & \searrow p \circ b_2 = n \circ b_3 & \\ b_1^* n^* \mathcal{L} & & & & b_3^* n^* \mathcal{L} \\ & \searrow b_1^*(\nu) & & \swarrow b_3^*(\nu) & \\ & b_1^* p^* \mathcal{L} & \xleftarrow{p \circ b_3 = p \circ b_1} & b_3^* p^* \mathcal{L} & \end{array}$$

Morphisms of H -equivariant Weil local systems $(\mathcal{L}, \nu) \rightarrow (\mathcal{L}', \nu')$ are morphisms of Weil local systems $\alpha : \mathcal{L} \rightarrow \mathcal{L}'$ for which the diagram

$$(3.5) \quad \begin{array}{ccc} n^* \mathcal{L} & \xrightarrow{n^*(\alpha)} & n^* \mathcal{L}' \\ \nu \downarrow & & \downarrow \nu' \\ p^* \mathcal{L} & \xrightarrow{p^*(\alpha)} & p^* \mathcal{L}' \end{array}$$

commutes. This defines $\text{Loc}_{\text{der}}(G)$, the category of G_{der} -equivariant Weil local systems on G . The reader will recognize this notion as the Weil local system version of equivariant sheaves for the action n of G_{der} on G , as can be found, for example, in [BL94, 0.2].

Let $\text{Loc}(G_{\text{ab}})$ be the category of Weil local systems on G_{ab} . If $\mathcal{L}_{\text{ab}} \in \text{Loc}(G_{\text{ab}})$ then $q^* \mathcal{L}_{\text{ab}} \in \text{Loc}(G)$ comes equipped with a canonical isomorphism $\nu(\mathcal{L}_{\text{ab}}) : n^* \mathcal{L} \rightarrow p^* \mathcal{L}$ defined by the following diagram of isomorphisms.

$$\begin{array}{ccc} n^* \mathcal{L} & \xrightarrow{\nu(\mathcal{L}_{\text{ab}})} & p^* \mathcal{L} \\ \parallel & & \parallel \\ n^*(q^* \mathcal{L}_{\text{ab}}) & \xrightarrow{q \circ n = q \circ p} & p^*(q^* \mathcal{L}_{\text{ab}}) \end{array}$$

Then $(q^* \mathcal{L}_{\text{ab}}, \nu(\mathcal{L}_{\text{ab}}))$ satisfies (3.3) and (3.4), so $(q^* \mathcal{L}_{\text{ab}}, \nu(\mathcal{L}_{\text{ab}})) \in \text{Loc}_{\text{der}}(G)$. Moreover, if $\alpha_{\text{ab}} : \mathcal{L}_{\text{ab}} \rightarrow \mathcal{L}'_{\text{ab}}$ is a morphism in $\text{Loc}(G_{\text{ab}})$ then $q^*(\alpha_{\text{ab}})$ satisfies the condition in (3.5), so $q^*(\alpha_{\text{ab}})$ is a morphism in $\text{Loc}_{\text{der}}(G)$. This defines the functor

$$L : \text{Loc}(G_{\text{ab}}) \rightarrow \text{Loc}_{\text{der}}(G)$$

LEMMA 3.3. — Suppose G is a smooth group scheme. The functor $L : \text{Loc}(G_{\text{ab}}) \rightarrow \text{Loc}_{\text{der}}(G)$ is an equivalence.

Proof. — The quotient $q : G \rightarrow G_{\text{ab}}$ is an G_{der} -torsor in the fppf topology by [Gab70, Theorem 3.2], and thus a G_{der} -torsor in the fpqc topology. The lemma is now a result from descent theory, arguing as in [FGI⁺05, Theorem 4.46] for example. \square

3.2.2. Equivariant linear character sheaves

With reference to Section 3.2.1, we define a G_{der} -equivariant linear character sheaf on G to be a triple (\mathcal{L}, μ, ν) , where (\mathcal{L}, μ) is a linear character sheaf and (\mathcal{L}, ν) is an

G_{der} -equivariant Weil local system. A morphism of G_{der} -equivariant linear character sheaves $(\mathcal{L}, \mu, \nu) \rightarrow (\mathcal{L}', \mu', \nu')$ is a morphism of G_{der} -equivariant Weil local systems $\alpha : \mathcal{L} \rightarrow \mathcal{L}'$ which is also a morphism of linear character sheaves; Let $\mathcal{CS}_{\text{der}}(G)$ be the category of G_{der} -equivariant linear character sheaves on G .

Consider the functor

$$q^* : \mathcal{CS}(G_{\text{ab}}) \rightarrow \mathcal{CS}(G)$$

given on objects by $(\mathcal{L}_{\text{ab}}, \mu_{\text{ab}}) \mapsto (q^*\mathcal{L}_{\text{ab}}, (q^2)^*\mu_{\text{ab}})$; this is an instance of [CR18, Lemma 1.4]. To see that $(q^*\mathcal{L}_{\text{ab}}, (q^2)^*\mu_{\text{ab}})$ is indeed a linear character sheaf on G , verify [CR18, CS.3]. Now set $L(\mathcal{L}_{\text{ab}}) = (\mathcal{L}, \nu)$, where $L : \text{Loc}(G_{\text{ab}}) \rightarrow \text{Loc}_{\text{der}}(G)$ is the comparison functor above, so $\mathcal{L} = q^*\mathcal{L}_{\text{ab}}$ and $\nu = \nu(\mathcal{L}_{\text{ab}})$. Then (\mathcal{L}, μ, ν) is an object in $\mathcal{CS}_{\text{der}}(G)$. If $\alpha_{\text{ab}} : (\mathcal{L}_{\text{ab}}, \mu_{\text{ab}}) \rightarrow (\mathcal{L}'_{\text{ab}}, \mu'_{\text{ab}})$ is a morphism in $\mathcal{CS}(G_{\text{ab}})$, then $q^*(\alpha_{\text{ab}}) : (\mathcal{L}, \mu) \rightarrow (\mathcal{L}', \mu')$ satisfies [CR18, CS4], so $\alpha = q^*(\alpha_{\text{ab}})$ is a morphism in $\mathcal{CS}(G)$. These observations define the comparison functor

$$q_{\text{ab}}^* : \mathcal{CS}(G_{\text{ab}}) \rightarrow \mathcal{CS}_{\text{der}}(G)$$

and also show that the functor $q^* : \mathcal{CS}(G_{\text{ab}}) \rightarrow \mathcal{CS}(G)$ factors according to the following commuting diagram of functors

$$(3.6) \quad \begin{array}{ccc} \mathcal{CS}(G) & \xleftarrow{q^*} & \mathcal{CS}(G_{\text{ab}}) \\ \text{forget} \uparrow & \swarrow q_{\text{ab}}^* & \\ \mathcal{CS}_{\text{der}}(G) & & \end{array}$$

The definition of $q_{\text{ab}}^* : \mathcal{CS}(G_{\text{ab}}) \rightarrow \mathcal{CS}_{\text{der}}(G)$ will be revisited in the proof of Proposition 3.5.

Set $G^2 = G \times G$, so $G_{\text{der}}^2 = G_{\text{der}} \times G_{\text{der}}$ and $G_{\text{ab}}^2 = G_{\text{ab}} \times G_{\text{ab}}$. Likewise define $n^2 : G_{\text{der}}^2 \times G^2 \rightarrow G^2$ and $p^2 : G_{\text{der}}^2 \times G \rightarrow G$.

LEMMA 3.4. — *If (\mathcal{L}, μ, ν) is a G_{der} -equivariant linear character sheaf on G then $\mu : m^*\mathcal{L} \rightarrow \mathcal{L} \boxtimes \mathcal{L}$ is a morphism of G_{der}^2 -equivariant Weil local systems on G^2 , as defined in Section 3.2.1.*

Proof. — Define

$$\begin{aligned} d : G_{\text{der}} \times G_{\text{der}} \times G \times G &\rightarrow G_{\text{der}} \times G \times G_{\text{der}} \times G \\ (h_1, h_2, g_1, g_2) &\mapsto (h_1, g_1, h_2, g_2) \\ n_2 : G_{\text{der}} \times G \times G_{\text{der}} \times G &\rightarrow G \times G \\ (h_1, g_1, h_2, g_2) &\mapsto (h_1 g_1, h_2 g_2) \\ p_2 : G_{\text{der}} \times G \times G_{\text{der}} \times G &\rightarrow G \times G \\ (h_1, g_1, h_2, g_2) &\mapsto (g_1, g_2). \end{aligned}$$

The following diagram defines the isomorphisms needed to see that both $m^*\mathcal{L}$ and $\mathcal{L} \boxtimes \mathcal{L}$ are G_{der}^2 -equivariant Weil local systems.

$$\begin{array}{ccc}
 n_2^*(m^*\mathcal{L}) & \dashrightarrow & p_2^*(m^*\mathcal{L}) \\
 n_2^*(\mu) \downarrow & & \downarrow p_2^*(\mu) \\
 n_2^*(\mathcal{L} \boxtimes \mathcal{L}) & \dashrightarrow & p_2^*(\mathcal{L} \boxtimes \mathcal{L}) \\
 n_2 = n^2 \circ d \downarrow & & \downarrow p_2 = p^2 \circ d \\
 d^*(n^*\mathcal{L} \boxtimes n^*\mathcal{L}) & \xrightarrow{d^*(\nu \boxtimes \nu)} & d^*(p^*\mathcal{L} \boxtimes p^*\mathcal{L})
 \end{array}$$

The dashed arrows both satisfy (3.3) and (3.4) as they apply here. This diagram also shows that $\mu : m^*\mathcal{L} \rightarrow \mathcal{L} \boxtimes \mathcal{L}$ is a morphism of G_{der}^2 -equivariant local systems, since it satisfies (3.5) as it applies here. \square

PROPOSITION 3.5. — *Suppose G is a smooth group scheme. Then pullback along $q : G \rightarrow G_{\text{ab}}$ defines an equivalence $\mathcal{CS}(G_{\text{ab}}) \rightarrow \mathcal{CS}_{\text{der}}(G)$.*

Proof. — Let $L^2 : \text{Loc}(G_{\text{ab}}^2) \rightarrow \text{Loc}_{\text{der}}(G^2)$ be the comparison functor for the quotient $q^2 : G^2 \rightarrow G_{\text{ab}}^2$. Then L^2 is also an equivalence by Lemma 3.3. Moreover, using Lemma 3.4, we may rewrite the comparison functor q_{ab}^* on objects by

$$\begin{aligned}
 \mathcal{CS}(G_{\text{ab}}) &\rightarrow \mathcal{CS}_{\text{der}}(G) \\
 (\mathcal{L}_{\text{ab}}, \mu_{\text{ab}}) &\mapsto (L(\mathcal{L}_{\text{ab}}), L^2(\mu_{\text{ab}})).
 \end{aligned}$$

and on morphisms by $\alpha \mapsto L(\alpha)$. The proposition now follows from the fact that both L and L^2 are equivalences. \square

3.2.3. Rigidification

We may now relate $\mathcal{CS}_{\text{ab}}(G)$ to $\mathcal{CS}(G_{\text{ab}})$.

PROPOSITION 3.6. — *The categories $\mathcal{CS}_{\text{ab}}(G)$ and $\mathcal{CS}(G_{\text{ab}})$ are equivalent.*

Proof. — In light of Proposition 3.5, it suffices to exhibit an equivalence between $\mathcal{CS}_{\text{der}}(G)$ and $\mathcal{CS}_{\text{ab}}(G)$.

We begin by defining a functor $\mathcal{CS}_{\text{ab}}(G) \rightarrow \mathcal{CS}_{\text{der}}(G)$. Let $j_{\text{der}} : G_{\text{der}} \rightarrow G$ be the kernel of $q : G \rightarrow G_{\text{ab}}$. Define $k : G_{\text{der}} \times G \rightarrow G \times G$ by $k(h, g) = (j_{\text{der}}(h), g)$. Then for $(\mathcal{L}, \mu, \nu) \in \mathcal{CS}_{\text{der}}(G)$, define $\beta : \mathcal{L}|_{G_{\text{der}}} \rightarrow (\bar{\mathbb{Q}}_{\ell})_{G_{\text{der}}}$ by the following diagram.

$$(3.7) \quad \begin{array}{ccc}
 & n^*\mathcal{L} \xrightarrow{\nu} p^*\mathcal{L} & \\
 \swarrow \scriptstyle m \circ k = n & & \searrow \\
 k^*m^*\mathcal{L} & & (\bar{\mathbb{Q}}_{\ell})_{G_{\text{der}}} \boxtimes \mathcal{L} \\
 \searrow \scriptstyle k^*(\mu) & & \nearrow \scriptstyle \beta \boxtimes \text{id}_{\mathcal{L}} \\
 & k^*(\mathcal{L} \boxtimes \mathcal{L}) = \mathcal{L}|_{G_{\text{der}}} \boxtimes \mathcal{L} &
 \end{array}$$

Then $\beta = r_{\text{der}} \circ j^*(\nu)$, where $r_{\text{der}} : (\mathcal{L}_1)_{G_{\text{der}}} \rightarrow (\bar{\mathcal{Q}}_\ell)_{G_{\text{der}}}$ is the isomorphism of constant local systems determined by the rigidification $r : \mathcal{L}_1 \rightarrow \bar{\mathcal{Q}}_\ell$ of \mathcal{L} determined by μ (see [CR18, Remark 1.11]) so that

$$(3.8) \quad \begin{array}{ccc} \mathcal{L}_1 & \xrightarrow{r} & \bar{\mathcal{Q}}_\ell \\ \downarrow \mu_{(1,1)} & & \parallel \\ \mathcal{L}_1 \otimes \mathcal{L}_1 & \xrightarrow{r \otimes r} & \bar{\mathcal{Q}}_\ell \otimes \bar{\mathcal{Q}}_\ell \end{array}$$

commutes where, as usual, $=$ denotes a canonical isomorphism. To show that $(\mathcal{L}, \mu, \beta) \in \mathcal{CS}_{\text{ab}}(G)$, we must see that β is a morphism in $\mathcal{CS}(G_{\text{der}})$ by showing that the following diagram commutes in $\text{Loc}(G_{\text{der}}^2)$.

$$(3.9) \quad \begin{array}{ccc} m_{\text{der}}^* \mathcal{L}|_{G_{\text{der}}} & \xrightarrow{m_{\text{der}}^* \beta} & (\bar{\mathcal{Q}}_\ell)_{G_{\text{der}}^2} \\ \parallel & & \parallel \\ \mathcal{L}|_{G_{\text{der}}} \boxtimes \mathcal{L}|_{G_{\text{der}}} & \xrightarrow{\beta \boxtimes \beta} & (\bar{\mathcal{Q}}_\ell)_{G_{\text{der}}} \boxtimes (\bar{\mathcal{Q}}_\ell)_{G_{\text{der}}} \end{array}$$

By Proposition 3.5, $\mathcal{L} = q^* \mathcal{L}_{\text{ab}}$, for some $(\mathcal{L}_{\text{ab}}, \mu_{\text{ab}}) \in \mathcal{CS}(G_{\text{ab}})$. Thus, $\mathcal{L}|_{G_{\text{der}}} = (\mathcal{L}_1)_{G_{\text{der}}}$ and the diagram above becomes a diagram of constant Weil local systems on G_{der}^2 . We can therefore test whether diagram commutes by evaluating all local systems at $(1, 1) \in G_{\text{der}}^2$. Doing this recovers (3.8) from (3.9), so (3.8) commutes. After confirming that morphisms that commute with μ and β also commute with μ and ν , this concludes the definition of the functor $\mathcal{CS}_{\text{ab}}(G) \rightarrow \mathcal{CS}_{\text{der}}(G)$.

The adjoint functor $\mathcal{CS}_{\text{der}}(G) \rightarrow \mathcal{CS}_{\text{ab}}(G)$ is given by a similar strategy: given $(\mathcal{L}, \mu, \beta) \in \mathcal{CS}_{\text{der}}(G)$, define (\mathcal{L}, μ, ν) again using (3.7). Then verify that (\mathcal{L}, ν) satisfies (3.4) as it applies here, so that $(\mathcal{L}, \nu) \in \text{Loc}_{\text{der}}(G)$. After confirming that morphisms that commute with μ and ν also commute with μ and β , this concludes the definition of the functor $\mathcal{CS}_{\text{der}}(G) \rightarrow \mathcal{CS}_{\text{ab}}(G)$. From these constructions, it is now clear that $\mathcal{CS}_{\text{der}}(G) \rightarrow \mathcal{CS}_{\text{ab}}(G)$ is an equivalence. \square

COROLLARY 3.7. — *If G is a smooth group scheme and $(\mathcal{L}, \mu) \in \mathcal{CS}(G)$, then the restriction of (\mathcal{L}, μ) to G_{der} is trivial if and only if $(\mathcal{L}, \mu) \cong q^*(\mathcal{L}_{\text{ab}}, \mu_{\text{ab}})$ in $\mathcal{CS}(G)$, for some $(\mathcal{L}_{\text{ab}}, \mu_{\text{ab}}) \in \mathcal{CS}(G_{\text{ab}})$.*

Proof. — Notation as in the proof of Proposition 3.5. Consider the following diagram.

$$\begin{array}{ccccc} \mathcal{CS}(G_{\text{der}}) & \xleftarrow{j_{\text{der}}^*} & \mathcal{CS}(G) & \xleftarrow{q^*} & \mathcal{CS}(G_{\text{ab}}) \\ & & \uparrow \text{forget} & \swarrow q_{\text{ab}}^* & \\ & & \mathcal{CS}_{\text{ab}}(G) & & \end{array}$$

Now, suppose $(\mathcal{L}, \mu) \in \mathcal{CS}(G)$ and there is an isomorphism $\beta : \mathcal{L}|_{G_{\text{der}}} \rightarrow (\bar{\mathcal{Q}}_\ell)_{G_{\text{der}}}$ so that $(\mathcal{L}, \mu, \beta) \in \mathcal{CS}_{\text{ab}}(G)$. By Proposition 3.5, there is some $(\mathcal{L}_{\text{ab}}, \mu_{\text{ab}}) \in \mathcal{CS}(G_{\text{ab}})$ with $(\mathcal{L}, \mu, \beta) \cong q_{\text{ab}}^*(\mathcal{L}_{\text{ab}}, \mu_{\text{ab}})$. Applying the forgetful functor $\mathcal{CS}_{G_{\text{der}}}(G) \rightarrow \mathcal{CS}(G)$ to this isomorphism, it follows that $(\mathcal{L}, \mu) \cong q^*(\mathcal{L}_{\text{ab}}, \mu_{\text{ab}})$ in $\mathcal{CS}(G)$, as desired.

Conversely, suppose $(\mathcal{L}, \mu) \in \mathcal{CS}(G)$ and $(\mathcal{L}, \mu) \cong q^*(\mathcal{L}_{\text{ab}}, \mu_{\text{ab}})$ in $\mathcal{CS}(G)$. Then

$$j_{\text{der}}^*(\mathcal{L}, \mu) \cong j_{\text{der}}^* q^*(\mathcal{L}_{\text{ab}}, \mu_{\text{ab}})$$

in $\mathcal{CS}(G_{\text{der}})$. Since $q \circ j_{\text{der}} = 1$, it follows that the restriction of (\mathcal{L}, μ) to G_{der} is trivial. \square

We may interpret this corollary as measuring how far q^* is from being essentially surjective. The next result shows that it is also not full. Let C denote the cokernel of the natural map

$$\text{Hom}(\pi_0(\bar{G})_{\text{Fr}}, \bar{\mathbb{Q}}_{\ell}^{\times}) \rightarrow \text{Hom}(\pi_0(\bar{G}_{\text{der}})_{\text{Fr}}, \bar{\mathbb{Q}}_{\ell}^{\times}),$$

where $\pi_0(\bar{G})_{\text{Fr}}$ denotes the coinvariants of the action of Frobenius on the component group of \bar{G}

COROLLARY 3.8. — *If G is a smooth group scheme and (\mathcal{L}, μ) is a linear character sheaf on G with trivial restriction to G_{der} , then the set of isomorphism classes of objects in $\mathcal{CS}(G_{\text{ab}})$ mapping to (\mathcal{L}, μ) under q^* is a principal homogeneous space for C .*

Proof. — By Proposition 3.5, it suffices to find the set of isomorphism classes in $\mathcal{CS}_{\text{ab}}(G)$ mapping to (\mathcal{L}, μ) under the forgetful functor. By the previous corollary this set is nonempty. If $(\mathcal{L}, \mu, \beta)$ and $(\mathcal{L}, \mu, \beta')$ both map to (\mathcal{L}, μ) , then $\beta' \circ \beta^{-1}$ is an automorphism of the constant sheaf on G_{der} . Conversely, if φ is an automorphism of $(\mathbb{Q}_{\ell})_{G_{\text{der}}}$ and $(\mathcal{L}, \mu, \beta) \in \mathcal{CS}_{\text{ab}}(G)$ then $(\mathcal{L}, \mu, \varphi \circ \beta) \in \mathcal{CS}_{\text{ab}}(G)$. As in [CR18, Theorem 3.9], the automorphism group is isomorphic to $\text{Hom}(\pi_0(\bar{G}_{\text{der}})_{\text{Fr}}, \bar{\mathbb{Q}}_{\ell}^{\times})$. Finally, we note that any automorphism α of $(\mathcal{L}, \mu) \in \mathcal{CS}(G)$ defines an isomorphism $(\mathcal{L}, \mu, \beta \circ \alpha|_{G_{\text{der}}}) \rightarrow (\mathcal{L}, \mu, \beta)$. Applying the analogue of [CR18, Theorem 3.9] again yields the desired result. \square

3.3. Objects and maps in commutative character sheaves

We are now in a position to prove that commutative character sheaves on G match perfectly with commutative character sheaves on G_{ab} . We start with a method that will allow us to situate the diagram in Definition 3.2 within $\mathcal{CS}_{\text{ab}}(G^2)$.

LEMMA 3.9. — *If $(\mathcal{L}, \mu, \beta) \in \mathcal{CS}_{\text{ab}}(G)$ then $\mu : m^*\mathcal{L} \rightarrow \mathcal{L} \boxtimes \mathcal{L}$, $\gamma : m^*\mathcal{L} \rightarrow \theta^*(m^*\mathcal{L})$ and $\vartheta : \mathcal{L} \boxtimes \mathcal{L} \rightarrow \theta^*(\mathcal{L} \boxtimes \mathcal{L})$ are morphisms in $\mathcal{CS}_{\text{ab}}(G^2)$.*

Proof. — Define $m^2 : G^2 \times G^2 \rightarrow G^2$ by $m^2(g_1, g_2, g'_1, g'_2) = (g_1 g'_1, g_2 g'_2)$. Also define $p_i^2 : G^2 \times G^2 \rightarrow G^2$ by $p_i^2(g_1, g_2, g'_1, g'_2) = (g_i, g'_i)$. First we show that $m^*\mathcal{L}$ is an object in $\mathcal{CS}(G^2)$ by equipping it with an isomorphism $\mu_m^2 : (m^2)^*(m^*\mathcal{L}) \rightarrow m^*\mathcal{L} \boxtimes m^*\mathcal{L}$ defined by the diagram below.

$$\begin{array}{ccc} (m^2)^*(m^*\mathcal{L}) & \xrightarrow{\mu_m^2} & m^*\mathcal{L} \boxtimes m^*\mathcal{L} \\ \downarrow (m^2)^*\mu & & \parallel \\ (m^2)^*(\mathcal{L} \boxtimes \mathcal{L}) & \xlongequal{\quad} (m^2)^*(p_1)^*\mathcal{L} \otimes (m^2)^*(p_2)^*\mathcal{L} \longrightarrow (p_1^2)^*m^*\mathcal{L} \otimes (p_2^2)^*m^*\mathcal{L} \end{array}$$

The pair $(m^*\mathcal{L}, \mu_m^2)$ satisfies the conditions appearing in [CR18, Definition 1.1]. The restriction of $m^*\mathcal{L}$ to $G_{\text{der}}^2 = G_{\text{der}} \times G_{\text{der}}$ is canonically isomorphic to $(\bar{\mathbb{Q}}_\ell)_{G_{\text{der}}^2}$ by

$$\begin{array}{ccc} (m^*\mathcal{L})|_{G_{\text{der}}^2} & \xrightarrow{\beta_m^2} & (\bar{\mathbb{Q}}_\ell)_{G_{\text{der}}^2} \\ \downarrow \mu|_{G_{\text{der}}^2} & & \uparrow \beta \boxtimes \beta \\ (\mathcal{L} \boxtimes \mathcal{L})|_{G_{\text{der}}^2} & \xlongequal{\quad} & (\mathcal{L}|_{G_{\text{der}}}) \boxtimes (\mathcal{L}|_{G_{\text{der}}}). \end{array}$$

This shows that $(m^*\mathcal{L}, \mu_m^2, \beta_m^2) \in \mathcal{CS}_{\text{ab}}(G^2)$. Similar work defines $(\mathcal{L} \boxtimes \mathcal{L}, \mu_{\boxtimes}^2, \beta_{\boxtimes}^2) \in \mathcal{CS}_{\text{ab}}(G^2)$. By construction, $\mu : m^*\mathcal{L} \rightarrow \mathcal{L} \boxtimes \mathcal{L}$ is a morphism in $\mathcal{CS}_{\text{ab}}(G^2)$. Similar work shows that $\gamma : m^*\mathcal{L} \rightarrow \theta^*(m^*\mathcal{L})$ and $\vartheta : \mathcal{L} \boxtimes \mathcal{L} \rightarrow \theta^*(\mathcal{L} \boxtimes \mathcal{L})$ are also morphisms in $\mathcal{CS}_{\text{ab}}(G^2)$. \square

Suppose G is commutative, so $G_{\text{der}} = 1$. Suppose $(\mathcal{L}, \mu, \beta)$ is an object in $\mathcal{CS}_{\text{ab}}(G)$. Then $\beta : \mathcal{L}_1 \rightarrow \bar{\mathbb{Q}}_\ell$ is an isomorphism in $\mathcal{CS}(1)$, which is unique by [CR18, Theorem 3.9]. Tracing through the construction of $\gamma : m^*\mathcal{L} \rightarrow \theta^*m^*\mathcal{L}$ from $\beta : \mathcal{L}_1 \rightarrow \bar{\mathbb{Q}}_\ell$, we find that $\gamma : m^*\mathcal{L} \rightarrow \theta^*m^*\mathcal{L}$ is the canonical isomorphism coming from the equation $m = m \circ \theta$. Thus, when G is commutative, Definition 3.2 agrees with Definition 2.1. The next result generalizes this observation.

THEOREM 3.10. — *Pull-back along the abelianization $q : G \rightarrow G_{\text{ab}}$ defines an equivalence of categories*

$$\mathcal{CCS}(G_{\text{ab}}) \rightarrow \mathcal{CCS}(G).$$

Proof. — By definition, $\mathcal{CCS}(G)$ is a full subcategory of $\mathcal{CS}_{\text{ab}}(G)$; likewise, $\mathcal{CCS}(G_{\text{ab}})$ is a full subcategory of $\mathcal{CS}_{\text{ab}}(G_{\text{ab}})$. We have just seen that $\mathcal{CS}_{\text{ab}}(G_{\text{ab}})$ is equivalent to $\mathcal{CS}(G_{\text{ab}})$. By Proposition 3.5, pullback along the abelianization $q : G \rightarrow G_{\text{ab}}$ induces an equivalence $q_{\text{ab}}^* : \mathcal{CS}(G_{\text{ab}}) \rightarrow \mathcal{CS}_{\text{ab}}(G)$. Thus, the functor $\mathcal{CS}_{\text{ab}}(G_{\text{ab}}) \rightarrow \mathcal{CS}_{\text{ab}}(G)$ induced by pullback along q is an equivalence. The functor $\mathcal{CCS}(G_{\text{ab}}) \rightarrow \mathcal{CCS}(G)$ under consideration is the restriction of $\mathcal{CS}_{\text{ab}}(G_{\text{ab}}) \rightarrow \mathcal{CS}_{\text{ab}}(G)$ to the subcategory $\mathcal{CCS}(G_{\text{ab}})$.

$$\begin{array}{ccc} & \mathcal{CS}(G_{\text{ab}}) & \\ & \nwarrow q_{\text{ab}}^* & \uparrow \text{equiv.} \\ \mathcal{CS}_{\text{ab}}(G) & \longleftarrow & \mathcal{CS}_{\text{ab}}(G_{\text{ab}}) \\ \uparrow & & \uparrow \\ \mathcal{CCS}(G) & \longleftarrow & \mathcal{CCS}(G_{\text{ab}}) \end{array}$$

To prove the theorem, it is now sufficient to show that $\mathcal{CCS}(G_{\text{ab}}) \rightarrow \mathcal{CCS}(G)$ is essentially surjective. Suppose $(\mathcal{L}, \nu, \beta) \in \mathcal{CCS}(G)$. Then $(\mathcal{L}, \nu, \beta) \in \mathcal{CS}_{\text{ab}}(G)$. Let $(\mathcal{L}_{\text{ab}}, \mu_{\text{ab}}) \in \mathcal{CS}(G_{\text{ab}})$ be given by the equivalences above. Let $\xi : m_{\text{ab}}^*\mathcal{L}_{\text{ab}} \rightarrow \theta^*m_{\text{ab}}^*\mathcal{L}_{\text{ab}}$ be the isomorphism attached to $(\mathcal{L}_{\text{ab}}, \mu_{\text{ab}}) \in \mathcal{CS}(G_{\text{ab}})$ as in Section 2. Let $\gamma : m^*\mathcal{L} \rightarrow \theta^*m^*\mathcal{L}$ be the isomorphism attached to $\beta : \mathcal{L}|_{G_{\text{der}}} \rightarrow (\bar{\mathbb{Q}}_\ell)_{G_{\text{der}}}$ as in Section 3.1.

By Lemma 3.9, the diagrams below are in $\mathcal{CS}(G_{\text{ab}})$ (right) and $\mathcal{CS}_{\text{ab}}(G)$ (left).

$$\begin{array}{ccc}
 m^*\mathcal{L} & \xrightarrow{\mu} & \mathcal{L} \boxtimes \mathcal{L} \\
 \gamma \downarrow & & \downarrow \vartheta \\
 \theta^*(m^*\mathcal{L}) & \xrightarrow{\theta^*\mu} & \theta^*(\mathcal{L} \boxtimes \mathcal{L})
 \end{array}
 \quad \xleftarrow{(q^2)^*_{\text{ab}}} \quad
 \begin{array}{ccc}
 m^*_{\text{ab}}\mathcal{L}_{\text{ab}} & \xrightarrow{\mu_{\text{ab}}} & \mathcal{L}_{\text{ab}} \boxtimes \mathcal{L}_{\text{ab}} \\
 \xi \downarrow & & \downarrow \vartheta \\
 \theta^*(m^*_{\text{ab}}\mathcal{L}_{\text{ab}}) & \xrightarrow{\theta^*\mu_{\text{ab}}} & \theta^*(\mathcal{L}_{\text{ab}} \boxtimes \mathcal{L}_{\text{ab}})
 \end{array}$$

The diagram on the left is the result of applying the functor $(q^2)^*_{\text{ab}}$ to the one on the right; in particular $\gamma = (q^2)^*_{\text{ab}}\xi$. Since $(q^2)^*_{\text{ab}}$ is an equivalence by Proposition 3.5, it follows that the diagram in Definition 3.2 commutes if and only if the diagram in Definition 2.1 commutes. In other words, $(\mathcal{L}, \mu, \beta) \in \mathcal{CCS}(G)$ if and only if $(\mathcal{L}_{\text{ab}}, \mu_{\text{ab}}) \in \mathcal{CCS}(G_{\text{ab}})$. \square

We may use Theorem 3.10 to give a description of the morphisms and the isomorphism classes of objects in $\mathcal{CCS}(G)$.

COROLLARY 3.11. — *The category $\mathcal{CCS}(G)$ is monoidal and there is a canonical isomorphism*

$$\mathcal{CCS}(G)_{/\text{iso}} \cong \text{Hom}\left(G_{\text{ab}}(k), \bar{\mathbb{Q}}_{\ell}^{\times}\right).$$

Every map in $\mathcal{CCS}(G)$ is either trivial or an isomorphism, and the automorphism group of any object in $\mathcal{CCS}(G)$ is canonically isomorphic to $\text{Hom}(\pi_0(\bar{G}_{\text{ab}})_{\text{Fr}}, \bar{\mathbb{Q}}_{\ell}^{\times})$.

Proof. — The first claim follows from Theorems 2.5 and 3.10. Let us write $(\mathcal{L}, \mu, \beta) \mapsto (\mathcal{L}_{\text{ab}}, \mu_{\text{ab}})$ to indicate the equivalence appearing in Theorem 3.10; then

$$\text{Aut}_{\mathcal{CCS}(G)}(\mathcal{L}, \mu, \beta) = \text{Aut}_{\mathcal{CCS}(G_{\text{ab}})}(\mathcal{L}_{\text{ab}}, \mu_{\text{ab}}).$$

By [CR18, Theorem 3.9], $\text{Aut}_{\mathcal{CCS}(G_{\text{ab}})}(\mathcal{L}_{\text{ab}}, \mu_{\text{ab}}) = \text{Hom}(\pi_0(\bar{G}_{\text{ab}})_{\text{Fr}}, \bar{\mathbb{Q}}_{\ell}^{\times})$. \square

3.4. Geometrizing characters trivial on the derived subgroup

Corollary 3.11 shows that commutative character sheaves on G provide a natural geometrization of characters of $G_{\text{ab}}(k)$. In Theorem 3.12 we take this one small step further by exploring the relation between characters of $G(k)$ and objects in $\mathcal{CCS}(G)$.

THEOREM 3.12. — *The trace of Frobenius $\text{Tr} : \mathcal{CCS}(G)_{/\text{iso}} \rightarrow G(k)^*$ fits into the following diagram,*

$$\begin{array}{ccccccc}
 \mathcal{CCS}(G_{\text{ab}})_{/\text{iso}} & \xrightarrow{\cong} & \mathcal{CCS}(G)_{/\text{iso}} \\
 \cong \downarrow \text{Tr} & & \downarrow \text{Tr} \\
 1 \longrightarrow \Delta_G^* & \longrightarrow & G_{\text{ab}}(k)^* & \longrightarrow & G(k)^* & \longrightarrow & G_{\text{der}}(k)^* \longrightarrow 1,
 \end{array}$$

where Δ_G is the image of the connecting homomorphism $G_{\text{ab}}(k) \rightarrow H^1(k, G_{\text{der}})$. Thus the category $\mathcal{CCS}(G)$ geometrizes characters of $G(k)$ in the following sense: for every group homomorphism $\chi : G(k) \rightarrow \bar{\mathbb{Q}}_{\ell}^{\times}$ that vanishes on $G_{\text{der}}(k)$, there is an object $(\mathcal{L}, \mu, \beta)$ in $\mathcal{CCS}(G)$ such that $t_{\mathcal{L}} = \chi$. While the geometrization of χ is not unique, the isomorphism classes of possibilities are enumerated by Δ_G^* .

Proof. — By the definition of Δ_G , we have a short exact sequence

$$1 \rightarrow G_{\text{der}}(k) \rightarrow G(k) \rightarrow G_{\text{ab}}(k) \rightarrow \Delta_G \rightarrow 1.$$

Applying $\text{Hom}(-, \bar{\mathbb{Q}}_\ell^\times)$, which is exact since $\bar{\mathbb{Q}}_\ell^\times$ is divisible, yields the bottom row in the statement.

By Theorem 3.10, the map $\mathcal{CCS}(G_{\text{ab}})_{/\text{iso}} \rightarrow \mathcal{CCS}(G)_{/\text{iso}}$ is an isomorphism. Moreover, since both $\mathcal{CCS}(G_{\text{ab}})_{/\text{iso}} \rightarrow \mathcal{CCS}(G)_{/\text{iso}}$ and $G_{\text{ab}}(k)^* \rightarrow G(k)^*$ are defined by pullback along q , the square in the statement of the theorem commutes. Finally, $\text{Tr} : \mathcal{CCS}(G_{\text{ab}})_{/\text{iso}} \rightarrow G_{\text{ab}}(k)^*$ is an isomorphism by Corollary 3.11. \square

4. Geometrizing Characters Nontrivial on the Derived Subgroup

If χ is a character of $G(k)$ vanishing on $G_{\text{der}}(k)$ then Theorem 3.12 shows that there is a *local system* \mathcal{L} such that $m^*\mathcal{L} \cong \mathcal{L} \boxtimes \mathcal{L}$ and $\text{Tr}(\mathcal{L}) = \chi$. In this section, we give methods for geometrizing characters that do not vanish on $G_{\text{der}}(k)$, though the results may not be local systems or conjugation-equivariant, a notion that we now define.

4.1. Conjugation equivariant Weil sheaves

A Weil sheaf \mathcal{L} is said to be *conjugation-equivariant* if $a^*\mathcal{L} \cong p_2^*\mathcal{L}$, where $a : G \times G \rightarrow G$ is conjugation $a(x, y) = xyx^{-1}$ and $p_2 : G \times G \rightarrow G$ is projection $p_2(x, y) = y$. If (\mathcal{L}, μ) is a linear character sheaf on G , then \mathcal{L} is conjugation-equivariant. Indeed, the isomorphism $\mu : m^*\mathcal{L} \rightarrow \mathcal{L} \boxtimes \mathcal{L}$ determines an isomorphism $a^*\mathcal{L} \rightarrow p_2^*\mathcal{L}$ of Weil sheaves on $G \times G$ by the following diagram.

$$\begin{array}{ccc}
 a^*\mathcal{L} & \xrightarrow{\hspace{2cm}} & p_2^*\mathcal{L} \\
 \left\| \begin{array}{c} a = m \circ (m \times (i \circ p_1)) \end{array} \right. & & \left\| \begin{array}{c} \mathcal{L} \otimes \mathcal{L}^\vee \cong (\bar{\mathbb{Q}}_\ell)_G \end{array} \right. \\
 (m \times (i \circ p_1))^* m^*\mathcal{L} & & p_1^*(\mathcal{L} \otimes \mathcal{L}^\vee) \otimes p_2^*\mathcal{L} \\
 \left\| \begin{array}{c} (m \times (i \circ p_1))^* \mu \downarrow \end{array} \right. & & \left\| \right. \\
 (m \times (i \circ p_1))^*(\mathcal{L} \boxtimes \mathcal{L}) & & p_1^*\mathcal{L} \otimes p_2^*\mathcal{L} \otimes p_1^*\mathcal{L}^\vee \\
 \left\| \begin{array}{c} m = p_1 \circ (m \times (i \circ p_1)) \end{array} \right. & \left\| \begin{array}{c} i \circ p_1 = p_2 \circ (m \times (i \circ p_1)) \end{array} \right. & \\
 m^*\mathcal{L} \otimes (i \circ p_1)^*\mathcal{L} & \xrightarrow[\mu \otimes \text{id}]{} & (\mathcal{L} \boxtimes \mathcal{L}) \otimes (i \circ p_1)^*\mathcal{L}
 \end{array}$$

Although every linear character sheaf on G is a conjugation-equivariant Weil local system on G , the converse is certainly not true.

4.2. Lusztig's character sheaves

For connected reductive groups G over k , Lusztig's character sheaves are simple perverse sheaves $\mathcal{IC}(C, \mathcal{L})$ on G that come equipped with an isomorphism $a^* \mathcal{IC}(C, \mathcal{L}) \rightarrow p_2^* \mathcal{IC}(C, \mathcal{L})$ in the triangulated category $D_c^b(G \times G; \bar{\mathbb{Q}}_\ell)$. Consequently, when $\mathcal{IC}(C, \mathcal{L})$ is also a Weil sheaf complex, its trace of Frobenius function $t_{\mathcal{IC}(C, \mathcal{L})} : G(k) \rightarrow \bar{\mathbb{Q}}_\ell$ is a class function, called the characteristic function of $\mathcal{IC}(C, \mathcal{L})$. By [Lus86, Corollary 25.7], Weil character sheaves on G determine a basis for the vector space of class functions on $G(k)$; in particular, any character of any representation of $G(k)$ can be expressed as a $\bar{\mathbb{Q}}_\ell$ -linear combination of characteristic functions of Weil character sheaves on G . Let us use the notation $D_{c,G}^b(G; \bar{\mathbb{Q}}_\ell)$ for the category of conjugation-equivariant objects in $D_c^b(G; \bar{\mathbb{Q}}_\ell)$, as defined in [BL94] for example, and $\mathrm{KD}_{c,G}^b(G; \bar{\mathbb{Q}}_\ell)$ for the Grothendieck group of $D_{c,G}^b(G; \bar{\mathbb{Q}}_\ell)$; we set $\mathrm{K}_{\bar{\mathbb{Q}}_\ell} D_{c,G}^b(G; \bar{\mathbb{Q}}_\ell) := \mathrm{KD}_{c,G}^b(G; \bar{\mathbb{Q}}_\ell) \otimes_{\mathbb{Z}} \bar{\mathbb{Q}}_\ell$. Thus, Lusztig's result shows that for every representation ρ of $G(k)$ there is some $\mathcal{F} \in \mathrm{K}_{\bar{\mathbb{Q}}_\ell} D_{c,G}^b(G; \bar{\mathbb{Q}}_\ell)$ such that $\mathrm{Tr} \rho = t_{\mathcal{F}}$. By [Lus04, Theorem 21.21], this result extends to all reductive groups G over k without the hypothesis that G is connected.

When specialized to the case of characters χ of $G(k)$, Lusztig's theory of character sheaves geometrizes χ using $\mathrm{K}_{\bar{\mathbb{Q}}_\ell} D_{c,G}^b(G; \bar{\mathbb{Q}}_\ell)$; of course, the resulting geometrization is generally not a local system. For an example when this method can be used to geometrize a character of $G(k)$ which is not trivial on $G_{\mathrm{der}}(k)$, consider $G = \mathrm{SL}_2$ over \mathbf{F}_3 . Then $G_{\mathrm{der}} = G$, but $G(k)_{\mathrm{der}}$ is the subgroup of semisimple elements, isomorphic to the quaternion group Q_8 and of index 3. There is thus a nontrivial character $\chi : G(k) \rightarrow \mu_3 \subset \bar{\mathbb{Q}}_\ell^\times$. Lusztig's character sheaves are constructed by perverse extension of local systems on a certain stratification of G . In the case of SL_2 , there are five strata:

C_3	regular semisimple elements
C_2	regular unipotent elements
C_2^-	$\{-u : \text{regular unipotent } u\}$
$\{1\}$	$\{1\}$
$\{2\}$	$\{2\}$

Each Weil character sheaf on SL_2 over \mathbf{F}_3 is the perverse extension of one of the following local systems: the rank-1 constant sheaf $\mathbf{1}_{C_3}$ (resp. $\mathbf{1}_{C_2}$, $\mathbf{1}_{C_2^-}$, $\mathbf{1}_{\{1\}}$, $\mathbf{1}_{\{2\}}$) on C_3 (resp. C_2 , C_2^- , $\{1\}$, $\{2\}$), or the non-trivial rank-1 local system \mathcal{E}_{C_2} (resp. $\mathcal{E}_{C_2^-}$) on C_2^- (resp. $\mathcal{E}_{C_2^-}$) trivialized by the double cover of C_2 (resp. C_2^-). The following table describes these perverse sheaves by showing how they decompose in the Grothendieck group into standard sheaves on SL_2 , and also gives the values of the trace of Frobenius on the 7 conjugacy classes in $\mathrm{SL}_2(\mathbf{F}_3)$.

Perverse Sheaf	Standard Sheaves	Trace of Frobenius values at conjugacy classes						
		$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	$\begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}$	$\begin{pmatrix} 0 & 1 \\ 2 & 0 \end{pmatrix}$	$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$	$\begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}$	$\begin{pmatrix} 2 & 1 \\ 0 & 2 \end{pmatrix}$	$\begin{pmatrix} 2 & 2 \\ 0 & 2 \end{pmatrix}$
$\mathcal{IC}(\mathbf{1}_{C_3})$	$\mathbf{1}_{\mathrm{SL}_2}[3]$	-1	-1	-1	-1	-1	-1	-1
$\mathcal{IC}(\mathbf{1}_{C_2})$	$\mathbf{1}_{C_2}[2] \oplus \mathbf{1}_{\{1\}}[2]$	1	0	0	1	1	0	0
$\mathcal{IC}(\mathcal{E}_{C_2})$	$\mathcal{E}_{C_2}[2]$	0	0	0	$\sqrt{-3}$	$-\sqrt{-3}$	0	0
$\mathcal{IC}(\mathbf{1}_{C_2^-})$	$\mathbf{1}_{C_2^-}[2] \oplus \mathbf{1}_{\{2\}}[2]$	0	1	0	0	0	1	1
$\mathcal{IC}(\mathcal{E}_{C_2^-})$	$\mathcal{E}_{C_2^-}[2]$	0	0	0	0	0	$\sqrt{-3}$	$-\sqrt{-3}$
$\mathcal{IC}(\mathbf{1}_{\{1\}})$	$\mathbf{1}_{\{1\}}[0]$	1	0	0	0	0	0	0
$\mathcal{IC}(\mathbf{1}_{\{2\}})$	$\mathbf{1}_{\{2\}}[0]$	0	1	0	0	0	0	0

Using the trace of Frobenius values, we find that

$$\begin{aligned}
 -\mathcal{IC}(\mathbf{1}_{C_3}) - \frac{3}{2}\mathcal{IC}(\mathbf{1}_{C_2}) + \frac{1}{2}\mathcal{IC}(\mathcal{E}_{C_2}) - \frac{3}{2}\mathcal{IC}(\mathbf{1}_{C_2^-}) \\
 - \frac{1}{2}\mathcal{IC}(\mathcal{E}_{C_2^-}) + \frac{3}{2}\mathcal{IC}(\mathbf{1}_{\{1\}}) + \frac{3}{2}\mathcal{IC}(\mathbf{1}_{\{2\}})
 \end{aligned}$$

geometrizes χ . Note that all seven character sheaves appear in this geometrization, even though we start just with a character; in particular, the geometrization of the character χ given by this method is not a local system.

We remark that for $p > 3$ and $G = \mathrm{SL}_2$, we have $G_{\mathrm{der}}(k) = G(k)_{\mathrm{der}}$. In this case, therefore, any character of $G(k)$ may be geometrized by a local system using Theorem 3.12, so Lusztig's character sheaves are not required.

4.3. Pushforward along the Lang isogeny

For connected, commutative groups, the standard method for geometrizing a character χ is to use the Lang isogeny $L : G \rightarrow G$, defined by $L(y) = y^{-1} \mathrm{Fr}_G(y)$. The pushforward $L_!(\bar{\mathbb{Q}}_\ell)_G$ of the constant sheaf decomposes into a direct sum of rank 1 local systems according to characters of the automorphism group $G(k)$ of the cover L , and the local system \mathcal{L} associated to χ^{-1} will have $\mathrm{Tr}(\mathcal{L}) = \chi$.

If G is nonabelian or disconnected, a similar strategy sometimes succeeds in geometrizing characters $\chi : G(k) \rightarrow \bar{\mathbb{Q}}_\ell^\times$. Define $\tilde{L} : G \rightarrow G$ by $\tilde{L}(y) = \mathrm{Fr}_G(y)y^{-1}$. The fibers of L are right cosets of $G(k)$ and the fibers of \tilde{L} are left cosets. Suppose that

(L.1) $G(k) \subset \mathrm{im}(\tilde{L}) \cap \mathrm{im}(L)$ (which will hold if G is connected).

Then we may associate to each χ a function $\tilde{\chi} : G(k) \rightarrow \bar{\mathbb{Q}}_\ell^\times$ as follows. If $z \in G(k)$, write $z = \tilde{L}(y)$ and define $\tilde{\chi}(z) = \chi(L(y))$. Note that $\tilde{\chi}$ is well defined since changing y to ya for $a \in G(k)$ has the effect of conjugating $L(y)$ by a^{-1} , which has no effect on the value $\chi(L(y))$. In order to obtain a local system with trace χ , we must also assume that

(L.2) $\tilde{\chi}$ is also a character.

PROPOSITION 4.1. — *Let G be a smooth group scheme over k and assume (L.1) and (L.2). Let $\mathcal{L} = (L_!(\bar{\mathbb{Q}}_\ell)_G)_{\tilde{\chi}^{-1}}$. Then $\mathrm{Tr}(\mathcal{L}) = \chi$.*

Proof. — The stalk of \mathcal{L} at $x \in G(k)$ is

$$\mathcal{L}_x = \left\{ s : L^{-1}(x) \rightarrow \bar{\mathbb{Q}}_\ell : s(ay) = \tilde{\chi}^{-1}(a)s(y) \text{ for all } a \in G(k), y \in L^{-1}(x) \right\}.$$

For any $y \in L^{-1}(x)$, note that $\mathrm{Fr}_G^{-1}(y) = yx^{-1}$ and $\tilde{L}(y) = yxy^{-1} \in G(k)$. Frobenius acts on $s \in \mathcal{L}_x$ by

$$\left({}^{\mathrm{Fr}}s \right)(y) = s\left(\mathrm{Fr}_G^{-1}(y)\right) = \tilde{\chi}^{-1}\left(yx^{-1}y^{-1}\right)s(y) = \tilde{\chi}\left(\tilde{L}(y)\right)s(y) = \chi(x)s(y),$$

thus proving the claim. \square

Note that the resulting sheaf is a local system, but is not necessarily conjugation-equivariant because the Lang map is not a homomorphism when G is nonabelian.

This method of geometrization also applies to the example considered above. Let $G = \mathrm{SL}_2$ over \mathbf{F}_3 and let $\chi : G(k) \rightarrow \mu_3 \subset \bar{\mathbb{Q}}_\ell^\times$ be a non-trivial character. In this case, $\tilde{\chi} = \chi^{-1}$. Since G is connected and $\tilde{\chi}$ is a character, Proposition 4.1 produces a local system \mathcal{L} on G which geometrizes χ . This \mathcal{L} is not conjugation-equivariant.

The same method of geometrization applies to $G = \mathrm{PGL}_2$ when k has odd characteristic. In this case, $G_{\mathrm{der}} = G$, but $G(k)_{\mathrm{der}} = \mathrm{PSL}_2(k)$ has index 2 in $G(k)$. For the non-trivial character χ of $G(k)$, $\tilde{\chi} = \chi$, so again we may use Proposition 4.1 to geometrize χ as a local system.

5. Application to type theory for p -adic groups

We now show how to use Theorem 3.12 to geometrize Yu type data and how to geometrize types for supercuspidal representations of tamely ramified p -adic groups.

5.1. Quasicharacters of smooth group schemes over certain Henselian traits

Recall that R is the ring of integers of a local field with finite residue field k . The maximal ideal of R will be denoted by \mathfrak{m} . Let \underline{G} be a smooth group scheme over R . Here we shall use [BGA18] for the definition and fundamental properties of the Greenberg transform. Let G be the Greenberg transform of \underline{G} ; then G is a group scheme over k and there is a canonical isomorphism

$$G(k) = \underline{G}(R).$$

Let $\varphi : \underline{G}(R) \rightarrow \bar{\mathbb{Q}}_\ell^\times$ be a quasicharacter. By the continuity of φ there is some $r \in \mathbb{N}$ and a factorization

$$\begin{array}{ccc} \underline{G}(R) & \xrightarrow{\varphi} & \bar{\mathbb{Q}}_\ell^\times \\ & \searrow & \nearrow \varphi_r \\ & \underline{G}(R/\mathfrak{p}^{r+1}) & \end{array}$$

The least such $r \in \mathbb{N}$ is called the *level* of φ . For any $r \in \mathbb{N}$, define $R_r := R/\mathfrak{p}^{r+1}$ and set $G_r := \mathrm{Gr}_r^R(\underline{G})$, the Greenberg transform of $\underline{G} \times_{\mathrm{Spec}(R)} \mathrm{Spec}(R_r)$. Then G_r is a smooth group scheme over k and $G_r(k) = \underline{G}(R_r)$.

PROPOSITION 5.1. — *Let \underline{G} be a smooth group scheme over R . Let $\varphi : \underline{G}(R) \rightarrow \bar{\mathbb{Q}}_\ell^\times$ be a quasicharacter and let r be the level of φ . If $\varphi_r : G_r(k) \rightarrow \bar{\mathbb{Q}}_\ell^\times$ is trivial on $G_{r,\text{der}}(k)$ then there is a commutative character sheaf $\mathcal{L}_r \in \mathcal{CCS}(G_r)$ such that*

$$t_{\mathcal{L}_r} = \varphi_r.$$

Proof. — This is a direct consequence of Theorem 3.12. \square

With φ as above, note that if r' is any integer greater or equal to the level r of φ , then the pullback of \mathcal{L}_r along $G_{r'} \rightarrow G_r$ is a linear character sheaf on $G_{r'}$. Thus, $\mathcal{L}_{r'}$ is conjugation-equivariant in the sense of Section 4.1.

Recall that the full Greenberg transform $G := \text{Gr}^R(\underline{G})$ is a group scheme over k such that $G(k) = \underline{G}(R)$; it comes equipped with a morphism $G \rightarrow G_r$. The Weil sheaf on G obtained from \mathcal{L}_r by pullback along the morphism of group schemes $G \rightarrow G_r$ is a quasicharacter sheaf on G , in the sense of [CR18, Definition 4.2], such that

$$t_{\mathcal{L}} = \varphi.$$

5.2. Jacobi theory over finite fields

For use below, we recall some facts about the Heisenberg–Weil representation.

Let V be a finite-dimensional vector space over a finite field k equipped with a symplectic pairing $\langle \cdot, \cdot \rangle : V \times V \rightarrow Z$, where Z is a one-dimensional vector space over k . Let V^\sharp be the Heisenberg group determined by $(Z, \langle \cdot, \cdot \rangle)$ [GH07, § 1.1]. Let $\text{Sp}(V)$ be the symplectic group determined by the symplectic pairing $\langle \cdot, \cdot \rangle$; this group acts on V^\sharp . The group $\text{Sp}(V) \ltimes V^\sharp$ is called the Jacobi group. From the construction above, it is clear that the Jacobi group may be viewed as the k -points of an algebraic group over k ; we will refer to that algebraic group as the Jacobi group.

Let $\psi : Z \rightarrow \bar{\mathbb{Q}}_\ell^\times$ be an additive character and let ω_ψ be the Heisenberg representation on V^\sharp with central character ψ [GH07, § 1.1]. The Heisenberg representation determines a representation π_ψ of $\text{Sp}(V)$ with the same representation space as ω_ψ and with the defining property: for each $g \in \text{Sp}(V)$, $\pi_\psi(g)$ determines an isomorphism of representations $\omega_\psi^g \rightarrow \omega_\psi$. Let $W_\psi = \pi_\psi \ltimes \omega_\psi$ be the Heisenberg–Weil representation of the Jacobi group $\text{Sp}(V) \ltimes V^\sharp$ given by ω_ψ and π_ψ [GH07, § 2.2].

By [GH07, Theorem 3.2.2.1] (see also [GH11, Theorem 4.5]), there is a Frobenius-stable conjugation-equivariant perverse sheaf \mathcal{K}_ψ on $\text{Sp}(V) \ltimes V^\sharp$ such that

$$(5.1) \quad t_{\mathcal{K}_\psi} = \text{Tr}(W_\psi).$$

In particular, this geometrization uses $D_{c, \text{Sp}(V) \ltimes V^\sharp}^b(\text{Sp}(V) \ltimes V^\sharp; \bar{\mathbb{Q}}_\ell)$, as recalled in Section 4.2, to geometrize $\text{Tr}(W_\psi)$.

5.3. Yu’s types and integral models

A Yu type datum $({}^\circ K^i, {}^\circ \rho^0, \varphi^i, d)$ consists of the following:

- (Y0) a sequence of compact groups ${}^\circ K^0 \subseteq {}^\circ K^1 \subseteq \cdots \subseteq {}^\circ K^d = {}^\circ K$;
- (Y1) a continuous representation ${}^\circ \rho^0$ of ${}^\circ K^0$;

(Y2) quasicharacters $\varphi^i : {}^\circ K^i \rightarrow \mathbb{C}^\times$, for $i = 0, \dots, d$.

The representation ${}^\circ \rho^0$ and the quasicharacters $(\varphi^0, \dots, \varphi^d)$ enjoy certain properties which allow Yu to construct a sequence of types $({}^\circ K^i, {}^\circ \rho_i)$, for $i = 1, \dots, d$. In order to prepare for the construction of the geometric types of Theorem 5.2 we review some further detail here. In Table 5.1 we explain how to convert the constructions appearing in this section into the notation of [Yu01].

First, Yu introduces

(Y3) compact groups $J_i \subset {}^\circ K$, for $i = 0, \dots, d$, such that ${}^\circ K^i = J_0 \cdots J_i$ and, for $i = 0, \dots, d-1$, a natural action of ${}^\circ K^i$ on J_{i+1} defining the groups ${}^\circ K^i \ltimes J_{i+1}$.

$$(5.2) \quad \begin{array}{ccccccc} & & 1 & & & & \\ & & \downarrow & & & & \\ & & {}^\circ K^i \cap J_{i+1} & & & & \\ & & \downarrow & & & & \\ 1 & \longrightarrow & J_{i+1} & \longrightarrow & {}^\circ K^i \ltimes J_{i+1} & \xrightarrow{p_i} & {}^\circ K^i \longrightarrow 1 \\ & & & & \downarrow \pi_{i+1} & & \\ & & & & {}^\circ K^{i+1} & & \\ & & & & \downarrow & & \\ & & & & 1 & & \end{array}$$

Next, Yu defines a group homomorphism (in fact, a quotient) $J_{i+1} \rightarrow V_{i+1}$ where V_{i+1} is a finite abelian group, the latter also given the structure of a k -vector space. The vector space V_{i+1} is then equipped with a symplectic pairing $\langle \cdot, \cdot \rangle_{i+1} : V_{i+1} \times V_{i+1} \rightarrow Z_{i+1}$, where Z_{i+1} is a one-dimensional vector space over k , itself equipped with an additive character $\psi_{i+1} : Z_{i+1} \rightarrow \mathbb{C}^\times$. This, in turn, is used to define a map $J_{i+1} \rightarrow V_{i+1}^\sharp$, where V_{i+1}^\sharp is the Heisenberg group determined by V_{i+1} , Z_{i+1} , $\langle \cdot, \cdot \rangle_{i+1}$ and ψ_{i+1} , as in Section 5.2. In fact, the quotient $J_{i+1} \rightarrow V_{i+1}^\sharp$ factors through a quotient $J_{i+1} \rightarrow H_{i+1}$ and an isomorphism $j_{i+1} : H_{i+1} \rightarrow V_{i+1}^\sharp$, where H_{i+1} is a Heisenberg p -group in the sense of [Yu01]. Finally, Yu constructs a group homomorphism $f_{i+1} : {}^\circ K^i \rightarrow \mathrm{Sp}(V_{i+1})$ such that the pair (f_{i+1}, j_{i+1}) is a symplectic action of ${}^\circ K^i$ on H_{i+1} in the sense of [Yu01]. Taken together, this defines

(Y4) a group homomorphism $h_{i+1} : {}^\circ K^i \ltimes J_{i+1} \rightarrow \mathrm{Sp}(V_{i+1}) \ltimes V_{i+1}^\sharp$ making the following diagram commute.

$$\begin{array}{ccccccc} 1 & \longrightarrow & J_{i+1} & \longrightarrow & {}^\circ K^i \ltimes J_{i+1} & \xrightarrow{p_i} & {}^\circ K^i \longrightarrow 1 \\ & & \downarrow & & \downarrow h_{i+1} & & \downarrow f_i \\ 1 & \longrightarrow & V_{i+1}^\sharp & \longrightarrow & \mathrm{Sp}(V_{i+1}) \ltimes V_{i+1}^\sharp & \longrightarrow & \mathrm{Sp}(V_{i+1}) \longrightarrow 1 \end{array}$$

We can now recall how Yu uses all this to construct the types $({}^\circ K^i, {}^\circ \rho_i)$; see [Yu01, § 4, 15]. The representations ${}^\circ \rho_i$ are defined recursively. For the base case $i = 0$, set ${}^\circ \rho_0 := {}^\circ \rho^0 \otimes \varphi^0$; see (Y1) above. Now fix i . Let W_{i+1} be the Heisenberg–Weil

Table 5.1. Notation conversion chart.

this paper	Jiu-Kang Yu, <i>Construction of tame supercuspidal representations</i>	[Yu01]
${}^\circ K^0$	${}^\circ K^0 = G^0(F)_y$	[Yu01, § 15]
${}^\circ K^{i+1}$	${}^\circ K^{i+1} = ({}^\circ K^0) \vec{G}^{(i+1)}(F)_{y, (0, s_0, \dots, s_i)}$	[Yu01, § 15]
${}^\circ \rho^0$	${}^\circ \rho^0$	[Yu01, § 15]
${}^\circ \rho_{i+1}$	${}^\circ \rho_{i+1}$	[Yu01, § 15]
φ^i	$\phi_i _{{}^\circ K^i}$	[Yu01, § 3]
J_{i+1}	$J^{i+1} = (G^i, G^{i+1})(F)_{y, (r_i, s_i)}$	[Yu01, § 3]
V_{i+1}	$J^{i+1}/J_+^{i+1} = (G^i, G^{i+1})(F)_{y, (r_i, s_i)}/$ $(G^i, G^{i+1})(F)_{y, (r_i, s_i^+)}$	[Yu01, § 3]
V_{i+1}^\sharp	$(G^i, G^{i+1})(F)_{y, (r_i, s_i)}/\ker\left(\widehat{\phi_i}\Big _{(G^i, G^{i+1})(F)_{y, (r_i, s_i^+)}}\right)$	[Yu01, § 4]
Z_{i+1}	$\ker(V_{i+1}^\sharp \rightarrow V_{i+1})$	[Yu01, § 11]
(f_{j+1}, j_{i+1})	(f, j)	[Yu01, §11]
$\langle \ , \ \rangle_{i+1}$	$\langle \ , \ \rangle$	[Yu01, § 11]

representation of the Jacobi group $\mathrm{Sp}(V_{i+1}) \ltimes V_{i+1}^\sharp$, whose restriction to V_{i+1}^\sharp has central character ψ_{i+1} . Pull-back along h_{i+1} to form $h_{i+1}^*(W_{i+1})$, a representation of ${}^\circ K^i \ltimes J_{i+1}$. Write $\inf({}^\circ \rho_i)$ for the representation of ${}^\circ K^i \ltimes J_{i+1}$ obtained by pulling back ${}^\circ \rho_i$ along ${}^\circ K^i \ltimes J_{i+1} \rightarrow {}^\circ K^i$. Consider the representation

$$(5.3) \quad {}^\circ \rho^{i+1} := h_{i+1}^*(W_{i+1}) \otimes \inf({}^\circ \rho_i)$$

of ${}^\circ K^i \ltimes J_{i+1}$. By [Yu01], the representation ${}^\circ \rho^{i+1}$ of ${}^\circ K^i \ltimes J_{i+1}$ is trivial on ${}^\circ K^i \cap J_{i+1}$ so ${}^\circ \rho^{i+1}$ descends to ${}^\circ K^{i+1}$. Set ${}^\circ \rho_{i+1} = {}^\circ \rho^{i+1} \otimes \varphi^{i+1}$. This completes the recursive definition of the Yu $({}^\circ K^i, {}^\circ \rho_i)$ for $i = 0, \dots, d$. By [Yu15, Proposition 10.2] there is a sequence

$$\underline{G}^0 \rightarrow \underline{G}^1 \rightarrow \dots \rightarrow \underline{G}^d = \underline{G}$$

of morphisms of affine smooth group schemes of finite type over R such that, on R -points it gives the sequence ${}^\circ K^0 \subseteq {}^\circ K^1 \subseteq \dots \subseteq {}^\circ K^d$ above. Indeed, this is the main result of [Yu15].

As explained in [Yu15, § 10.4], there is morphism of affine smooth group schemes of finite type over R

$$\underline{J}^i \rightarrow \underline{G},$$

for each $i = 0, \dots, d$, such that $\underline{J}^i(R) = J_i$ as a subgroup of G and such that the image of the R -points under the multiplication map $\underline{J}^0 \times \dots \times \underline{J}^i \rightarrow \underline{G}$ is ${}^\circ K^i$, for $i = 0, \dots, d$. There is a natural action of \underline{G}^i on \underline{J}^{i+1} in the category of smooth affine group schemes over R so that the group scheme

$$\underline{G}^i \ltimes \underline{J}^{i+1} \quad \text{gives} \quad (\underline{G}^i \ltimes \underline{J}^{i+1})(R) = {}^\circ K^i \ltimes J_{i+1}.$$

Write \underline{J}_k^{i+1} for the special fibre $\underline{J}^{i+1} \times_{\mathrm{Spec}(R)} \mathrm{Spec}(k)$ of \underline{J}^{i+1} . The vector space V_{i+1} may be realized as the k -points on a variety V^{i+1} over k , where V^{i+1} , appears as a quotient $\underline{J}_k^{i+1} \rightarrow V^{i+1}$ of algebraic groups over k . Then the quotient $J_{i+1} \rightarrow V_{i+1}$ is realized as the composition

$$\underline{J}^{i+1}(R) \rightarrow \underline{J}^{i+1}(k) = \underline{J}_k^{i+1}(k) \rightarrow V^{i+1}(k) = V_{i+1}.$$

Likewise, the Heisenberg p -group H_{i+1} , appearing in Section 5.3, may be realized as a quotient of algebraic groups, and $\underline{J}_k^{i+1} \rightarrow H^{i+1}$ as the composition

$$\underline{J}^{i+1}(R) \rightarrow \underline{J}^{i+1}(k) = \underline{J}_k^{i+1}(k) \rightarrow H_k^{i+1}(k) = H_{i+1}.$$

Finally, the group homomorphism $f_i : J_0 \cdots J_i \rightarrow \mathrm{Sp}(V_{i+1})$ may be made geometric in much the same way. Writing \underline{G}_k^i for the special fibre $\underline{G}^i \times_{\mathrm{Spec}(R)} \mathrm{Spec}(k)$ of \underline{G}^i , and writing $\underline{G}_k^{i,\mathrm{red}}$ for the reductive quotient of \underline{G}_k^i , there is a quotient of algebraic groups $\underline{G}_k^{i,\mathrm{red}} \rightarrow W_k^{i+1}$ so that $f_i : J_0 \cdots J_i \rightarrow \mathrm{Sp}(V_{i+1})$ is realized as the composition

$$\underline{G}^i(R) \rightarrow \underline{G}^i(k) = \underline{G}_k^i(k) \rightarrow \underline{G}_k^{i,\mathrm{red}}(k) \rightarrow W_k^{i+1}(k) = \mathrm{Sp}(V_{i+1}).$$

With all this, we may revisit the quotients appearing in Section 5.3:

$$\begin{array}{ccccccc} 1 & \longrightarrow & \underline{J}^{i+1} & \longrightarrow & \underline{G}^i \ltimes \underline{J}^{i+1} & \longrightarrow & \underline{G}^i \longrightarrow 1 \\ & & \uparrow & & \uparrow & & \uparrow \\ 1 & \longrightarrow & \underline{J}_k^{i+1} & \longrightarrow & \underline{G}_k^i \ltimes \underline{J}_k^{i+1} & \longrightarrow & \underline{G}_k^i \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ 1 & \longrightarrow & V_{i+1}^\# & \longrightarrow & \mathrm{Sp}(V_{i+1}) \ltimes V_{i+1}^\# & \longrightarrow & \mathrm{Sp}(V_{i+1}) \longrightarrow 1, \end{array}$$

where the last two rows are now understood as forming a diagram in the category of algebraic groups over k . This realizes the Jacobi group $\mathrm{Sp}(V_{i+1}) \ltimes V_{i+1}^\#$ as a quotient of the special fibre of the smooth group scheme $\underline{G}^i \ltimes \underline{J}^{i+1}$ over R .

We may now revisit the ingredients in the construction of the representation ρ of $\underline{G}(R)$ along the lines indicated by Yu and recalled in Section 5.3.

- (M0) The compact groups ${}^\circ K^i$ appearing in (Y0) have been replaced by the smooth group schemes \underline{G}^i .
- (M1) The continuous representation ${}^\circ \rho^0$ of ${}^\circ K^0$ appearing in (Y1) may be interpreted as a representation of $\underline{G}^0(R)$ obtained by inflation along $\underline{G}^0(R) \rightarrow \underline{G}^0(k)$ from a representation ϱ_0 of $\underline{G}^0(k) = \underline{G}_k^0(k)$. In fact, ϱ_0 is itself obtained by pulling back a representation ϱ_0^{red} ; this is a consequence of [Yu15, § 10.5 (i)].
- (M2) The quasicharacters φ^i appearing in (Y2) are now quasicharacters of $\underline{G}^i(R)$, for $i = 0, \dots, d$. In fact, if r_i is the level of φ^i , as it appears in Section 5.1, then φ^i is obtained by pulling back a quasicharacter $\varphi_i := \varphi_{r_i}^i$ of the smooth group scheme $G_i := G_{r_i}^i := \mathrm{Gr}_{r_i}^R(\underline{G}^i)$ along $G^i \rightarrow G_i$.

(M3) Diagram (5.2) in (Y3) is now replaced by the following diagram of smooth group schemes over R .

$$(5.4) \quad \begin{array}{ccccccc} & & & 1 & & & \\ & & & \downarrow & & & \\ & & & \underline{G}^i \times_{\underline{G}} \underline{J}^{i+1} & & & \\ & & & \downarrow & & & \\ 1 & \longrightarrow & \underline{J}^{i+1} & \longrightarrow & \underline{G}^i \ltimes \underline{J}^{i+1} & \longrightarrow & \underline{G}^i \longrightarrow 1 \\ & & & & \downarrow & & \\ & & & & \underline{G}^{i+1} & & \\ & & & & \downarrow & & \\ & & & & 1 & & \end{array}$$

(M4) The representation $h_{i+1}^*(W_{i+1})$ appearing in (Y4) is now obtained by pulling back a representation along

$$(\underline{G}^i \ltimes \underline{J}^{i+1})(R) \rightarrow (\underline{G}^i \ltimes \underline{J}^{i+1})(k).$$

Let w_{i+1} be that representation of $(\underline{G}^i \ltimes \underline{J}^{i+1})(k) = (\underline{G}_k^i \ltimes \underline{J}_k^{i+1})(k)$. Then w_{i+1} is itself obtained by pulling back the representation W_{i+1} along the k -points of the quotient

$$\underline{G}_k^i \ltimes \underline{J}_k^{i+1} \rightarrow \mathrm{Sp}(V_{i+1}) \ltimes V_{i+1}^\sharp.$$

This brings us back to the point made in [Yu15, § 10.5] as quoted in the Introduction to this paper.

5.4. Geometrization of characters of certain types

We may now give the main result of Section 5, Theorem 5.2. Since Yu's theory refers to complex representations, and since our geometrization uses ℓ -adic sheaves, we grit our teeth and fix an isomorphism $\mathbb{C} \approx \bar{\mathbb{Q}}_\ell$.

As we recalled in Section 5.3, a Yu type datum consists of compact groups ${}^\circ K^i$, a representation ${}^\circ \rho^0$ of ${}^\circ K^0$ and quasicharacters φ^i of ${}^\circ K^i$, for $i = 0, \dots, d$; see (Y0), (Y1) and (Y2). In Section 5.3 we also saw how a Yu type datum determines smooth group schemes \underline{G}^i , a representation ϱ_0^{red} of the reductive quotient $(\underline{G}^0)_k^{\mathrm{red}}$ of the special fibre \underline{G}_k^0 of \underline{G}^0 , and quasicharacters φ_i of the group of k -rational points on the level- r_i Greenberg transform G_i of \underline{G}^i ; see (M0), (M1) and (M2). Theorem 5.2 places the following conditions on the Yu type datum:

- (H1) for each $i = 0, \dots, d$, either
- (a) G_i is reductive, or
 - (b) the quasicharacter φ_i of G_i is trivial on $G_{i, \mathrm{der}}(k)$.

Hypothesis **(H1a)** may be used to geometrize each quasicharacter φ_i inside the Grothendieck group $K_{\bar{\mathbb{Q}}} D_{c, G_i}^b(G_i; \bar{\mathbb{Q}}_\ell)$. Alternatively, Hypothesis **(H1b)** allows us to use Proposition 5.1 to geometrize each quasicharacter φ_i using local systems on G_i , or more precisely, using commutative character sheaves on G_i . Using Sections 4.1 and 4.2 we see that, in both cases, the resulting geometrization may be interpreted as an element in $K_{\bar{\mathbb{Q}}} D_{c, G_i}^b(G_i; \bar{\mathbb{Q}}_\ell)$. Note that when Hypothesis **(H1b)** applies, it provides a considerably simpler geometrization than when Hypothesis **(H1a)** applies.

We give examples illustrating Hypothesis **(H1b)**: first, an example of a quasicharacter that does not vanish on $G_{i, \text{der}}(k)$, and then a class of Yu data where it is always satisfied. Let $G = \text{SL}_2$ over \mathbb{Q}_2 . Then the hyperspecial maximal compact ${}^\circ K^0$ reduces to $\text{SL}_2(\mathbb{Z}/2^{d+1}\mathbb{Z})$ at depth d . The derived subgroup of $\text{SL}_2(\mathbb{Z}/2\mathbb{Z})$ has index 2, while it has index 4 in positive depth. There is therefore a character φ^0 of K of order 4 and depth 1 factoring through the quotient of $\text{SL}_2(\mathbb{Z}/4\mathbb{Z})$ by its derived subgroup. Taking the trivial representation for ${}^\circ \rho^0$ gives a datum not satisfying **(H1b)**. Note that this datum does not satisfy **(H1a)** either since the depth 1 Greenberg transform of the hyperspecial ${}^\circ K^0$ is not reductive.

In contrast, there is a broad class of representations that automatically satisfy **(H1b)**. Recall that a representation is *toral* ([Adl98] and [Kal19, § 6]) if it arises from a Yu datum of the form $(S \subset G, {}^\circ \rho^0 = 1, (\varphi^{d-1}, 1))$ where S is an elliptic maximal torus and φ^{d-1} is G -generic character of $S(F)$ of positive depth. Since S is abelian, it has trivial derived subgroup and thus **(H1b)** is satisfied. Toral representations include epipelagic representations [RY14] and the representations considered by Reeder in [Ree08].

THEOREM 5.2. — *Let $({}^\circ K^i, {}^\circ \rho^0, \varphi^i, d)$ be a Yu type datum that satisfies Hypotheses **(H1)**, and let ${}^\circ \rho_i$ be the representation of ${}^\circ K^i$ constructed from it in Section 5.3. Then for each $i = 0, \dots, d$ there is an element $\mathcal{F}_i \in K_{\bar{\mathbb{Q}}} D_{c, G^i}^b(G^i; \bar{\mathbb{Q}}_\ell)$ such that*

$$t_{\mathcal{F}_i} = \text{Tr}({}^\circ \rho_i).$$

Proof. — Recall that $G^i(k) = \underline{G}^i(R) = {}^\circ K^i$, canonically. Let r_i be the level of φ^i as defined in Section 5.1. Set $r = \max\{r_i \mid i = 0, \dots, d\}$.

By [Lus04, Theorem 21.21], there is $A \in K_{\bar{\mathbb{Q}}} D_{c, (\underline{G}^0)_k^{\text{red}}}^b((\underline{G}^0)_k^{\text{red}}; \bar{\mathbb{Q}}_\ell)$ such that

$$t_A = \text{Tr } \varrho_0^{\text{red}}.$$

Let $A^0 \in K_{\bar{\mathbb{Q}}} D_{c, (\underline{G}^0)_k}^b((\underline{G}^0)_k; \bar{\mathbb{Q}}_\ell)$ be obtained by pullback along the quotient $(\underline{G}^0)_k \rightarrow (\underline{G}^0)_k^{\text{red}}$. Then

$$t_{A^0} = \text{Tr } \varrho_0.$$

The special fibre $(\underline{G}^0)_k$ of the smooth group scheme \underline{G}^0 is itself a smooth group scheme, and may be identified with the Greenberg transform $Q^0 = \text{Gr}_0^R(\underline{G}^0)$ [CR18, § 4.3]. With $r \in \mathbb{N}$ as above, let A_r^0 be the equivariant Weil sheaf on the algebraic

group G_r^i obtained by pull-back from A^0 along the affine morphism $G_r^i \rightarrow Q^0$. Factor

$$(5.5) \quad \begin{array}{ccc} G^0(k) & \xrightarrow{\mathrm{Tr}(\circ \rho^0)} & \bar{\mathbb{Q}}_\ell \\ & \searrow & \nearrow \mathrm{Tr}(\circ \rho^0)_r \\ & G_r^0(k) & \end{array}$$

Observe that $\mathrm{Tr}(\circ \rho^0)_r$ may be recovered from A_r^0 :

$$t_{A_r^0} = \mathrm{Tr}(\circ \rho^0)_r$$

Consider the Jacobi group $\mathrm{Sp}(V_{i+1}) \ltimes V_{i+1}^\sharp$ and the Heisenberg–Weil representation W_{i+1} appearing in Section 5.3. Let \mathcal{K}^{i+1} be the conjugation equivariant Weil sheaf on the Jacobi group, recalled in Section 5.2, such that

$$t_{\mathcal{K}^{i+1}} = \mathrm{Tr}(W_{i+1}).$$

Recall from Section 5.3 that $\mathrm{Sp}(V_{i+1}) \ltimes V_{i+1}^\sharp$ is a quotient of the special fibre of the smooth group scheme $\underline{G}^i \ltimes \underline{J}^{i+1}$. Let \mathcal{K}_0^{i+1} be the Weil sheaf on the special fibre of $\underline{G}^i \ltimes \underline{J}^{i+1}$ obtained from W_{i+1} by pullback. Let \mathcal{K}_r^{i+1} be the equivariant Weil sheaf on $\mathrm{Gr}_r^R(\underline{G}^i \ltimes \underline{J}^{i+1})$ obtained from \mathcal{K}_0^{i+1} by pullback along the affine morphism $\mathrm{Gr}_r^R(\underline{G}^i \ltimes \underline{J}^{i+1}) \rightarrow \mathrm{Gr}_0^R(\underline{G}^i \ltimes \underline{J}^{i+1})$.

We now define $\mathcal{A}_r^i \in \mathbf{K}_{\bar{\mathbb{Q}}_\ell} \mathbf{D}_{c, G_r^i}^b$ for $i = 0, \dots, d$ recursively, following the construction of the representations $\circ \rho^i$, as reviewed in Section 5.3. First, set $\mathcal{A}_r^0 = A_r^0$ and note that (5.5) commutes with $\mathrm{Tr}(\circ \rho^0)_r$ replaced by $t_{\mathcal{A}_r^0}$. Using Hypothesis **(H1)**, let \mathcal{L}_i be the geometrization of the quasicharacter $\varphi_i := \varphi_{r_i}^i$ appearing in **(M2)**. If **(H1a)** applies, then $\mathcal{L}_i \in \mathbf{K}_{\bar{\mathbb{Q}}_\ell} \mathbf{D}_{c, G_i}^b(G_i; \bar{\mathbb{Q}}_\ell)$, using [Lus04, Theorem 21.21]; if **(H1b)** applies, then $\mathcal{L}_i \in \mathcal{CCS}(G_i)$, using Proposition 5.1. In either case, $\mathcal{L}_i \in \mathbf{K}_{\bar{\mathbb{Q}}_\ell} \mathbf{D}_{c, G_i}^b(G_i; \bar{\mathbb{Q}}_\ell)$ and

$$t_{\mathcal{L}_i} = \varphi_i.$$

For each i , let \mathcal{L}_r^i be the pull-back of \mathcal{L}_i along $G_r^i \rightarrow G_i$; then \mathcal{L}_r^i is a linear character sheaf and

$$t_{\mathcal{L}_r^i} = \varphi_r^i.$$

Now, suppose \mathcal{A}_r^i on G_r^i is defined such that

$$\begin{array}{ccc} G^i(k) & \xrightarrow{\mathrm{Tr}(\circ \rho^i)} & \bar{\mathbb{Q}}_\ell \\ & \searrow & \nearrow t_{\mathcal{A}_r^i} \\ & G_r^i(k) & \end{array}$$

commutes. Applying the Greenberg functor Gr_r^R to (5.4) gives

$$(5.6) \quad \begin{array}{ccccccc} & & & & 1 & & \\ & & & & \downarrow & & \\ & & & & G_r^i \times_{G_r} J_r^{i+1} & & \\ & & & & \downarrow & & \\ 1 & \longrightarrow & J_r^{i+1} & \longrightarrow & G_r^i \ltimes J_r^{i+1} & \xrightarrow{p_r^i} & G_r^i \longrightarrow 1 \\ & & & & \downarrow \pi_r^{i+1} & & \\ & & & & G_r^{i+1} & & \\ & & & & \downarrow & & \\ & & & & 1 & & \end{array}$$

where $J_r^{i+1} := \mathrm{Gr}_r^R(\underline{J}^{i+1})$ and $G_r^i := \mathrm{Gr}_r^R(\underline{G}^i)$. By [BGA18, Proposition 14.2], the sequences are exact. Consider $\mathcal{B}_r^{i+1} \in \mathbf{K}_{\bar{\mathbb{Q}}_\ell} \mathbf{D}_{c, G_r^i \ltimes J_r^{i+1}}^b(G_r^i \ltimes J_r^{i+1}; \bar{\mathbb{Q}}_\ell)$ defined by

$$\mathcal{B}_r^{i+1} := \mathcal{K}_r^{i+1} \otimes (p_r^i)^* (\mathcal{A}_r^i \otimes \mathcal{L}_r^i).$$

Comparing with (5.3), we see that $t_{\mathcal{B}_r^{i+1}}$ is precisely the function obtained by factoring the character of ${}^\circ\rho^{i+1}$ through $(\underline{G}^i \ltimes \underline{J}^{i+1})(R) \rightarrow (\underline{G}^i \ltimes \underline{J}^{i+1})(R_r)$ using the canonical identification $(G_r^i \times_{G_r} J_r^{i+1})(k) = (\underline{G}^i \ltimes \underline{J}^{i+1})(R_r)$. In particular, $t_{\mathcal{B}_r^{i+1}}$ is constant on $(G_r^i \times_{G_r} J_r^{i+1})(k)$, taking the value $\dim {}^\circ\rho^{i+1}$. With reference to the morphism $\pi_r^{i+1} : G_r^i \ltimes J_r^{i+1} \rightarrow G_r^{i+1}$ from (5.6), define

$$\mathcal{C}_r^{i+1} := (\pi_r^{i+1})_! (\mathcal{B}_r^{i+1}).$$

Then $\mathcal{C}_r^{i+1} \in \mathbf{K}_{\bar{\mathbb{Q}}_\ell} \mathbf{D}_{c, G_r^{i+1}}^b(G_r^{i+1}; \bar{\mathbb{Q}}_\ell)$ and

$$t_{\mathcal{C}_r^{i+1}}(x) = \sum_{y \in (\pi_r^{i+1})^{-1}(x)} t_{\mathcal{B}_r^{i+1}}(y).$$

Since $t_{\mathcal{B}_r^{i+1}}$ is constant on $(G_r^i \times_{G_r} J_r^{i+1})(k)$, it follows that

$$t_{\mathcal{C}_r^{i+1}} = n t_{\mathcal{B}_r^{i+1}}$$

on $G_r^{i+1}(k)$ with n equal to the product of $\#(G_r^i \times_{G_r} J_r^{i+1})(k)$ and $\dim {}^\circ\rho^{i+1}$. Let \mathcal{A}_r^{i+1} be the element of $\mathbf{K}_{\bar{\mathbb{Q}}_\ell} \mathbf{D}_{c, G_r^{i+1}}^b(G_r^{i+1}; \bar{\mathbb{Q}}_\ell)$ given by $\mathcal{A}_r^{i+1} = \frac{1}{n} \mathcal{C}_r^{i+1}$. This completes the inductive definition of \mathcal{A}_r^i so that the following diagram commutes.

$$\begin{array}{ccc} G_r^{i+1}(k) & \xrightarrow{\mathrm{Tr}({}^\circ\rho^{i+1})} & \bar{\mathbb{Q}}_\ell \\ & \searrow & \nearrow t_{\mathcal{A}_r^{i+1}} \\ & G_r^{i+1}(k) & \end{array}$$

Now set $\mathcal{F}_r^i = \mathcal{A}_r^i \otimes \mathcal{L}_r^i$, for $i = 0, \dots, d$. Then $\mathcal{F}_r^i \in \mathbf{K}_{\bar{\mathbb{Q}}_\ell} \mathbf{D}_{c, G_r^i}^b(G_r^i; \bar{\mathbb{Q}}_\ell)$ such that

$$\begin{array}{ccc} G^i(k) & \xrightarrow{\mathrm{Tr}(\circ \rho_i)} & \bar{\mathbb{Q}}_\ell \\ & \searrow & \nearrow t_{\mathcal{F}_r^i} \\ & G_r(k) & \end{array}$$

commutes. Define $\mathcal{F}^i \in \mathbf{K}_{\bar{\mathbb{Q}}} \mathbf{D}_{c, G^i}^b(G^i; \bar{\mathbb{Q}}_\ell)$ by pulling back \mathcal{F}_r^i along $G^i \rightarrow G_r^i$. Then

$$t_{\mathcal{F}^i} = \mathrm{Tr}(\circ \rho_i),$$

as desired. \square

BIBLIOGRAPHY

- [Adl98] Jeffrey D. Adler, *Refined anisotropic K -types and supercuspidal representations*, Pac. J. Math. **185** (1998), no. 1, 1–32. \uparrow 1415
- [Ber70] Jean-Étienne Bertin, *Exposé VI_B : Généralités sur les préschémas en groupes*, Propriétés générales des schémas en groupes. Séminaire de Géométrie Algébrique du Bois Marie, Schémas en groupes I (SGA3) (Michel Demazure and Alexander Grothendieck, eds.), Lecture Notes in Mathematics, vol. 151, Springer, 1970, pp. 318–410. \uparrow 1396
- [BGA18] Alessandra Bertapelle and Cristian D. González-Avilés, *The Greenberg functor revisited*, Eur. J. Math. **4** (2018), no. 4, 1340–1389. \uparrow 1409, 1417
- [BK98] Colin J. Bushnell and Philip C. Kutzko, *Smooth representations of reductive p -adic groups: structure theory via types*, Proc. Lond. Math. Soc. **77** (1998), no. 3, 582–634. \uparrow 1390
- [BL94] Joseph Bernstein and Valery Lunts, *Equivariant sheaves and functors*, Lecture Notes in Mathematics, vol. 1578, Springer, 1994. \uparrow 1399, 1407
- [CR18] Clifton Cunningham and David Roe, *From the function-sheaf dictionary to quasicharacters of p -adic tori*, J. Inst. Math. Jussieu **17** (2018), no. 1, 1–37. \uparrow 1391, 1392, 1393, 1394, 1395, 1396, 1400, 1402, 1403, 1404, 1405, 1410, 1415
- [Del77] Pierre Deligne, *Cohomologie étale. Séminaire de géométrie algébrique du Bois-Marie SGA 4 1/2 par P. Deligne, avec la collaboration de J.F. Boutot, A. Grothendieck, L. Illusie et J. L. Verdier*, Lecture Notes in Mathematics, vol. 569, Springer, 1977. \uparrow 1391
- [Del80] ———, *La conjecture de Weil. II*, Publ. Math., Inst. Hautes Étud. Sci. (1980), no. 52, 137–252. \uparrow 1393
- [FGI⁺05] Barbara Fantechi, Lothar Göttsche, Luc Illusie, Kleiman Steven L., Nitin Nitsure, and Angelo Vistoli, *Fundamental algebraic geometry*, Mathematical Surveys and Monographs, vol. 123, American Mathematical Society, 2005. \uparrow 1399
- [Fin18] Jessica Fintzen, *Types for tame p -adic groups*, <http://arxiv.org/abs/1810.04198>, 2018. \uparrow 1390
- [Fin19] ———, *On the construction of tame supercuspidal representations*, <http://arxiv.org/abs/1908.09819>, 2019. \uparrow 1390
- [Gab70] Pierre Gabriel, *Exposé VI_A : Généralités sur les groupes algébriques*, Propriétés générales des schémas en groupes. Séminaire de Géométrie Algébrique du Bois Marie, Schémas en groupes I (SGA3) (Michel Demazure and Alexander Grothendieck, eds.), Lecture Notes in Mathematics, vol. 151, Springer, 1970, pp. 287–317. \uparrow 1396, 1399

- [GH07] Shamgar Gurevich and Ronny Hadani, *The geometric Weil representation*, Sel. Math., New Ser. **13** (2007), no. 3, 465–481. ↑1392, 1410
- [GH11] ———, *The categorical Weil representation*, <http://arxiv.org/abs/1108.0351>, 2011. ↑1410
- [Gro67] Alexandre Grothendieck, *Éléments de géométrie algébrique IV. étude locale des schémas et des morphismes de schémas. IV*, Publ. Math., Inst. Hautes Étud. Sci. (1967), no. 32, 1–361, rédigé avec la collaboration de Jean Dieudonné. ↑1393
- [Hak18] Jeffrey Hakim, *Constructing tame supercuspidal representations*, Represent. Theory **22** (2018), 45–86. ↑1392
- [Kal19] Tasho Kaletha, *Regular supercuspidal representations*, J. Am. Math. Soc. **32** (2019), 1071–1170. ↑1392, 1415
- [Kam09] Masoud Kamgarpour, *Stacky abelianization of algebraic groups*, Transform. Groups **14** (2009), no. 4, 825–846. ↑1392, 1394
- [Kim07] Ju-Lee Kim, *Supercuspidal representations: an exhaustion theorem*, J. Am. Math. Soc. **20** (2007), no. 2, 273–320. ↑1390
- [Lus85] George Lusztig, *Character sheaves. I*, Adv. Math. **56** (1985), 193–237. ↑1392
- [Lus86] ———, *Character sheaves. V*, Adv. Math. **61** (1986), no. 2, 103–155. ↑1407
- [Lus04] ———, *Character sheaves on disconnected groups. IV*, Represent. Theory **8** (2004), 145–178. ↑1407, 1415, 1416
- [Ree08] Mark Reeder, *Supercuspidal L -packets of positive depth and twisted Coxeter elements*, J. Reine Angew. Math. **620** (2008), 1–33. ↑1415
- [RY14] Mark Reeder and Jiu-Kang Yu, *Epipelagic representations and invariant theory*, J. Am. Math. Soc. **27** (2014), no. 2, 437–477. ↑1415
- [Yu01] Jiu-Kang Yu, *Construction of tame supercuspidal representations*, J. Am. Math. Soc. **14** (2001), no. 3, 579–622. ↑1390, 1411, 1412
- [Yu15] ———, *Smooth models associated to concave functions in Bruhat–Tits theory*, Autour des schémas en groupes – École d’Été franco-asiatique de géométrie algébrique et de théorie des nombres. Volume III, Panoramas et Synthèses, vol. 47, Société Mathématique de France, 2015, http://smf4.emath.fr/Publications/PanoramasSyntheses/2016/47/html/smf_pano-synth_47_227-258.php, pp. 227–258. ↑1390, 1392, 1412, 1413, 1414

Manuscript received on 3rd July 2019,
 revised on 20th October 2020,
 accepted on 6th January 2021.

Recommended by Editor B. Edixhoven.
 Published under license CC BY 4.0.



This journal is a member of Centre Mersenne.



Clifton CUNNINGHAM
 Department of Mathematics and Statistics,
 University of Calgary,
 2500 University Drive Northwest,

Calgary, AB, (Canada), T2N 1N4
cunning@math.ucalgary.ca

David ROE
Department of Mathematics,
Massachusetts Institute of Technology,
77 Massachusetts Ave,
Cambridge, MA, (United States), 02139
roed@mit.edu