



ANNALES
HENRI LEBESGUE

THIBAUT DELCROIX

SIMON JUBERT

AN EFFECTIVE WEIGHTED K-STABILITY CONDITION FOR POLYTOPES AND SEMISIMPLE PRINCIPAL TORIC FIBRATIONS

UNE CONDITION DE K-STABILITÉ À
POIDS EFFECTIVE POUR LES POLYTOPES
ET LES FIBRATIONS PRINCIPALES
SEMISIMPLES TORIQUES

ABSTRACT. — The second author has shown that existence of extremal Kähler metrics on semisimple principal toric fibrations is equivalent to a notion of weighted uniform K-stability, read off from the moment polytope. The purpose of this article is to prove various sufficient conditions of weighted uniform K-stability which can be checked effectively and explore the low dimensional new examples of extremal Kähler metrics it provides.

Keywords: semisimple principal toric fibration, extremal Kähler metric, weighted cscK metric, uniform K-stability, projective bundle.

2020 *Mathematics Subject Classification:* 14M25, 32Q15, 32Q26, 53C55.

DOI: <https://doi.org/10.5802/ahl.161>

(*) The first author is partially funded by ANR-21-CE40-0011 JCJC project MARGE and ANR-18-CE40-0003-01 JCJC project FIBALGA, as well as PEPS JCJC INSMI CNRS projects 2021 and 2022. The second author was supported by PhD fellowships of the UQAM and of the Université de Toulouse III - Paul Sabatier.

RÉSUMÉ. — Le second auteur a démontré l'équivalence entre l'existence de métriques de Kähler extrémales sur les fibrations principales semisimples toriques et une notion de K-stabilité uniforme à poids, lue sur le polytope moment. Dans cet article, nous obtenons plusieurs conditions suffisantes de K-stabilité uniforme à poids qui peuvent être vérifiées de manière effectives, et nous explorons les nouveaux exemples que ces conditions fournissent en petite dimension.

1. Introduction

Calabi's work has been extremely influential in Kähler geometry, his name being still associated to some of the most fundamental objects of interest. The present article is motivated by two of these, Calabi's extremal Kähler metrics and Calabi's ansatz.

Extremal Kähler metrics provide a natural notion of canonical Kähler metrics in a given Kähler class on a compact Kähler manifold X : they are the metrics that achieve the minimum of the L^2 -norm of the scalar curvature. Kähler metrics with constant scalar curvature (cscK metrics for short) are special cases of such metrics, but Calabi showed in [Cal82] that there may exist extremal Kähler metrics when there exists no cscK metrics at all, by exhibiting extremal Kähler metrics on Hirzebruch surfaces. In order to show this, Calabi relied on the simple yet powerful idea that one should search for extremal Kähler metrics among those Kähler metrics that behave well with respect to the geometry of the manifold.

This was not a new idea of course. Matsushima showed for example [Mat57] that cscK metrics must behave well with respect to biholomorphism. More precisely, the automorphism group of X must be the complexification of the isometry group of the cscK metric, if it exists. This is preventing Hirzebruch surfaces from admitting cscK metrics as their automorphism group is non-reductive.

Calabi went further and restricted to metrics that respect the structure of \mathbb{P}^1 -bundles of Hirzebruch surfaces. He was then able to translate, for such metrics, the extremal property into a simple ODE and to solve it, showing the existence of extremal Kähler metrics. His construction was later referred to as Calabi's ansatz, used in various situations and generalized in various directions. It would be easy to fill pages with a bibliographical review of these, but it is not the purpose of this introduction. We only stress that a common theme is usually the desire to get explicit existence results or criterions. An influential illustration is [ACGTF08], where a variant of Calabi's ansatz was used to show that on various \mathbb{P}^1 -bundles, existence of extremal Kähler metrics reduces to checking the positivity of a polynomial on $[-1, 1]$, the so-called extremal polynomial. In the series of papers leading to [ACGTF08] (see also [HS02]), the general idea of Calabi's ansatz was actually pushed way further, allowing for example to consider certain fibrations with toric fiber.

The interest for such fibrations was significantly renewed last year, when the second author proved in [Jub21], using the breakthrough results of Chen and Cheng [CC18], that a uniform version of the Yau–Tian–Donaldson conjecture holds for semisimple principal toric fibrations, a very large class of toric fibrations. While it allows to translate the question of existence of extremal Kähler metrics on such manifolds into

a question of convex geometry on their moment polytopes, it is not yet an explicitly checkable criterion, as the conditions to check still form an infinite dimensional space. Motivated by the more practical philosophy behind Calabi's ansatz, we prove in the present paper various sufficient conditions of existence of extremal Kähler metrics which may be easily checked. Our approach is based on an initial idea by Zhou and Zhu [ZZ08], exploited in greater generality by the first author in [Del22].

The previous paragraphs are meant as an introduction to our results, and it should be stressed that it presents as such a biased and very incomplete historical account of the study of extremal Kähler metrics on manifolds with large symmetry. We refer to Székelyhidi's book [Szé14], Gauduchon's lecture notes [Gau10] for a general introduction to extremal Kähler metrics, and to Donaldson's remarkable survey [Don08] and Apostolov's lecture notes [Apo22] for some of the more specific aspects of manifolds with large symmetry. More recent developments very related to our work will be discussed at the beginning of Section 3.

Let us now highlight in the remainder of this introduction our main results. For this, a few notations are needed, and the full details will be given in Section 3. Semisimple principal toric fibrations are certain holomorphic fiber bundles $\pi : Y \rightarrow B$ where the basis $B = \prod_a B_a$ is a product of Hodge manifolds (B_a, ω_a) with constant scalar curvature s_a , and where the fiber X is toric under a compact torus \mathbb{T} . They are constructed from certain types of principal \mathbb{T} -bundles, essentially determined by the data of a tuple (p_a) of one-parameter subgroups of \mathbb{T} . In this paper, a one-parameter subgroup $p_a : \mathbb{S}^1 \rightarrow \mathbb{T}$ of \mathbb{T} will be identified with the element of the Lie algebra of \mathbb{T} determined by the image of $1 \in \mathbb{R} = \text{Lie}(\mathbb{S}^1)$ under the differential of p_a at the neutral element. In particular, it defines a linear function on the dual of the Lie algebra of \mathbb{T} . On such manifolds, a Kähler class is called compatible if it decomposes as the sum of a relative Kähler class induced by a Kähler class $[\omega_X]$ on X , and a sum of real multiples $c_a \pi^*[\omega_a]$ of the pull-backs of the Kähler classes $[\omega_a]$. An admissible Kähler class contains admissible Kähler metrics, that behave well with respect to the fibration structure.

THEOREM 1.1. — *Assume that Y is a semisimple principal toric fibration, that the toric fiber X is Fano equipped with the Kähler class $[\omega_X] = t2\pi c_1(X)$, and let $[\omega_Y]$ be an admissible Kähler class. Assume that for all a , $2 \dim(B_a)c_a \geq ts_a$ and that at every vertex x of the moment polytope P of $(X, [\omega_X])$,*

$$2(\dim(Y) + 1) + \sum_a \frac{ts_a - 2 \dim(B_a)c_a}{p_a(x) + c_a} - tl_{\text{ext}}(x) \geq 0$$

where l_{ext} is the extremal affine function. Then there exists an extremal Kähler metric in $[\omega_Y]$.

Recall that, when a maximal torus of automorphisms of Y is fixed, the scalar curvature of an invariant extremal Kähler metric, if it exists, is a holomorphy potential of a well defined vector field called the extremal vector field. In the statement above, the extremal function is encoding the extremal vector field, and a choice of maximal torus of automorphisms of Y is implicitly assumed. We will explain why it reduces to an affine function on the polytope P in Section 3.

We actually prove a much more general sufficient condition, Theorem 2.6, that does not require the fiber to be Fano. Since we obtain already a wealth of new examples with this particular case, and it is a natural generalization of the \mathbb{P}^1 -bundle case, we focus on this result for the introduction.

In the case when the fibration is Fano itself, and not only its fiber, then $t = 1$ and $s_a = 2 \dim(B_a)c_a$ so one gets a particularly simple criterion:

COROLLARY 1.2. — *A Fano semisimple principal toric fibration Y admits an extremal Kähler metric in $c_1(Y)$ if its extremal affine function l_{ext} satisfies*

$$\sup l_{\text{ext}} \leq 2(\dim(Y) + 1)$$

and the latter obviously needs only be verified at vertices of the moment polytope.

We provide, for the reader's convenience, an elementary Python program implementing the sufficient condition from Theorem 1.1 in the case when there is only one factor in B and the fiber is of dimension one or two. It would be easy to imitate these to allow greater flexibility in the data. It may be used either with all the data given numerically, or some of the data treated as variable. We use this to our advantage to prove the existence of extremal Kähler metrics in a wide range of Kähler classes for some examples of fibrations.

PROPOSITION 1.3. — *Let $Y = \mathbb{P}_B(\mathcal{O}_B \oplus H^{-p_1} \oplus H^{-p_2})$, where B is a Kähler–Einstein Fano threefold, H is the smallest integral divisor of $2\pi c_1(B)$ and $1 \leq p_1 \leq p_2$. Then there exists an extremal Kähler metric in the Kähler class $c_1(X) + \lambda c_1(B)$ for $\lambda \geq 9p_2$, where $c_1(X)$ and $c_1(B)$ respectively denote the relative first Chern class and the pull-back of the first Chern class, by an abuse of notations.*

Here, Y is a semisimple principal fibration over the base B , with fiber the projective space $X = \mathbb{P}^2$. More generally, projectivizations of direct sums of line bundles can often be considered as semisimple principal fibrations, as explained in Section 3.3.

The article is organized as follows. In Section 2, we prove a general sufficient condition for weighted uniform K-stability of labelled polytopes, and consider the special case of monotone polytopes. Section 3 explains the geometric origin of weighted uniform K-stability of labelled polytopes, with a particular emphasis on semisimple principal toric fibrations. In Section 4, we put together the two aspects to prove Theorem 1.1 and Corollary 1.2 using the monotone case of Section 2, as well as more general statements. We present various examples of applications of the sufficient condition in Section 5, including Proposition 1.3. Finally, we include in an appendix elementary Python programs computing the sufficient condition for fibrations with only one factor in the basis, and a one or two dimensional Fano fiber.

Acknowledgements

The authors are very grateful to Vestislav Apostolov and Eveline Legendre for their valuable comments on the manuscript. The authors also thank the referee for his useful suggestions and corrections.

2. Weighted K-stability of labelled polytopes: a sufficient condition

2.1. Weighted K-stability of labelled polytopes

Let V be an affine space of dimension ℓ , equipped with a fixed Lebesgue measure dx . A *labelled polytope* in V is a pair (P, \mathbf{L}) where P is a (compact, convex) polytope in V and $\mathbf{L} = (L_j)_{j=1}^d$ is a minimal set of defining affine functions for P , that is,

$$P = \{x \in V \mid \forall j, \quad L_j(x) \geq 0\}$$

where d is the number of facets (codimension one faces) of P . We denote by $F_j := \{x \in P \mid L_j(x) = 0\}$ the facet of P defined by L_j .

DEFINITION 2.1. — *The labelled boundary measure $d\sigma$ is the measure on ∂P whose restriction to the facet F_j is defined by $dL_j \wedge d\sigma = -dx$.*

Note that the labelled boundary measure depends heavily on the choice of labelling (L_j) . For example, for any tuple (r_j) of positive real numbers, the tuple $(L'_j) = (r_j L_j)$ is another labelling of P . The associated labelled boundary measure $d\sigma'$ satisfies $d\sigma' = \frac{1}{r_j} d\sigma$ on F_j . In particular, if the r_j are not all equal, there is no obvious relation between $d\sigma$ and $d\sigma'$. Similarly, the notion of weighted uniform K-stability that we are about to define depends heavily on the choice of labelling.

Let v be a continuous, positive function on P , and let w be a continuous function on P . Following [Don02, Lah19, LLS16], we define the (v, w) -Donaldson–Futaki invariant of the labelled polytope (P, \mathbf{L}) as the functional \mathcal{F} on the space of continuous functions on P such that

$$(2.1) \quad \mathcal{F}(f) := 2 \int_{\partial P} f(x)v(x)d\sigma - \int_P f(x)w(x)dx.$$

Following [His20, NS21], we also set

$$\|f\|_J = \inf_{l \in \text{Aff}(V)} \int_P \left(f + l - \inf_P(f + l) \right) dx$$

where $\text{Aff}(V)$ denotes the space of affine functions on V .

DEFINITION 2.2. — *A labelled polytope (P, \mathbf{L}) is (v, w) -uniformly K-stable if there exists a $\lambda > 0$ such that for any continuous convex functions f on P ,*

$$(2.2) \quad \mathcal{F}(f) \geq \lambda \|f\|_J.$$

Remark 2.3. — Note that \mathcal{F} is linear, and the right-hand side of (2.2) is always non-negative, hence the following is a necessary condition for (2.2) to hold:

$$(2.3) \quad \forall f \in \text{Aff}(V), \mathcal{F}(f) = 0.$$

We will explain in Section 3 the geometric significance of this notion for various choices of v and w , let us for now just highlight that when v and w are constant, the functional \mathcal{F} first appeared in [Don02] as an expression of the (Donaldson–)Futaki invariant for toric test configurations in the study of K-stability of toric manifolds, whence the name.

We denote by $\mathcal{CV}^0(P)$ the space of continuous convex functions on P , and by $\mathcal{CV}^1(P)$ the space of all convex functions f on P which are the restrictions to P of a continuously differentiable function defined on an open subset of V containing P . Note that by uniform approximation by smooth functions, it is enough to consider only functions in $\mathcal{CV}^1(P)$ to check condition (2.2).

In order to deal more efficiently with the right hand side of (2.2), following [Don02], we consider the following normalization of functions. We choose a point x_0 in the interior P^0 of the polytope P . It allows to choose a linear complement $\mathcal{CV}_*^1(P)$ to $\text{Aff}(V)$ in $\mathcal{CV}^1(P)$, defined by

$$(2.4) \quad \mathcal{CV}_*^1(P) := \left\{ f \in \mathcal{CV}^1(P) \mid \forall x, f(x) \geq f(x_0) = 0 \right\}.$$

Then, any $f \in \mathcal{CV}^1(P)$ can be written uniquely as $f = f^* + f_0$, where f_0 is affine and $f^* \in \mathcal{CV}_*^1(P)$, and we will use these notations in the following. By linearity, $\mathcal{F}(f) = \mathcal{F}(f^*)$ if \mathcal{F} vanishes on $\text{Aff}(V)$.

LEMMA 2.4. — *The labelled polytope (P, \mathbf{L}) is (v, w) -uniformly K -stable if and only if there exists $\lambda > 0$ such that for all $f \in \mathcal{CV}^1(P)$,*

$$(2.5) \quad \mathcal{F}(f) \geq \lambda \|f^*\|_{L^1}$$

where $\|\cdot\|_{L^1}$ denotes the L^1 -norm on P with respect to the Lebesgue measure dx .

Proof. — From [NS21, Proposition 4.1(3)], there exists a constant $C_1 > 0$ such that for all continuous convex functions on P ,

$$\|f\|_J \leq \|f\|_{L^1} \leq C_1 \|f\|_J$$

The equivalence between condition (2.2) and condition (2.5) follows immediately. \square


Remark 2.5. — Condition (2.5) is the condition that we will effectively use in the sequel, so one might wonder why we introduced the first definition. The point is that by Lemma 2.4, condition (2.5) is independent of the choice of x_0 , and condition (2.2) makes it perfectly clear. In the more familiar unweighted case, the equivalence between various notions of K -stability of polytopes was fully worked out by Nitta and Saito [NS21].

2.2. The sufficient condition

Let $\mathcal{C}^0(P, \mathbb{R})$ denote the space of continuous functions on P , and let $\mathcal{C}^1(P, \mathbb{R})$ denote the space of functions that are the restriction to P of continuously differentiable functions defined in an open subset of V containing P .

Recall that F_j denotes the facet of P defined by L_j . For each j , let P_j be the cone with basis F_j and vertex x_0 as illustrated in Figure 2.2. For a function $f \in \mathcal{C}^1(P, \mathbb{R})$, we denote by $d_x f$ its differential at $x \in P$. The following is the main technical result of our paper, it imitates quite closely part of the proof by Zhou and Zhu [ZZ08] of a coercivity criterion for the modified Mabuchi functional on toric manifolds.

Figure 2.1. The cone decomposition



THEOREM 2.6. — Let $v \in \mathcal{C}^1(P, \mathbb{R})$ be a positive function on P and let $w \in \mathcal{C}^0(P, \mathbb{R})$. Assume that \mathcal{F} vanishes on $\text{Aff}(V)$ and that for all $j = 1, \dots, d$, for all $x \in P_j$,

$$(2.6) \quad \frac{1}{L_j(x_0)} \left(v(x)(\ell + 1) + d_x v(x - x_0) \right) - \frac{w(x)}{2} \geq 0,$$

then (P, \mathbf{L}) is (v, w) -uniformly K -stable.

Proof. — Since $L_j(x) = 0$ for $x \in F_j$, we have $L_j(x_0) = d_x L_j(x_0 - x)$ for all $x \in F_j$. In particular,

$$\int_{F_j} f(x)v(x)d\sigma = \int_{F_j} f(x)v(x) \frac{-d_x L_j(x - x_0)}{L_j(x_0)} d\sigma.$$

For each facet F of ∂P_j different from F_j , and $x \in F$, the interior product $\iota_{x-x_0}(dx)$ vanishes since it vanishes on the affine space spanned by F . If we further use that $-dL_j \wedge d\sigma = dx$ on F_j , we obtain

$$\int_{F_j} f(x)v(x)d\sigma = \frac{1}{L_j(x_0)} \int_{\partial P_j} f(x)v(x)\iota_{x-x_0}(dx).$$

Hence by Stokes theorem we obtain

$$\int_{F_j} f(x)v(x)d\sigma = \frac{1}{L_j(x_0)} \int_{P_j} \left(v(x)d_x f(x - x_0) + \ell f(x)v(x) + f(x)d_x v(x - x_0) \right) dx.$$

Summing the previous identities over j we get

$$(2.7) \quad \begin{aligned} \mathcal{F}(f) &= \sum_{j=1}^d \frac{2}{L_j(x_0)} \int_{P_j} \left(d_x f(x - x_0) - f(x) \right) v(x) dx \\ &+ \sum_{j=1}^d \int_{P_j} \left(\frac{2}{L_j(x_0)} \left((\ell + 1)v(x) + d_x v(x - x_0) \right) - w(x) \right) f(x) dx. \end{aligned}$$

Assume condition (2.6) is satisfied and (P, \mathbf{L}) is not (v, w) -uniformly K -stable.

We will show contradiction to a stronger condition than condition (2.5). Namely, assume that condition (2.6) is satisfied and that there exists a sequence of $\{f_k\}_{k \in \mathbb{N}}$ in $\mathcal{CV}^1(P)$ such that

$$(2.8) \quad \lim_{k \rightarrow \infty} \mathcal{F}(f_k^*) = 0 \quad \text{and} \quad \forall k \in \mathbb{N}, \int_{\partial P} v(x) f_k^*(x) d\sigma = 1.$$

Recall from [Don02, Lemma 5.1.3] (see [NS21, Proposition 5.1.2] for a detailed proof and explicit constant C) that there exists a positive constant $C > 0$ such that for all $f \in \mathcal{CV}^1(P)$,

$$\int_{\partial P} f^*(x) d\sigma \geq C \|f^*\|_{L^1}.$$

As a consequence, since $v > 0$ on P , there exists a constant $C' > 0$ such that for all $f \in \mathcal{CV}^1(P)$,

$$\int_{\partial P} v(x) f^*(x) d\sigma \geq C' \|f^*\|_{L^1}.$$

In particular, the sequence $\{f_k^*\}$ has bounded L^1 norm. By [Don02, Corollary 5.2.5], $\{f_k^*\}_{k \in \mathbb{N}}$ converges (up to a sub-sequence, still denoted by f_k^*) locally uniformly in P^0 to a convex function f_∞^* which still satisfies $\inf f_\infty^* = f_\infty^*(x_0) = 0$. Since in addition all f_k^* are smooth and convex we have $d_x f_k^*(x - x_0) - f_k^*(x) \geq 0$. Then, since condition (2.6) is assumed to hold, all terms of the sum in (2.7) are non-negative. Evaluating (2.7) at f_k^* and passing to the limit reveals that $\lim_{k \rightarrow \infty} d_x f_k^*(x - x_0) - f_k^*(x) = 0$ almost everywhere in P^0 , showing that f_∞^* is affine on P^0 . Using again $\inf f_\infty^* = f_\infty^*(x_0) = 0$, we conclude that f_∞^* is the zero function on P^0 .

The local uniform convergence of $\{f_k^*\}$ to the zero function shows that

$$\lim_{k \rightarrow \infty} \int_P f_k^*(x) w(x) dx = 0$$

that is,

$$\lim_{k \rightarrow \infty} 2 \int_{\partial P} f_k^*(x) v(x) d\sigma - \mathcal{F}(f_k^*) = 0$$

which is in contradiction with condition (2.8).

From this contradiction it follows that there exists a constant $\mu > 0$ such that for all $f \in \mathcal{CV}^1(P)$,

$$\begin{aligned} \mathcal{F}(f) &\geq \mu \int_{\partial P} v(x) f^*(x) d\sigma \\ &\geq \mu C' \|f^*\|_{L^1} \end{aligned}$$

which concludes the proof of the Theorem 2.6. □

Remark 2.7. — We stress that the property of (v, w) -uniform K -stability is independent of the choice of $x_0 \in P^0$ in the previous section, but condition (2.6) depends on that choice. It is possible and useful in practical uses of the condition to vary this x_0 according to the data of the problem, see 5.3.1.

Remark 2.8. — Condition (2.6) depends continuously on the labelled polytope, the weights v and w , and the choice of x_0 .

2.3. Case of monotone polytopes

Let us recall the terminology of monotone polytopes, used in [Leg16].

DEFINITION 2.9. — *A labelled polytope (P, \mathbf{L}) is monotone if there exists an $x_0 \in P^0$ such that $L_1(x_0) = L_2(x_0) = \dots = L_d(x_0)$.*

There is thus an obvious choice of x_0 in that case. Our sufficient condition indeed becomes much simpler in that case, since the decomposition of the polytope may essentially be forgotten.

COROLLARY 2.10. — *Let (P, \mathbf{L}) be a monotone labelled polytope with $L_1(x_0) = L_2(x_0) = \dots = L_d(x_0) = t$. Let $v \in \mathcal{C}^1(P, \mathbb{R})$ such that v is positive on P and let $w \in \mathcal{C}^0(P, \mathbb{R})$. Assume that \mathcal{F} vanishes on $\text{Aff}(V)$ and that for all $x \in P$,*

$$(2.9) \quad \frac{1}{t} \left(v(x)(\ell + 1) + d_x v(x - x_0) \right) - \frac{w(x)}{2} \geq 0,$$

then (P, \mathbf{L}) is (v, w) -uniformly K-stable.

The conditions involved form a finite set of conditions to check, contrary to the definition of (v, w) -uniform K-stability. It is furthermore easy to implement in a computer program, *via* formal or numerical computations depending on the data (P, \mathbf{L}, v, w) . The same is true for the more general Theorem 2.6, but the decomposition in cones makes it a bit more tedious.

3. Geometric origin of weighted K-stability of polytopes

3.1. Weighted cscK toric manifolds

The results from Section 2 are motivated by the study of the existence of weighted cscK metrics on toric manifolds, as studied in [Jub21].

Let \mathbb{T} be an ℓ -dimensional compact torus. We denote by \mathfrak{t} its Lie algebra and by $\Lambda \subset \mathfrak{t}$ the lattice of generators of circle subgroups, so that $\mathbb{T} = \mathfrak{t}/2\pi\Lambda$. Let (X, ω, \mathbb{T}) be a compact Kähler toric manifold. Denote by μ the moment map of X with respect to the action of \mathbb{T} , and let $P = \mu(X) \subset \mathfrak{t}^*$ be the moment polytope. The polytope P is a Delzant polytope [Del88], and in particular, there is a natural choice of labelling \mathbf{L} of P such that all the differentials dL_j of the defining affine functions L_j are primitive elements in the lattice Λ .

Remark 3.1. — We focus here on smooth manifolds, but let us mention that the cases of orbifolds or pairs would also be natural settings to consider. In these situations, the labelling could be more general (see [Abr01] for orbifolds and [DGSW18] for pairs), thus justifying our choice to allow arbitrary labellings in the previous section.

In the context of toric manifold, the v -weighted scalar curvature was introduced in [LLS16]. To avoid introducing too much notation, we give the definition of [Lah19], which makes sense for general compact Kähler manifold and coincide with the one

of [LLS16] in the toric context. Let $\mathcal{C}^\infty(P, \mathbb{R})$ denote the space of restrictions to P of smooth functions defined on an open set containing P , and $\mathcal{C}^\infty(P, \mathbb{R}_{>0})$ the subspace of positive functions.

DEFINITION 3.2 (Weighted cscK metrics). —

(1) For $v \in \mathcal{C}^\infty(P, \mathbb{R}_{>0})$, define the v -scalar curvature of ω as the function

$$\text{Scal}_v(\omega) := v(\mu)\text{Scal}(\omega) + 2\Delta_\omega(v(\mu)) + \text{Tr}(G_\omega \circ (\text{Hess}(v) \circ \mu)),$$

where $\text{Scal}(\omega)$ is the usual scalar curvature of the Riemannian metric g_ω associated to ω , Δ_ω is the Riemannian Laplacian of g_ω , $\text{Hess}(v)$ is the Hessian of v viewed as a bilinear form on \mathfrak{t}^* whereas G_ω is the bilinear form with smooth coefficients on \mathfrak{t} , given by the restriction of g_ω on fundamental vector fields.

(2) If furthermore $w \in \mathcal{C}^\infty(P, \mathbb{R})$, then ω is a (v, w) -cscK metric if

$$\text{Scal}_v(\omega) = w \circ \mu$$

In general, no YTD correspondence is proved for the existence of weighted cscK metrics on toric manifolds. However by analogy with the unweighted cscK case, there is a known candidate for the corresponding K-stability condition, which translates on the polytope as Definition 2.2. In fact, the direction from existence of weighted cscK metrics to K-stability was proved in general by Li, Lian, Sheng [LLS17].

THEOREM 3.3 ([LLS17, Theorem 2.1]). — *If ω is a (v, w) -cscK metric, then (P, \mathbf{L}) is (v, w) -uniformly K-stable.*

The converse direction is in general much harder, but is known for special choices of weights.

- If v and w are constants, this is the uniform YTD conjecture for cscK metrics on toric manifolds. If v is constant and w is affine, this is the uniform YTD conjecture for extremal metrics on toric manifolds. Both these conjectures were proved recently [Apo22, His20, Leg19, Li20, LLS21, NS21] thanks to the breakthrough of Chen–Cheng [CC18], its adaptation by He to the extremal setting [He19], and earlier works, notably [Don02, ZZ08].
- If only v is constant, the converse of Theorem 3.3 is known for all $w \in \mathcal{C}^\infty(P, \mathbb{R})$ by [LLS21].
- For v -solitons on Fano toric manifolds, which correspond to choosing an arbitrary weight $v \in \mathcal{C}^\infty(P, \mathbb{R}_{>0})$ and $w(x) = 2(\ell v(x) + d_x v(x))$ (see [AJL22, Proposition 1]), it was proved first in [BB13] that the converse of Theorem 3.3 holds for general weight v , and much earlier in [WZ04] for the weight corresponding to Kähler–Ricci solitons. We note that [LH20] proved that the general uniform YTD conjecture holds for v -solitons on general Fano manifolds, and refer to Section 4.4 for a discussion of v -solitons on semisimple principal toric fibrations.
- Finally, as we shall explain in details in the next sections, the converse of Theorem 3.3 was proven by the second author for weights corresponding to extremal Kähler metrics on semisimple principal toric fibrations [Jub21].

3.2. Construction of semisimple principal toric fibration

In this section we briefly recall the construction of semisimple principal toric fibrations introduced in [ACGTF04]. We take the point of view of [AJL22], which generalized the construction when the fiber is not necessarily toric.

Let \mathbb{T} be an ℓ -dimensional compact torus with Lie algebra \mathfrak{t} . Let $\pi_B : Q \rightarrow B$ be a principal \mathbb{T} -bundle over a n -dimensional product of cscK manifold $(B, J_B, \omega_B) := \prod_{a=1}^k (B_a, J_a, \omega_a)$ satisfying the Hodge condition $\omega_a \in H^2(B_a, \mathbb{Z})$. We supposed that Q is equipped with a principal connection $\theta \in \Omega^1(Q) \otimes \mathfrak{t}$ whose curvature satisfies

$$d\theta = \sum_{a=1}^k \pi_B^*(\omega_a) \otimes p_a,$$

where $p_a \in \mathfrak{t}$ define one-parameter subgroups of \mathbb{T} . Let (X, J_X, ω_X) be a toric projective manifold under the action of \mathbb{T} . Since \mathbb{T} acts on various spaces, to avoid confusion, we under-script the space on which \mathbb{T} acts, e.g. \mathbb{T}_X acts on X . We consider the $2(\ell + n)$ dimensional smooth manifold

$$(3.1) \quad Y := (X \times Q)/\mathbb{T},$$

where the $\mathbb{T}_{X \times Q}$ -action on $(X \times Q)$ is given by $\gamma \cdot (x, q) = (\gamma \cdot x, \gamma^{-1} \cdot q)$, $x \in X$, $q \in Q$ and $\gamma \in \mathbb{T}$. Let $\mathcal{H} := \text{ann}(\theta) \subset TQ$ be the horizontal distribution on the principal bundle Q with respect to θ . We consider the smooth section of the endomorphism of $T(X \times Q)$

$$J_Y := J_X \oplus J_B,$$

where J_B acts on Q via the unique horizontal lift of vector fields on B to \mathcal{H} . The section J_Y is invariant with respect to the $\mathbb{T}_{Q \times X}$ -action and it is shown in [AJL22, Section 5] that J_Y descends to a complex structure (still denoted by J_Y) on Y .

Let $P \subset \mathfrak{t}^*$ be the Delzant polytope associated to $(X, J_X, \omega_X, \mathbb{T})$ [Del88]. By definition the \mathbb{T} -action on X is hamiltonian and we denote by $\mu : X \rightarrow P$ its moment map. We consider the 2-form on $X \times Q$

$$(3.2) \quad \omega_Y := \omega_X + \sum_{a=1}^k (p_a \circ \mu + c_a) \pi_B^*(\omega_a) + d\mu \circ \theta,$$

where the k -tuple of real constants (c_a) are such that for all a , $p_a \circ \mu + c_a > 0$ and $d\mu \circ \theta$ is a 2-form understood as the contraction of the \mathfrak{t} -valued one form θ and the \mathfrak{t}^* -valued one form $d\mu$. It is shown in [AJL22, Section 5] that ω_Y is basic with respect to $\mathbb{T}_{X \times Q}$ and as such, it is the pullback of a Kähler form (still denoted by ω_Y) on Y . The \mathbb{T}_X -action on X induces an action on $X \times Q$ by the natural \mathbb{T}_X -action on the first factor. This action commutes with $\mathbb{T}_{X \times Q}$ and therefore descends to a \mathbb{T}_Y -action on Y . The Kähler form ω_Y on Y defined via the basic 2-form (3.2) on $X \times Q$ is \mathbb{T}_Y -invariant (see [AJL22, Section 5] for more details).

The Kähler metrics ω_a on B_a , the connection form θ and the constants $p_a \in \mathfrak{t}$ are fixed. The Kähler manifold $(Y, J_Y, \omega_Y, \mathbb{T})$ is a fiber bundle over B with fiber the toric Kähler manifold $(X, J_X, \omega_X, \mathbb{T})$. Following [ACGTF11] we define:

DEFINITION 3.4. — *The Kähler manifold $(Y, J_Y, \omega_Y, \mathbb{T})$ defined above is called a semisimple principal toric fibration. The \mathbb{T} -invariant Kähler metric ω_Y on Y defined*

from the \mathbb{T} -invariant Kähler metric ω_X on X is called a compatible Kähler metric. A Kähler class $\alpha_Y \in H^2(Y, \mathbb{R})$ containing a compatible Kähler metric is called a compatible Kähler class.

In this setup, the constants c_a can vary and they parameterize the compatible Kähler classes.

Let P^0 be the interior of P and $\mathring{X} := \mu^{-1}(P^0)$ be the dense open subset of X of regular orbits for the \mathbb{T}_X -action. The \mathbb{T} -action on X extends to an effective holomorphic action of the complexified torus $\mathbb{T}^{\mathbb{C}} := \mathbb{T} \otimes \mathbb{C}$. Fixing any point $x_0 \in \mathring{X}$, we can identify (\mathring{X}, J_X) and the orbit $\mathbb{T}^{\mathbb{C}} \cdot x_0 \cong (\mathbb{C}^*)^{\ell}$ of x_0 :

$$(3.3) \quad (\mathring{X}, J_X) \cong (\mathbb{C}^*)^{\ell}.$$

Restricting the $\mathbb{T}_{X \times Q}$ -action to $\mathring{X} \times Q$, we define

$$\mathring{Y} := (\mathring{X} \times Q) / \mathbb{T}.$$

By (3.3) \mathring{Y} is \mathbb{T} -equivariantly biholomorphic to a $(\mathbb{C}^*)^{\ell}$ -bundle over B . Since (\mathring{X}, J_X) compactifies to (X, J_X) , the complex principal $(\mathbb{C}^*)^{\ell}$ -bundle \mathring{Y} compactifies to (Y, J_Y) .

Remark 3.5. — Semisimple principal toric fibrations $(Y, J_Y, \omega_Y, \mathbb{T})$ constructed above correspond to semisimple rigid toric fibration introduced and studied in [ACG06, ACGTF04, ACGTF08] when there is no blow-down and the basis B is a global product of cscK Hodge manifolds.

A particular case, which will be of interest for us, is the case of Fano semisimple principal toric fibrations. We recall a characterization of these from [AJL22]

LEMMA 3.6 ([AJL22, Lemma 5.11]). — *Assume that each B_a is a Fano Kähler-Einstein manifold. Let ω_a denote a Kähler-Einstein metric on B_a so that $I_a[\omega_a] = 2\pi c_1(B_a)$, where I_a denotes the Fano index of B_a . We fix a principal bundle with connection (Q, θ) as before (with associated data (p_a)). We further assume that (X, ω_X) is a Fano toric manifold with a \mathbb{T} -invariant Kähler form $\omega_X \in 2\pi c_1(X)$, with the natural choice of moment map μ . If for all a , $p_a \circ \mu + I_a > 0$, then the semisimple principal fibration Y associated to the above data is a Fano manifold, and ω_Y is in $2\pi c_1(Y)$ for the k -tuple $(c_a) = (I_a)$.*

Note that in the above situation, the scalar curvature of ω_a is indeed constant, equal to $2n_a I_a$ where n_a is the complex dimension of B_a .

3.3. Projectivization of sum of line bundles as semisimple principal toric fibration

We now provide an effective way of constructing examples of semisimple principal toric fibrations.

Let $(B, \omega_B) := \prod_{a=1}^k (B_a, \omega_a)$ be a product of compact complex manifolds B_a endowed with cscK metrics ω_a with $[\omega_a]$ primitive element of $H^2(B_a, \mathbb{Z})$. We consider

holomorphic line bundles $\mathcal{L}_i \rightarrow B, i = 1, \dots, \ell$, and we suppose that their first Chern classes satisfy

$$-2\pi c_1(\mathcal{L}_i) = \sum_{a=1}^k p_{ai}[\omega_a],$$

where by definition p_{ai} is the ω_a -degree of \mathcal{L}_i^* . The natural \mathbb{C}_i^* -action on \mathcal{L}_i induces an action of $\mathbb{S}_i^1 \subset \mathbb{C}_i^*$, providing a \mathbb{T} -bundle $\pi : Q \rightarrow B$, where $\mathbb{T} = \prod_i \mathbb{S}_i^1$ is a compact ℓ -torus. We choose a Hermitian metric h_i on \mathcal{L}_i and consider the norm function $r_i(u) := (h_i(u, u))^{\frac{1}{2}}$ for any u in \mathcal{L}_i . On $\tilde{\mathcal{L}}_i$, the \mathbb{C}^* -bundle obtained from \mathcal{L}_i by removing the zero section, r_i is positive and we let $t_i = \log(r_i)$. We fix a basis $\xi = (\xi_i)_{i=1}^\ell$ of the Lie algebra \mathfrak{t} of \mathbb{T} and we denote by $\xi^{\tilde{\mathcal{L}}_i}$ the generator of the \mathbb{S}_i^1 -action on $\tilde{\mathcal{L}}_i$. We then consider the \mathfrak{t} -valued one-form $t := \sum_{i=1}^\ell t_i \xi_i$ and we define a connection one-form θ on Q as the restriction of $d^c t$ to Q , seen as the \mathbb{T} -bundle of unit element on each (\mathcal{L}_i, h_i) . For all $i = 1, \dots, \ell$, it satisfies

$$\theta(\xi^{\tilde{\mathcal{L}}_i}) = \xi_i.$$

We obtain by construction

$$\begin{aligned} d\theta &= \sum_{i=1}^\ell \xi_i \otimes \pi^*(\omega_{h_i}) = \sum_{i=1}^\ell \xi_i \otimes \left(\sum_{a=1}^k p_{ai} \pi^*(\omega_a) \right) \\ (3.4) \quad &= \sum_{a=1}^k p_a \otimes \pi^*(\omega_a), \end{aligned}$$

where ω_{h_i} is the opposite of the curvature form of the Chern connection of (\mathcal{L}_i, h_i) and $p_a = \sum_{i=1}^\ell p_{ai} \xi_i$. We consider the ℓ -projective space $(\mathbb{P}^\ell, \omega_{\mathbb{P}^\ell}, \mathbb{T})$ endowed of a toric \mathbb{T} -action with respect to a fixed Kähler metric $\omega_{\mathbb{P}^\ell}$. We fix the principal \mathbb{T} -bundle Q with its connection one form θ , the cscK Kähler manifolds (B_a, ω_a) and the toric Kähler manifold $(\mathbb{P}^\ell, \omega_{\mathbb{P}^\ell}, \mathbb{T})$. From these data, we define a semisimple principal toric fibration $Y := Q \times_{\mathbb{T}} \mathbb{P}^\ell$. By construction, Y is biholomorphic to the total space of the projective bundle $\mathbb{P}(E), E := \mathcal{O} \oplus \bigoplus_{i=1}^\ell \mathcal{L}_i$.

Suppose $\omega_{\mathbb{P}^\ell}$ belongs to the first Chern class $2\pi c_1(\mathbb{P}^\ell)$ of \mathbb{P}^ℓ and denote by P the canonical ℓ -simplex associate to $(\mathbb{P}^\ell, 2\pi c_1(\mathbb{P}^\ell), \mathbb{T}^\ell)$ via Delzant correspondence [Del88]. By (3.4), any compatible Kähler metric on Y is of the form

$$(3.5) \quad \omega_Y = \sum_{a=1}^k \left(\sum_{i=1}^\ell p_{ai} x_i + c_a \right) \pi^*(\omega_a) + \omega_{\mathbb{P}^\ell}, \mathbf{x} = (x)_{i=1}^\ell \in P$$

with

$$(3.6) \quad -c_a > \sum_{i=1}^\ell p_{ai},$$

In the above formulas, by abuse of notation, $\omega_{\mathbb{P}^d}$ denotes both the Kähler metric on \mathbb{P}^ℓ and its induced metric in $2\pi c_1(O_E(\ell + 1))$. The tuples (c_a) satisfying (3.6), parametrize the compatible Kähler classes.

Furthermore, suppose that B is a product of Kähler-Einstein manifolds $(B, \omega_B) := \prod_{a=1}^k (B_a, \omega_a)$. By Lemma 3.6, if we choose c_a equal to the Fano index I_a of B_a , the

corresponding compatible Kähler form ω_Y defined in (3.5) belongs to the first Chern class $2\pi c_1(Y)$. In particular, if

$$(3.7) \quad -I_a > \sum_{i=1}^{\ell} p_{ai},$$

Y is a Fano manifold with compatible first Chern class.

3.4. Extremal metrics on semisimple principal toric fibrations

We begin this section by recalling a general characterisation of extremal metrics on compact Kähler manifold (Y, J_Y) in a fixed Kähler class α_Y . By definition a Kähler metric ω is extremal if the symplectic gradient of its scalar curvature is real holomorphic, i.e.

$$\mathcal{L}_{\omega^{-1}(Scal(\omega))} J_Y = 0.$$

By a well-known result of Calabi [Cal85] an extremal metric needs to be invariant by a maximal torus \mathbb{K} in the reduced automorphism group $\text{Aut}_{\text{red}}(Y)$ of Y . Let us fix such a torus \mathbb{K} . For any \mathbb{K} -invariant Kähler metric ω on Y the action of \mathbb{K} on (Y, ω) is hamiltonian and we denote by μ_ω the corresponding moment map. By a result of Guillemin–Sternberg [GS82] the image of μ_ω is a convex compact polytope P in the dual of the Lie algebra \mathfrak{k} of \mathbb{K} . For any \mathbb{K} -invariant metric ω in α_Y we normalize μ_ω in such way that its image equals P . It then follows from [FM95] and [Lah19, Lemma 1] that a \mathbb{K} -invariant metric $\omega \in \alpha_Y$ is extremal if and only if

$$Scal(\omega) = l_{\text{ext}}(\mu_\omega),$$

where l_{ext} is the unique affine extremal function such that the Futaki invariant vanishes

$$(3.8) \quad \mathbf{F}(l) := \int_Y l(\mu_\omega)(Scal(\omega) - l_{\text{ext}}(\mu_\omega)) \frac{\omega^n}{n!} = 0,$$

for any $l \in \text{Aff}(\mathfrak{k}^*)$.

We now suppose that $(Y, J_Y, \omega_Y, \mathbb{T})$ is a semisimple principal toric fibration with fiber $(X, J_X, \omega_X, \mathbb{T})$ and that $\alpha_Y := [\omega_Y]$ is a compatible Kähler class on Y . Let P denotes the Delzant polytope associated to $(X, J_X, \omega_X, \mathbb{T})$. In general, \mathbb{T}_Y is not maximal in $\text{Aut}_{\text{red}}(Y)$. However, by [ACGTF11, Proposition 1], any compatible Kähler metric (3.2) is invariant by a maximal torus $\mathbb{K}_Y \subset \text{Aut}_{\text{red}}(Y)$ containing \mathbb{T}_Y such that the following exact sequence holds

$$(3.9) \quad \{0\} \longrightarrow \mathfrak{t} \longrightarrow \mathfrak{k} \longrightarrow \mathfrak{k}_B \rightarrow \{0\},$$

where $\mathfrak{t} := \text{Lie}(\mathbb{T}_Y)$, $\mathfrak{k} := \text{Lie}(\mathbb{K}_Y)$ and $\mathfrak{k}_B := \text{Lie}(\mathbb{K}_B)$ where $\mathbb{K}_B \subset \text{Aut}_{\text{red}}(B)$ is a maximal torus such that $\omega_B := \sum_{a=1}^k \omega_a$ is \mathbb{K}_B -invariant (without loss of generality by Lichnerowicz–Matsushima Theorem).

By construction (3.1), there is an embedding of the space of \mathbb{T} -invariant smooth functions $\mathcal{C}_{\mathbb{T}}^\infty(X)$ on X in the space of \mathbb{T} -invariant smooth functions $\mathcal{C}_{\mathbb{T}}^\infty(Y)$ on Y

$$(3.10) \quad \mathcal{C}_{\mathbb{T}}^\infty(X) \subset \mathcal{C}_{\mathbb{T}}^\infty(Y).$$

Moreover, by computations in [ACG06, p. 380], the scalar curvature $Scal(\omega_Y)$ of compatible Kähler metric ω_Y on Y corresponding to a Kähler metric ω_X on X belongs to $\mathcal{C}_{\mathbb{T}}^\infty(X) \subset \mathcal{C}_{\mathbb{T}}^\infty(Y)$ and is equal to

$$(3.11) \quad Scal(\omega_Y) = \sum_{a=1}^k \frac{s_a}{p_a \circ \mu_{\omega_X} + c_a} + \frac{1}{v(\mu_{\omega_X})} Scal_v(\omega_X),$$

where s_a are the constant scalar curvatures of ω_a and $n_a := \dim(B_a)$, $Scal_v(\omega_X)$ is the v -weighted scalar curvature of ω_X (see Definition 3.2) with respect to the weight $v(x) := \prod_{a=1}^k (p_a(x) + c_a)^{n_a}$. It then follows from (3.9) and (3.11) that (3.8) reduces to an equation on X and is equivalent to

$$\int_X \left(Scal_v(\omega_X) - w(\mu_{\omega_X}) \right) l(\mu_{\omega_X}) \omega_X^\ell = 0,$$

for every $l \in \text{Aff}(\mathfrak{t}^*)$, where the moment maps μ_{ω_X} are normalized such that $\mu_{\omega_X}(X) = P$ and $w(x) := (l_{\text{ext}}(x) - \sum_{a=1}^k \frac{s_a}{p_a(x) + c_a})v(x)$. To summarize we have the following (see [ACGTF11, Section 3.5] or [AJL22, Lemma 5.14] for more details).

PROPOSITION 3.7. — *The affine extremal function l_{ext} of a compatible Kähler class belongs to the image of the map $\text{Aff}(\mathfrak{t}^*) \hookrightarrow \text{Aff}(\mathfrak{k}^*)$ induced by (3.9).*

Remark 3.8. — It follows from Proposition 3.7, [AJL22, (27)] and [Lah19, Lemma 15] that the Futaki invariant (3.8) restricted to $\text{Aff}(\mathfrak{t}^*) \subset \text{Aff}(\mathfrak{k}^*)$ coincides, up to a positive multiplicative constant, to the weighted Donaldson–Futaki invariant (2.1) for the weights (3.12).

We now state the main existence result of this section, which is one of the main results of [Jub21].

THEOREM 3.9 ([Jub21, Theorem 3]). — *Let $(Y, J_Y, \omega_Y, \mathbb{T})$ be a semisimple principal bundle with Kähler toric fiber $(X, J_X, \omega_X, \mathbb{T})$ and denote by P its moment polytope. Then there exists an extremal Kähler metric in $[\omega_Y]$ if and only if P is (v, w) -uniformly K -stable, where*

$$(3.12) \quad \begin{aligned} v(x) &= \prod_{a=1}^k (p_a(x) + c_a)^{n_a} \\ w(x) &= \left(l_{\text{ext}}(x) - \sum_{a=1}^k \frac{s_a}{p_a(x) + c_a} \right) v(x), \end{aligned}$$

and l_{ext} is the unique affine function such that (2.3) holds for (v, w) . Equivalently, there exists a (v, w) -cscK metric in $[\omega_X]$.

Proof. — We only sketch the proof of the direction “ (v, w) -uniform K -stability implies existence of a (v, w) -weighted cscK metric” which is key in the present paper, and refer to the original paper [Jub21] for details and the converse direction. We fix the weights (v, w) given by (3.12). By Proposition 3.7 and (3.11), compatible extremal metrics on Y correspond to (v, w) -cscK metrics on X via (3.2). Then, the existence of (v, w) -cscK metric in $[\omega_X]$ implies the existence of a (compatible) extremal metric in $[\omega_Y]$. The main ingredient is the existence result of cscK metric of Chen–Cheng [CC18, CC21a, CC21b], extended by He [He19] to the extremal case:

THEOREM 3.10 (Chen–Cheng, He [CC18, CC21a, CC21b, He19]). — *Let α be a Kähler class on a compact Kähler manifold Y and \mathbb{K} be a maximal torus in the reduced automorphism group $\text{Aut}_{\text{red}}(Y)$. Then there exists an extremal metric in α if and only if the \mathbb{K} -relative Mabuchi energy $\mathcal{M}^{\mathbb{K}}$ is $\mathbb{K}^{\mathbb{C}}$ -coercive, i.e. there exists $C > 0$ and $D > 0$ such that*

$$\mathcal{M}^{\mathbb{K}}(\varphi) \geq C \inf_{\gamma \in \mathbb{K}^{\mathbb{C}}} d_1(0, \gamma \cdot \varphi) - D,$$

for every $\varphi \in \mathring{\mathcal{K}}_{\mathbb{K}}(Y, \tilde{\omega}_0)$.

In the above statement, $\tilde{\omega}_0 \in \alpha$ is a fixed \mathbb{K} -invariant Kähler metric,

$$\mathcal{M}^{\mathbb{K}} : \mathcal{K}_{\mathbb{K}}(Y, \tilde{\omega}_0) \longrightarrow \mathbb{R}$$

is the \mathbb{K} -relative Mabuchi energy [Gua99, Mab86] defined on the space of \mathbb{K} -relative Kähler potential $\mathcal{K}_{\mathbb{K}}(Y, \tilde{\omega}_0) := \{\varphi \in \mathcal{C}_{\mathbb{K}}^{\infty}(Y) \mid \tilde{\omega}_{\varphi} := \tilde{\omega}_0 + dd^c\varphi > 0\}$, $\mathring{\mathcal{K}}_{\mathbb{K}}(Y, \tilde{\omega}_0) \subset \mathcal{K}_{\mathbb{K}}(Y, \tilde{\omega}_0)$ is the space of normalized potential (with respect to the vanishing of the Aubin–Mabuchi functional), d_1 is the Darvas distance [Del88] and $\mathbb{K}^{\mathbb{C}} := \mathbb{K} \otimes \mathbb{C}$ is the complexification of \mathbb{K} . Moreover $\mathbb{K}^{\mathbb{C}}$ acts on $\mathring{\mathcal{K}}_{\mathbb{K}}(Y, \tilde{\omega}_0)$ via the natural action on \mathbb{K} -invariant Kähler metrics in $[\tilde{\omega}_0]$. Originally [He19], the coercivity condition in Theorem 3.10, was expressed in term of the complexification of a maximal compact connected subgroup of $\text{Aut}_{\text{red}}(Y)$ (not necessarily commutative). As observed in [Jub21, Section 5], the same arguments as in [CC18, He19] provide this statement.

The proof of Theorem 3.9 is divided in two steps:

- (1) show that the (v, w) -uniform stability with respect to the weights (3.12) of the polytope P implies the $\mathbb{T}^{\mathbb{C}}$ -coercivity of the *weighted Mabuchi energy* $\mathcal{M}_{v,w}$ corresponding to $(X, J_X, [\omega_X], \mathbb{T})$;
- (2) adapt the continuity path (3.13) involved in the proof of Cheng–Cheng and He to obtain the existence of a compatible extremal metric in $[\omega_Y]$ (or equivalently a (v, w) -weighted cscK metric in $[\omega_X]$).

The weighted Mabuchi energy $\mathcal{M}_{v,w}$ mentioned above is the one introduced by Lahdili [Lah19].

Step 1. — The first step is an adaptation of [Don02, ZZ08] which established this result in the unweighted case, i.e. when $v = 1$ and $w = l_{\text{ext}}$. Fix $\omega_0 \in [\omega_X]$ a \mathbb{T} -invariant Kähler metric. On a toric Kähler manifold, a \mathbb{T} -invariant Kähler metric $\omega \in [\omega_0]$ can be defined via two functions: a \mathbb{T} -invariant ω_0 -relative Kähler potential $\varphi \in \mathcal{K}_{\mathbb{T}}(X, \omega_0)$ and a symplectic potential $u \in \mathcal{S}(P, \mathbf{L})$, which, by definition, $\mathcal{S}(P, \mathbf{L})$ is the space of smooth strictly convex functions on P^0 which satisfy the so-called *Abreu boundary conditions* [Abr03, Gui94]. There is a well-known correspondence between a symplectic and a Kähler potential defining the same Kähler metric [ACG06, ACGTF04, Don02, Gui94]. Via this correspondence we can consider the weighted Mabuchi energy $\mathcal{M}_{v,w}$ as a functional on $\mathcal{S}(P, \mathbf{L})$. A first step is to show, as in the case $v = 1$ [Don02, Proposition 3.3.4], that $\mathcal{M}_{v,w}$ extends to the space $\mathcal{CV}^{\infty}(P)$ of smooth convex functions on P^0 and continuous on P , and that a symplectic potential $u \in \mathcal{S}(P, \mathbf{L})$ defining a (v, w) -weighted cscK metric realizes the minimum of $\mathcal{M}_{v,w}$ over $\mathcal{CV}^{\infty}(P)$. This is proven in [Jub21, Proposition 7.7]. The idea is then, as in [ZZ08], to compare $\mathcal{M}_{v,w}$ and \mathcal{M}_{v,w_0} , where the weight w_0 is the v -weighted

scalar curvature $Scal_v(u_0)$ of (the Kähler metric defined by) any fixed symplectic potential u_0 . Then u_0 trivially solves

$$Scal_v(u) = w_0, \quad u \in \mathcal{S}(P, \mathbf{L}),$$

In other words, there exists a (v, w_0) -cscK metric in $[\omega_X]$. We deduce that \mathcal{M}_{v, w_0} is bounded from below on $\mathcal{CV}^\infty(P)$ by the discussion above. Consequently, by comparing $\mathcal{M}_{v, w}$ and \mathcal{M}_{v, w_0} and using our hypothesis, we can show that [Jub21, Proposition 7.9] (or [ZZ08] for $v = 1$)

$$\mathcal{M}_{v, w}(u) \geq C \|u^*\|_{L_1} - D,$$

for any $u \in \mathcal{S}(P, \mathbf{L})$. There $D > 0$ and $C > 0$ are uniform in u and $\|\cdot\|_{L_1}$ is the L_1 -norm on P with respect to the Lebesgue measure. Denote by d_1^X the Darvas distance on $\mathcal{K}_{\mathbb{T}}(X, \omega_0)$. For a (normalized) Kähler potential $\varphi \in \mathcal{K}_{\mathbb{T}}(X, \omega_0)$ corresponding to (normalized) symplectic potential $u \in \mathcal{S}(P, \mathbf{L})$ via the correspondence described above, we can show that $d_1^X(0, \varphi) \leq A\|u\| + B$, for some positive constant A and B . We deduce that

$$\mathcal{M}_{v, w}(\varphi) \geq C' \inf_{\gamma \in \mathbb{T}^c} d_1^X(0, \gamma \cdot \varphi) - D'.$$

for any φ (normalized) \mathbb{T} -invariant Kähler potential, where C' and D' are positive constant. We refer to [Jub21, Section 7.6] for the precise normalization of symplectic and Kähler potentials.

Step 2. — The proof of the direction “coercivity implies existence” in Theorem 3.10 is based on the resolution of the following continuity path

$$(3.13) \quad t(Scal(\tilde{\omega}_\varphi) - l_{ext}(\mu_{\tilde{\omega}_\varphi})) = (1 - t)(\Lambda_{\tilde{\omega}_\varphi}(\tilde{\chi}) - n - \ell),$$

where $t \in [0, 1]$, $\tilde{\chi}$ is a fixed Kähler metric in $[\tilde{\omega}_0]$, $\dim(Y) = n + \ell$, $\varphi \in \mathcal{K}_{\mathbb{K}}(Y, \tilde{\omega}_0)$ and $\Lambda_{\tilde{\omega}_\varphi}(\tilde{\chi})$ is the symplectic trace of $\tilde{\chi}$ with respect to $\tilde{\omega}_\varphi$. Chen–Cheng and He showed that there exists $t_0 \in (0, 1)$ such that

$$\tilde{S} := \{t \in [t_0, 1] \text{ such that } \exists \varphi_t \in \mathcal{K}_{\mathbb{K}}(Y, \tilde{\omega}_0) \text{ solution of (3.13)}\}$$

is non-empty, open and closed in $[t_0, 1]$.

We come back to the case of semisimple principal toric fibration. We fix a \mathbb{T} -invariant Kähler metric $\omega_0 \in [\omega_X]$ and the induced compatible Kähler metric $\tilde{\omega}_0 \in [\omega_Y]$, see (3.2). It follows from (3.10) and (3.9) (see [ACGTF11, Lemma 7] for a proof) that

$$(3.14) \quad \mathcal{K}_{\mathbb{T}}(X, \omega_0) \subset \mathcal{K}_{\mathbb{K}}(Y, \tilde{\omega}_0),$$

where \mathbb{K} is a maximal torus in $\text{Aut}_{\text{red}}(Y)$ containing \mathbb{T} such that (3.9) holds. Moreover, for every $\varphi \in \mathcal{K}_{\mathbb{T}}(X, \omega_0)$, seen as function on Y , the Kähler metric $\tilde{\omega}_\varphi := \tilde{\omega}_0 + dd^c\varphi$ is a compatible Kähler metric in $[\omega_Y]$ in the sense of Definition 3.4. Also, $\tilde{\omega}_\varphi$ on Y is the metric induced by ω_φ on X via (3.2) (see [ACGTF11, Lemma 7] or [AJL22, Lemma 5.5]). Following [ACGTF11], we then refer to the image of (3.14) as the space of *compatible Kähler potentials*. Additionally, (3.11), (3.14), [AJL22, (27)] and Proposition 3.7 (see also [AJL22, Lemma 5.10]) show that

$$(3.15) \quad \mathcal{M}^{\mathbb{K}}|_{\mathcal{K}_{\mathbb{T}}(X, \omega_0)} = \mathcal{M}_{v, w}.$$

Let d_1^Y be the Darvas distance on $\mathcal{K}_{\mathbb{K}}(Y, \tilde{\omega}_0)$ [Del88]. By [AJL22, Corollary 6.5] and since $v > 0$,

$$(3.16) \quad d_1^X(0, \varphi) \geq A d_1^Y(0, \varphi)$$

for any $\varphi \in \mathcal{K}_{\mathbb{T}}(X, \omega_0) \subset \mathcal{K}_{\mathbb{K}}(Y, \tilde{\omega}_0)$, where $A > 0$. By (3.11), the restriction of (3.13) to the space of compatible Kähler potentials $\mathcal{K}_{\mathbb{T}}(X, \omega_0) \subset \mathcal{K}_{\mathbb{K}}(Y, \tilde{\omega}_0)$ is equal to

$$(3.17) \quad t(\text{Scal}_v(\omega_\varphi) - w(\mu_{\omega_\varphi})) = (1 - t)(\Lambda_{\tilde{\omega}_\varphi}(\tilde{\chi}) - n - \ell),$$

We now want to show that there exists $t_0 \in (0, 1)$ such that

$$S := \{t \in [t_0, 1] \text{ such that } \exists \varphi_t \in \mathcal{K}_{\mathbb{T}}(X, \omega_0) \subset \mathcal{K}_{\mathbb{K}}(Y, \tilde{\omega}_0) \text{ solution of (3.17)}\}$$

is non-empty, open and closed in $[t_0, 1]$. Since any $\mathbb{T}^{\mathbb{C}}$ -orbit is included in a $\mathbb{K}^{\mathbb{C}}$ -orbit, the $\mathbb{T}^{\mathbb{C}}$ -coercivity is stronger than the $\mathbb{K}^{\mathbb{C}}$ -coercivity. Then, by (3.15), (3.16) and *Step 1*, $\mathcal{M}^{\mathbb{K}}$ is $\mathbb{K}^{\mathbb{C}}$ -coercive on the space of compatible potentials $\mathcal{K}_{\mathbb{T}}(X, \omega_0)$ suitably normalized. As observed in [Jub21, Lemma 6.3], we can choose $\tilde{\chi}$ such that (3.17) is an equation on X and admits a solution $\varphi_{t_0} \in \mathcal{K}_{\mathbb{T}}(X, \omega_0)$ for some $t_0 \in (0, 1)$, showing that S is non-empty. The openness follows from an application of the Implicit Function Theorem [Jub21, Proposition 6.4]. From the *openness*, the closedness of $\mathcal{K}_{\mathbb{T}}(X, \omega_0)$ in $\mathcal{K}_{\mathbb{K}}(Y, \tilde{\omega}_0)$ (see [Jub21, Section 3.4]) and *Step 1*, following the proof of Theorem 3.10, we obtain a sequence of compatible Kähler metric $\tilde{\omega}_{\varphi_j}$ such that $\bar{\omega}_j := \gamma_j^*(\tilde{\omega}_{\varphi_j})$, $\gamma_j \in \mathbb{K}^{\mathbb{C}}$, converge to an extremal metric $\bar{\omega}_1$. The space of (normalized) compatible Kähler potential $\mathring{\mathcal{K}}_{\mathbb{T}}(X, \omega_0)$ is not stable under the action of $\mathbb{K}^{\mathbb{C}}$, then either $\bar{\omega}_j$ or $\bar{\omega}_1$ is compatible in general. However, we can show (see [Jub21, Proof of Proposition 6.5]) that $\bar{\omega}_1$ is of the form of (3.2), for a possibly different principal connection θ and Kähler metrics ω_a . Moreover, we can argue that the Kähler metric $\omega_1 \in [\omega_X]$ defining $\bar{\omega}_1$ via (3.2) is a weighted (v, w) -cscK metric for the same weights (3.12), see [Jub21, Proof of Proposition 6.5]. \square

Note that condition (2.3) corresponds to the vanishing of the modified Futaki character, and l_{ext} encodes the extremal vector field. In particular the extremal metric above is cscK if and only if l_{ext} is constant.

Remark 3.11. — It is remarkable that the condition depends on the base only through the constants (s_a) and the existence of a principal \mathbb{T} -bundle with connection with corresponding data (p_a) . In particular, when we obtain an existence result for extremal Kähler metrics, we usually actually obtain the existence of extremal Kähler metrics over a full deformation family of cscK manifolds. This is exactly this fact which is used in the Proof of Proposition 5.1 to obtain an existence condition of extremal metric on $\mathbb{C}\mathbb{P}^2$ -bundles over any Kähler–Einstein Fano threefold depending only on the cohomology class and the degrees of the line bundles.

4. Geometric applications of the sufficient condition

4.1. The general statement

We use the same notations as in Section 3 for semisimple principal toric fibrations and the same notations as in Section 2 for the decomposition of polytopes.

COROLLARY 4.1 (of Theorem 2.6). — *The semisimple principal toric fibration (Y, ω_Y) admits an extremal Kähler metric in $[\omega_Y]$ if there exists an $x_0 \in P^0$ and corresponding cone decomposition $P = \cup_j P_j$ such that for all j and for all $x \in P_j$*

$$\frac{1}{L_j(x_0)} \left(\ell + 1 + \sum_{a=1}^k \frac{n_a p_a(x - x_0)}{p_a(x) + c_a} \right) - \frac{1}{2} \left(l_{\text{ext}}(x) - \sum_{a=1}^k \frac{s_a}{p_a(x) + c_a} \right) \geq 0.$$

Proof. — By Theorem 3.9, the sufficient condition of Theorem 2.6 translates as a sufficient condition of existence of extremal Kähler metrics. To obtain the statement above, it suffices to note that for the weight v involved, we have

$$d_x v(y) = \left(\sum_{a=1}^k \frac{n_a p_a(y)}{p_a(x) + c_a} \right) v(x).$$

so that in the condition in Theorem 2.6, we can factor by $v(x)$ which is positive everywhere. □

4.2. Fibrations with Fano fiber

We now turn to the fibrations with Fano fiber, in order to use Corollary 2.10. With the same notations as in Section 3.4, we now assume furthermore that the toric fiber is a Fano manifold, and that the Kähler class $[\omega_X]$ is a multiple of the anticanonical class $2\pi c_1(X)$. As a consequence, the moment polytope P is a dilation of a reflexive lattice polytope. This implies that the labelled polytope (P, \mathbf{L}) corresponding to the lattice polytope P is monotone, with a preferred point x_0 and $L_1(x_0) = \dots = L_d(x_0) = t$. Assuming without loss of generality that the (anti-)canonical normalization is used for the moment polytope of the fiber, we may further assume that $x_0 = 0$, and $t = \frac{[\omega]}{2\pi c_1(X)}$.

COROLLARY 4.2. — *The semisimple principal toric fibration $(Y, [\omega_Y])$ with Fano toric fiber admits an extremal Kähler metric in $[\omega_Y]$ if $\forall x \in P$,*

$$(4.1) \quad 2 \left(\ell + \sum_a n_a \right) + 2 + \sum_a \frac{t s_a - 2n_a c_a}{p_a(x) + c_a} - t l_{\text{ext}}(x) \geq 0$$

Note that $\ell + \sum_a n_a = \dim(Y)$.

Proof. — Since all $L_j(x_0)$ are equal to t , the condition from Corollary 4.1 further simplifies to

$$2\ell + 2 + \sum_{a=1}^k \frac{2n_a p_a(x) + t s_a}{p_a(x) + c_a} - t l_{\text{ext}}(x) \geq 0 \quad \forall x \in P$$

as for Corollary 2.10. Writing $2n_a p_a(x) = 2n_a(p_a(x) + c_a) - 2n_a c_a$ yields the statement. □

While simple enough, and tractable with numerical optimization techniques, the inequality involved is a polynomial inequality in several variables, whose degree can be equal to the dimension of the basis plus one. It is difficult to solve formally, but there is a further reduction that allows to get a simpler condition which can be checked by a finite number of evaluations of polynomial functions.

COROLLARY 4.3. — Assume furthermore that for all a , $c_a \geq \frac{ts_a}{2n_a}$. Then the semisimple principal toric fibration $(Y, [\omega_Y])$ admits an extremal Kähler metric in $[\omega_Y]$ if inequation (4.1) is satisfied at every vertex of P .

Proof. — The inverse of an affine function is convex on the locus where this affine function is positive. Hence under the condition in the statement, the function $\frac{ts_a - 2n_a c_a}{p_a + c_a}$ is concave on P . Condition (4.1) thus amounts to checking the non-negativity of a concave function on a convex polytope: it is enough to check the non-negativity on vertices. \square

Remark 4.4. — In the case of a *simple* principal toric fibration, that is, if there is only one factor in the basis, then the condition becomes extremely simple for classes with $c_a \geq \frac{ts_a}{2n_a}$: it is enough to check a degree two polynomial inequation on every vertex of the moment polytope. This is actually used in most of the examples of Section 5, see e.g. Proposition 5.1 or Proposition 5.3.

Remark 4.5. — We can write a similar statement for the general case of toric fibrations, by working on the cone decomposition. In that case the conditions to impose are: for all j , for all a , $L_j(x_0)s_a - 2n_a(p_a(x_0) + c_a) \leq 0$ and condition (4.1) is satisfied at all vertices of P_j , that is, some vertices of P and x_0 .

4.3. Extremal metrics in the anticanonical class

An important special case when the toric fiber is Fano is given by the semisimple principal toric fibrations which are themselves Fano. By our general sufficient condition, we obtain a very simple condition for the existence of extremal Kähler metrics on Fano toric fibrations.

COROLLARY 4.6. — A Fano semisimple principal toric fibration Y admits an extremal Kähler metric in $c_1(Y)$ if its extremal function l_{ext} satisfies:

$$(4.2) \quad \sup l_{\text{ext}} \leq 2(\dim(Y) + 1)$$

Proof. — By Lemma 3.6, for all a , $s_a = 2n_a c_a$ and the condition from Corollary 4.2 becomes

$$2 \dim(Y) + 2 - l_{\text{ext}} \geq 0 \quad \text{on } P \quad \square$$

Of course, as in Corollary 4.3, it is enough to check this condition on vertices of the polytope. Furthermore, if l_{ext} is constant, it is equal to $2 \dim(Y)$ since the class is the anticanonical one. As a consequence, the condition is strictly satisfied:

$$(4.3) \quad 2 \dim(Y) + 2 - l_{\text{ext}} = 2 > 0.$$

In particular, we recover that a Fano toric fibration with vanishing Futaki invariant admits a Kähler-Einstein metric :

PROPOSITION 4.7 ([AJL22]). — Let (Y, ω_Y) be a Fano semisimple principal toric fibration with vanishing Futaki invariant. Then there exists a Kähler-Einstein metric in $2\pi c_1(Y)$.

More interestingly, we have the following consequence.

PROPOSITION 4.8. — *Let (Y, ω_Y) be a Fano semisimple principal fibration. Then on a neighborhood of the anticanonical class, a compatible Kähler class admits a cscK metric if and only if its Futaki invariant vanishes.*

Proof. — By Remark 2.8, the non-negativity condition in Theorem 2.6 varies continuously with the weight. By equation (4.3), that condition is strictly satisfied at the anticanonical class, so it is satisfied on a neighborhood of this class. The only added assumption in Theorem 2.6 translates as vanishing of the Futaki invariant. \square

Remark 4.9. — Of course, if the Futaki invariant of the anticanonical class vanishes, this is already known by Lebrun–Simanca [LS93] and [AJL22]. Similarly, if the anticanonical class is strictly K-unstable, nearby classes will be as well. However, in the present setting, working directly with the condition it is not hard to find an explicit neighborhood which works. Furthermore, the statement applies even when we do not know whether there exists an extremal Kähler metric or not in the anticanonical class. In the current state of knowledge, it could also happen that the anticanonical class is K-semistable (for the notion of relative K-stability adapted to extremal Kähler metrics), and the above proposition would still apply in that case. This is a further illustration of a phenomenon observed in [Del22].

4.4. Weighted solitons on Fano semisimple principal toric fibrations

Recall a v -soliton is a Kähler metric ω such that

$$\text{Ric}(\omega) - \omega = \frac{1}{2} dd^c \log(v),$$

where $\text{Ric}(\omega)$ is the Ricci form of ω . On Fano semisimple principal toric fibrations Y with fiber X , a Kähler metric $\omega_Y \in 2\pi c_1(Y)$ is a v -soliton if and only if its corresponding metric $\omega_X \in 2\pi c_1(X)$ is $(v v_0, \tilde{v})$ -cscK (see [AJL22, Lemma 2.2, Lemma 5.11]) for the weights $\tilde{v} := 2(\ell_{v_0}(x)v(x) + d_x(v_0 v)(x))$ and v_0 is defined in (3.12). Since the polytope must be reflexive hence monotone, one has $x_0 = 0$, $t = 1$ and condition (2.9) becomes $v \geq 0$ on the polytope, which is obviously satisfied. Moreover, by [Jub21, Proposition 7.8], the (v, \tilde{v}) -uniform K-stability implies the coercivity of the corresponding weighted (v, \tilde{v}) -Mabuchi functional. Thanks to [LH20, Theorem 3.5], we obtain the existence of a weighted soliton in $2\pi c_1(Y)$. This result was obtained in [AJL22, Theorem 3], by applying K-stability arguments from [LH20]. The proof proposed above allows to remain fully on the differential geometric side.

COROLLARY 4.10 ([AJL22, Theorem 3]). — *Let Y be a Fano semisimple principal toric fibration with associate Delzant polytope P . Consider the weighted Donaldson–Futaki invariant \mathcal{F} for the weights corresponding to v -solitons defined above. Then, if \mathcal{F} vanishes, there exists a v -soliton in $2\pi c_1(Y)$.*

5. Examples

5.1. Examples of bases

In this section, we comment on examples of possible bases for the semisimple principal toric fibration construction. This allows to determine possible values of

s_a to plug into the condition. The easiest way to get a cscK basis is to choose a Kähler–Einstein manifold, equipped with a multiple of its first Chern class when it is definite, and with an arbitrary Kähler class for Calabi–Yau manifolds.

For canonically polarized manifolds, there always exists a Kähler–Einstein metric in $-2\pi c_1(X)$, and there exists such manifolds in every dimension. In particular, the value $s_a = -\frac{2n_a}{k_a}$ are always allowed, for $k_a \in \mathbb{Z}_{>0}$. For manifolds with zero first Chern class, there always exist Kähler–Einstein metrics with zero scalar curvature. For the positive curvature case, since the projective space of dimension n is a Kähler–Einstein manifold of index $n+1$, all the values $s_a = 2\frac{n_a(n_a+1)}{k_a}$ are allowed, for $k_a \in \mathbb{Z}_{>0}$. More generally, for a Kähler–Einstein Fano basis of dimension n_a and index I_a , then all the values $s_a = 2\frac{n_a I_a}{k_a}$ are allowed, for $k_a \in \mathbb{Z}_{>0}$. Note that the Fano index of an n -dimensional Fano manifold is always an integer between 1 and $n+1$. Here are a couple known results on existence of Fano Kähler–Einstein manifolds when n is small or I is large:

- if $I = n + 1$ then $X = \mathbb{P}^n$ is the n -dimensional projective space, and it is Kähler–Einstein,
- if $I = n$ then $X = Q^n$ is the n -dimensional quadric, and it is Kähler–Einstein,
- if $n = 1$ then $X = \mathbb{P}^1$, $I = 2$ and it is Kähler–Einstein,
- if $n = 2$ then \mathbb{P}^2 (index 3), $X = \mathbb{P}^1 \times \mathbb{P}^1$ (index 2) and the blowups of \mathbb{P}^2 (index 1) at three or more points are Kähler–Einstein,
- if $n = 3$, then the existence of Kähler–Einstein metrics on a general member of a deformation family of smooth Fano threefolds was recently settled in [ACC⁺23], and the families where the general member is *not* Kähler–Einstein are the following, in the labelling used in [ACC⁺23], 2.23, 2.26, 2.28, 2.30, 2.31, 2.33, 2.35, 2.36, 3.14, 3.16, 3.18, 3.21, 3.22, 3.23, 3.24, 3.26, 3.28, 3.29, 3.30, 3.31, 4.5, 4.8, 4.9, 4.10, 4.11, 4.12, 5.2.

5.2. \mathbb{P}^2 -fiber over Fano threefold

We consider the 2-dimensional projective space $(\mathbb{P}^2, \mathbb{T}^2, 2\pi c_1(\mathbb{P}^2))$. Identifying the lattice Λ of \mathbb{T}^2 with \mathbb{Z}^2 , we consider its labelled moment polytope (P, \mathbf{L}) in \mathbb{R}^2

$$(5.1) \quad P = \left\{ (x_1, x_2) =: x \in \mathbb{R}^2 \mid L_1(x) \geq 0, L_2(x) \geq 0, L_3(x) \geq 0 \right\},$$

where $L_1(x) := x_1 + 1$, $L_2(x) := x_2 + 1$, $L_3(x) := -x_1 - x_2 + 1$. Let (B, ω_B) be a KE Fano threefold with $\alpha_B := [\omega_B]$ primitive element of $H^2(B, \mathbb{Z})$ proportional to the first Chern class $2\pi c_1(B)$. Let $\mathcal{L}_i \rightarrow B$ be a holomorphic line bundle of degree $-p_i$ proportional to the anticanonical line bundle $-K_B$, i.e. $-p_i \alpha_B = 2\pi c_1(\mathcal{L}_i)$. We consider a *simple* principal toric fibration (i.e. the basis has only one factor) $\pi : Y := \mathbb{P}(\mathcal{L}_0 \oplus \mathcal{L}_1 \oplus \mathcal{L}_2) \rightarrow B$. Since the holomorphic class of Y is invariant by tensoring $E := \mathcal{L}_0 \oplus \mathcal{L}_1 \oplus \mathcal{L}_2$ with a line bundle, we can suppose without loss of generality that $\mathcal{L}_0 = \mathcal{O}$ is the trivial line bundle and $p_i \geq 0$, $i = 1, 2$. When B is a local Kähler product of nonnegative cscK metric and $p_1 = p_2 > 0$ or $p_2 > p_1 = 0$, it is known [ACGTF08, Proposition 11], that there exists an extremal metric in every compatible Kähler classes. We then suppose $p_2 \geq p_1 > 0$.

The compatible Kähler classes are parametrized by constants c and are of the form

$$(5.2) \quad \alpha_c := 2\pi c_1(O_E(3)) + c\pi^*(\alpha_B),$$

As introduced in Section 5.1, since B is a Fano threefold, the only possible Fano indices I are 1, 2, 3 or 4. In the case where B is the quadric Q_3 or the projective space \mathbb{P}^3 (i.e. if $I = 3$ or $I = 4$ respectively), Leray–Hirsch Theorem shows that $H^2(Y, \mathbb{R}) \cong \mathbb{R}^2$. It follows that, up to scaling, all Kähler classes are compatible, i.e. of the form of (5.2). It is known that [ACGTF11, Theorem 4] for c sufficiently large, the class α_c is extremal. The following Proposition gives a precise value for c , depending on p_1 and p_2 , from which α_c admits an extremal metric.

PROPOSITION 5.1. — *Let $Y = \mathbb{P}_B(\mathcal{O}_B \oplus H^{-p_1} \oplus H^{-p_2})$, where B is a Kähler–Einstein Fano threefold, H is the smallest integral divisor of $2\pi c_1(B)$ and $1 \leq p_1 \leq p_2$. Then there exists an extremal metric in α_c for $c \geq 7p_2$.*

Proof. — Since the arguments are identical for each Fano index I , we give the proof only for $I = 4$.

By Corollary 4.3, for $c \geq 4$, it is sufficient to check (4.1) evaluated in each vertex $v_1 := (-1, 2)$, $v_2 := (-1, -1)$, $v_3 := (2, -1)$ of the polytope P .

Using Program 2 in Appendix A, we find that the LHS of (4.1) evaluated in v_1 is a rational fraction in the variables c, p_1, p_2 :

$$\text{LHS of (4.1)} = \frac{P(c, p_1, p_2)}{Q(c, p_1, p_2)}.$$

We give the explicit expression of the polynomials P and Q in Appendix B. Suppose now $c \geq 7p_2$ and $p_2 \geq p_1 \geq 1$. Then we can find two polynomials

$$R(c) := 12250c^{10} - 73500c^9 - 295470c^8 + 1296540c^7 - 3657150c^6 + 3776220c^5 - 6537672c^4 + 5624964c^3 - 6193584c^2 + 85920232c - 1889568$$

and

$$S(c) := 6125c^{10} + 18375c^9 + 6615c^8 + 19845c^7 + 127575c^6 + 382725c^5 + 17496c^4 + 52488c^3 - 288684c^2 - 866052c$$

such that

$$0 < R(c) \leq P(c, p_1, p_2)$$

and

$$0 < S(c) \text{ and } S(c) \geq Q(c, p_1, p_2).$$

It implies that

$$\text{LHS of (4.1)} = \frac{P(c, p_1, p_2)}{Q(c, p_1, p_2)} \geq \frac{R(c)}{S(c)} \geq 0.$$

We proceed analogously for the vertex v_2 and v_3 . We conclude the proof by involving Corollary 4.3. □

Remark 5.2. — In Proposition 5.1, we obtain a lower bound on c depending only on p_1 and p_2 . For given values of p_1 and p_2 it is possible to obtain a more optimal result. Indeed, suppose p_1 and p_2 are fixed. Then, the LHS of (4.1) is a rational fraction F depending only on the variable c . We then only need to look for constant α

such that F is non-negative for $c \geq \alpha$. For example, if $B = \mathbb{P}^3$, respectively $B = Q_3$, $p_1 = 1$ and $p_2 = 2$, (4.1) show the existence of an extremal metric in α_c for $c \geq 7.09$, respectively $c \geq 9.08$. We refer to Appendix A for further examples of application of the sufficient condition on simple principal \mathbb{P}^2 -fibrations.

5.3. Comments on the rank one case

5.3.1. Varying x_0 and prescribing weighted scalar curvature on \mathbb{P}^1

As noted in Remark 2.7, it can be useful to vary the base point x_0 . In this short paragraph, we want to illustrate this phenomenon in the simplest possible case, that is, when working on the one-dimensional polytope $[-1, 1] \subset \mathbb{R}$ with the weights $v \equiv 1$ and arbitrary w . We further choose the lattice labelling of $[-1, 1]$ induced by the lattice $\mathbb{Z} \subset \mathbb{R}$ (in other word, we work on the anticanonical moment polytope of \mathbb{P}^1). More precisely, the labelling (L_1, L_2) is given by $L_1(x) = 1 + x$ and $L_2(x) = 1 - x$. Since $v \equiv 1$, we have $d_x v \equiv 0$, hence condition (2.6) from Theorem 2.6 translates as

$$\frac{1}{4}w|_{[-1, x_0]} \leq \frac{1}{1+x_0} \quad \text{and} \quad \frac{1}{4}w|_{[x_0, 1]} \leq \frac{1}{1-x_0}$$

The latter condition is illustrated in Figure 5.1, and it is obviously less restrictive if one can choose x_0 than the uniform condition corresponding to the obvious choice of $x_0 = 0$ for the monotone lattice polytope $[-1, 1]$.




Figure 5.1. Varying x_0

We end this paragraph by recalling that $(1, w)$ -uniform stability of the lattice polytope $[-1, 1]$ translates to existence of certain canonical Kähler metrics on \mathbb{P}^1 thanks to [LLS21].

5.3.2. Extremal metrics on \mathbb{P}^1 -bundles

We have focused on applications of our sufficient condition to semisimple principal toric bundles with dimension two toric fiber. This is because in the case of a one-dimensional toric fiber, quite a few strong results have been shown in [ACGTF08].

For example, it is proved in [ACGTF08, Proposition 11] that if all factors (B_a, ω_a) of the basis have non-negative constant scalar curvature, and the fiber is one-dimensional, then there exists an extremal Kähler metric in all compatible Kähler classes.

There cannot be such a result if some factors of the basis have negative constant scalar curvature, as shown by examples in [ACGTF08]. More importantly, some of these examples motivated the initial introduction of the notion of uniform K-stability, as they are likely relatively K-polystable in the sense of [Szé07], but do not admit extremal Kähler metrics.

On the positive side, by [ACGTF08, Theorem 1], there always exist extremal Kähler metrics on a semisimple principal \mathbb{P}^1 -fibration, when all the c_a are large enough, an example of existence of extremal Kähler metrics in an adiabatic regime for fibrations. However, it is not so easy to derive explicit Kähler classes with extremal Kähler metrics from this asymptotic proof. A possible approach to get explicit classes with extremal Kähler metrics would be to compute the extremal polynomial (in the terminology of [ACGTF08]) and check when it is positive. This is less practical than our sufficient condition, which involves only checking the positivity of a polynomial at two points. We provide in the appendix an elementary computer program which checks whether our sufficient condition is satisfied for a simple principal \mathbb{P}^1 -fibration, which could easily be adapted to the case of a semisimple principal \mathbb{P}^1 -fibration.

5.3.3. A more explicit example

Consider B a three-dimensional canonically polarized manifold, equipped with its Kähler–Einstein metric in $-2\pi c_1(X)$, whose scalar curvature is thus equal to -6 . We consider the sufficient condition for existence of extremal Kähler metrics in admissible Kähler classes on the \mathbb{P}^1 -bundles $\mathbb{P}(\mathcal{O}_B \oplus K_B^m)$. Up to rescaling and symmetry, this amounts to checking (v, w) -uniform K-stability of the reflexive lattice polytope $[-1, 1] \subset \mathbb{R}$ with respect to the weights

$$v(x) = (px + c)^3 \quad \text{and} \quad w(x) = \left(l_{\text{ext}}(x) - \frac{-6}{px + c} \right) (px + c)^3$$

where $p \in \mathbb{Q}$, $c \in \mathbb{R}$ and $c > p > 0$. Our sufficient condition allows to obtain the following explicit families of extremal Kähler classes. We only show an example with very rough estimates to illustrate the results, but of course one could get much more classes by using more precise estimates in the proof, and even more classes by using the sufficient condition in Theorem 2.6 in its full generality.

PROPOSITION 5.3. — *With the above notations, if $c \geq 15p$, then $[-1, 1]$ is (v, w) -uniformly K-stable. The corresponding Kähler classes on the \mathbb{P}^1 -bundles $\mathbb{P}(\mathcal{O} \oplus K_B^m)$ admit extremal Kähler metrics.*

Proof. — Using Program 1 in the appendix or straightforward but tedious computations, we obtain up to elementary simplifications that the sufficient condition reads as

$$75c^7 - 300c^6 - 65c^5p^2 + 160c^4p^2 - 15c^3p^4 - 180c^2p^4 - 27cp^6 + 48p^6$$

is greater than

$$\left| -75c^6p + 5c^4p^3 + 80c^3p^3 - 105c^2p^5 + 15p^7 \right|$$

Without attempting to give an optimal result, we may as well check that it is greater than

$$75c^6p + 5c^4p^3 + 80c^3p^3 + 105c^2p^5 + 15p^7$$

since c and p are positive. Writing $c = \alpha p$ for some $\alpha > 1$ and simplifying by p^6 , we get a linear inequation in p

$$(5.3) \quad pA + B \geq 0$$

where

$$A = 75\alpha^7 - 75\alpha^6 - 65\alpha^5 - 5\alpha^4 - 15\alpha^3 - 105\alpha^2 - 27\alpha - 15$$

$$B = -300\alpha^6 + 160\alpha^4 - 80\alpha^3 - 180\alpha^2 + 48$$

Since $\alpha > 1$, the coefficient A is larger than $(75\alpha - 307)\alpha^6$ and in particular, it is non-negative for $\alpha \geq \frac{307}{75}$. Using the same lower bound for the leading coefficient, inequation (5.3) is certainly satisfied at $p = 1$ if

$$(75\alpha - 307)\alpha^6 - 300\alpha^6 - 160\alpha^4 - 80\alpha^3 - 180\alpha^2 - 48 \geq 0$$

Using again $\alpha > 1$ and very rough estimates, this is implied by the inequality

$$(75\alpha - 1075)\alpha^6 \geq 0$$

The latter is satisfied at least for $\alpha \geq 15$, and since $15 \geq \frac{307}{75}$, we obtain that if $\alpha \geq 15$, the sufficient condition is satisfied for all $p \geq 1$. \square

Appendix A. An elementary Python program

We provide, as a courtesy to the reader, elementary Python programs using SymPy that checks the sufficient condition from Corollary 4.3 for *simple* principal toric fibrations (that is, the basis has only one factor) with Fano toric fiber X of dimension one or two such that $[\omega_X]$ a multiple of $2\pi c_1(X)$.

The only data from the simple principal toric bundle needed to compute the condition is:

- from the basis, the dimension $n \in \mathbb{Z}$ and scalar curvature $s \in \mathbb{Q}$
- from the Fano toric fiber of dimension $\ell \in \{1, 2\}$, the reflexive moment polytope $P \subset \mathbb{R}^\ell = \mathbb{Z}^\ell \otimes \mathbb{R}$, and the multiple $t = \frac{[\omega_X]}{2\pi c_1(X)} \in \mathbb{R}$
- the one-parameter subgroup p from the principal bundle, identified with an integer $p \in \mathbb{Z}$ if the fiber is one-dimensional, and with an element $p = (p_1, p_2) \in \mathbb{Z}^2$ if the fiber is of dimension two,
- and the constant $c \in \mathbb{R}$ defining the admissible Kähler class.

We wish to compute the expression given by the right-hand side of (4.1)

$$\text{test} = 2(\ell + n + 1) + \frac{ts - 2nc}{p(x) + c} - tl_{\text{ext}}(x)$$

in order to check the condition. For this, it suffices to compute the extremal function l_{ext} by solving the linear system which defines it. Our short programs compute l_{ext} ,

then `test`, then evaluate `test` at the vertices of P and returns the minimum if all data are explicitly given. If the minimum returned by the program is non-negative, the data correspond to a simple principal toric fibration with an admissible Kähler class and $c > \frac{ts}{2n}$, then there exists an extremal Kähler metric. We may also let some of the data remain unknown and treat them as variables.

```

1 import sympy as sym
2 # variable on the line (here the fiber is one-dimensional)
3 x = sym.symbols('x')
4 # data of the simple principal toric fibration
5 p, c = sym.symbols('p,c')
6 n, s, t = 3, -6, 1
7 # weights
8 l = c+p*x
9 v, w0 = l**n, -s*l**(n-1) # for now, unknown l_ext is replaced with
   zero
10 # Donaldson-Futaki invariant with weights (v,w0)
11 def DFO(f):
12     interior=sym.integrate(f*w0, (x, -t, t))
13     facets=(f*v).subs(x,-t)+(f*v).subs(x,t)
14     return(interior+facets)
15 # Compute the extremal function l_ext
16 X=sym.Matrix(2, 1, [1, x])
17 M=sym.Matrix(2, 2, lambda i,j:
18     sym.integrate(X[i,0]*X[j,0]*v, (x, -t, t)))
19 V=sym.Matrix(2, 1, [DFO(1), DFO(x)])
20 Lext=M.LUsolve(V)
21 lext=((Lext.T)*X)[0,0]
22 # Compute expression test at the two vertices and print it
23 test=2*(1+1+n)+(t*s-2*n*c)/1-t*lext
24 print(sym.factor(test.subs(x,-t)))
25 print(sym.factor(test.subs(x,t)))

```

Program 1. Rank one simple principal toric fibrations

Program 1 prints the condition to check when c and p are variables, $n = 3$, $s = -6$ and $t = 1$, as used in Proposition 5.3. By modifying Line 5 and 6, one can obtain the conditions for an arbitrary simple principal \mathbb{P}^1 -bundle.

```

1 import sympy as sym
2 # variables on the plane
3 x1, x2 = sym.symbols('x1,x2')
4 # data of toric fibration and admissible Kahler class
5 c, p1, p2, n, s, t = 12, 1, 2, 3, 18, 1
6 ## weights associated to the data
7 l=c+p1*x1+p2*x2
8 v=l**n
9 w0=-s*l**(n-1) # for now, unknown l_ext replaced with zero
10 # list of vertices of the polytope
11 vert= [[2*t,-t], [-t,-t], [-t,2*t]]
12 # Donaldson-Futaki invariant with weights (v,w0)

```

```

13 def DF0(f):
14     interior=sym.integrate(sym.integrate(f*w0,
15     (x2,-t,t-x1)),(x1,-t,2*t))
16     facet1=sym.integrate((2*f*v).subs(x2,-t),(x1,-t,2*t))
17     facet2=sym.integrate((2*f*v).subs(x2,t-x1),
18     (x1,-t,2*t))
19     facet3=sym.integrate((2*f*v).subs(x1,-t),(x2,-t,2*t))
20     return(interior+facet1+facet2+facet3)
21 # Compute the extremal function l_ext
22 X=sym.Matrix(3, 1, [1, x1, x2])
23 M=sym.Matrix(3, 3, lambda i,j:
24     sym.integrate(sym.integrate(X[i,0]*X[j, 0]*v,
25     (x2,-t,t-x1)),(x1,-t,2*t))
26 V=sym.Matrix(3, 1, [DF0(1), DF0(x1), DF0(x2)])
27 Lext=M.LUsolve(V)
28 lext=((Lext.T)*X)[0,0]
29 # Compute and print the minimum of expression test on
30 # vertices
31 test=2*(1+2+n)+(t*s-2*n*c)/1-t*lext
32 test_vertices=test.subs(x1,vert[0][0]).subs(x2,vert[0][1])
33 for i in range(1,len(vert)):
34     test_vertices=sym.Min(test_vertices,
35     test.subs(x1,vert[i][0]).subs(x2,vert[i][1]))
36 print("The minimum of expression test on vertices is ",
37 test_vertices)

```

Program 2. Simple principal \mathbb{P}^2 toric fibrations

Program 2 computes the condition when all the data are given the fixed values $(c, p_1, p_2, n, s, t) = (12, 1, 2, 3, 18, 1)$. Changing the values on the right-hand side of Line 5 allows to check the sufficient condition for arbitrary fixed values. If one wants one or several of the above quantities to be treated as variables, for example c , p_1 and p_2 , it suffices to remove these and the corresponding values on the right in Line 5 and add the line

```

6 c, p1, p2 = sym.symbols('c,p1,p2')

```

Since the program will now compute values of `test` as symbolic expressions, it will no longer be able to determine the minimum. One should thus replace Lines 28–32 for example by

```

28 print(sym.separatevars(test.subs(x1,vert[2][0]).
29     subs(x2,vert[2][1])))

```

to get the expressions from Appendix B, to be used in the proof of Proposition 5.1.

Similarly, it is very easy to modify the program to consider another Fano toric surface as fiber (Recall that there are five smooth Fano toric surfaces: $\mathbb{P}^1 \times \mathbb{P}^1$ and the blowups of \mathbb{P}^2 at up to three fixed points under the torus action). It suffices to modify Lines 10–18 according to the desired polytope. For example, if one wants to work with fiber the first Hirzebruch surface (i.e. the blowup of \mathbb{P}^2 at one point), then it suffices to replace Lines 10–18 with

```

10 # list of vertices of the polytope
11 vert= [[-t,-t], [t,-t], [t,0], [-t,2t]]
12 # Donaldson-Futaki invariant with weights (v,w0)
13 def DF0(f):
14     interior=sym.integrate(sym.integrate(f, (x2, -t, t-x1)),
15         (x1, -t, t))
16     facet1=sym.integrate(f.subs(x2,-t), (x1, -t, t))
17     facet2=sym.integrate(f.subs(x2,t-x1), (x1, -t, t))
18     facet3=sym.integrate(f.subs(x1,-t), (x2, -t, 2t))
19     facet4=sym.integrate(f.subs(x1,t), (x2, -t, 0))
20     return(interior+facet1+facet2+facet3+facet4)

```

Appendix B. Complement of proof of Proposition 5.1

$P(c, p_1, p_2)$

$$\begin{aligned}
&:= 12250c^{10} + 24500c^9p_1 - 39690c^8p_1^2 + 18060c^7p_1^3 - 22470c^6p_1^4 - 31752c^5p_1^5 \\
&\quad - 53376c^4p_1^6 + 22740c^3p_1^7 - 57024c^2p_1^8 + 1312p_1^9 - 49000c^9p_2 \\
&\quad + 34650c^7p_1^2p_2 + 286860c^6p_1^3p_2 + 152460c^5p_1^4p_2 + 360972c^4p_1^5p_2 \\
&\quad - 59520c^3p_1^6p_2 + 230112c^2p_1^7p_2 + 18288cp_1^8p_2 - 464p_1^9p_2 - 127890c^8p_2^2 \\
&\quad - 212310c^7p_1p_2^2 - 615510c^6p_1^2p_2^2 - 373212c^5p_1^3p_2^2 - 921924c^4p_1^4p_2^2 \\
&\quad - 425376c^3p_1^5p_2^2 - 160632cp_1^6p_2^2 - 19296p_1^8p_2^2 + 141540c^7p_2^3 + 657300c^6p_1p_2^3 \\
&\quad + 603288c^5p_1^2p_2^3 + 1408632c^4p_1^3p_2^3 + 390936c^3p_1^4p_2^3 + 571536c^2p_1^5p_2^3 \\
&\quad + 349440cp_1^6p_2^3 + 41376p_1^7p_2^3 - 328650c^6p_2^4 - 531720c^5p_1p_2^4 - 1421136c^4p_1^2p_2^4 \\
&\quad - 806100c^3p_1^3p_2^4 - 829080c^2p_1^4p_2^4 - 497592cp_1^5p_2^4 - 22416p_1^6p_2^4 + 212688c^5p_2^5 \\
&\quad - 43812c^3p_1^5p_2^2 + 860184c^4p_1p_2^5 + 849456c^3p_1^2p_2^5 + 906192c^2p_1^3p_2^5 \\
&\quad + 485712cp_1^4p_2^5 - 7488p_1^5p_2^5 - 286728c^4p_2^6 - 527016c^3p_1p_2^6 - 725760c^2p_1^2p_2^6 \\
&\quad - 329952cp_1^3p_2^6 + 127890c^8p_1p_2 + 7352cp_1^9 + 22656p_1^4p_2^6 + 150576c^3p_2^7 \\
&\quad + 363168c^2p_1p_2^7 + 156096cp_1^2p_2^7 - 25728p_1^3p_2^7 - 90792c^2p_2^8 - 46368cp_1p_2^8 \\
&\quad + 16992p_1^2p_2^8 + 10304cp_2^9 - 7040p_1p_2^9 + 1408p_2^{10} + 132300c^7p_1^2 + 105840c^6p_1^3 \\
&\quad - 11340c^5p_1^4 + 125496c^4p_1^5 + 151200c^3p_1^6 - 79056c^2p_1^7 + 60048cp_1^8 \\
&\quad - 12096p_1^9 - 396900c^7p_1p_2 - 449820c^6p_1^2p_2 - 260820c^5p_1^3p_2 - 374220c^4p_1^4p_2 \\
&\quad - 420336c^3p_1^5p_2 + 358992c^2p_1^6p_2 - 364176cp_1^7p_2 + 17712p_1^8p_2 + 396900c^7p_2^3 \\
&\quad + 714420c^6p_1p_2^3 + 601020c^5p_1^2p_2^3 + 378756c^4p_1^3p_2^3 + 743904c^3p_1^4p_2^3 \\
&\quad - 557280c^2p_1^5p_2^3 + 734832cp_1^6p_2^3 + 84240p_1^7p_2^3 - 476280c^6p_2^3 - 680400c^5p_1p_2^3 \\
&\quad - 282744c^4p_1^2p_2^3 - 728784c^3p_1^3p_2^3 + 99792c^2p_1^4p_2^3 - 1073520cp_1^5p_2^3 \\
&\quad - 287280p_1^6p_2^3 + 340200c^5p_2^4 + 45360c^4p_1p_2^4 + 568512c^3p_1^2p_2^4 + 829440c^2p_1^3p_2^4
\end{aligned}$$

$$\begin{aligned}
& + 1551312cp_1^4p_2^4 - 244944c^3p_1p_2^5 - 241056c^2p_2^7 - 736128cp_1p_2^7 - 18144c^4p_2^5 \\
& + 415152p_1^5p_2^4 - 279072p_1^2p_2^7 + 184032cp_2^8 + 139968p_1p_2^8 - 31104p_2^9 \\
& - 1175472c^2p_1^2p_2^5 - 1732752cp_1^3p_2^5 - 358992p_1^4p_2^5 + 81648c^3p_2^6 \\
& + 843696c^2p_1p_2^6 + 1436400cp_1^2p_2^6 + 323568p_1^3p_2^6
\end{aligned}$$

$Q(c, p_1, p_2)$

$$\begin{aligned}
& := 6125c^9 + 2205c^7p_1^2 + 210c^6p_1^3 + 14175c^5p_1^4 - 7812c^4p_1^5 + 24c^3p_1^6 \\
& + 9072c^2p_1^7 - 5004cp_1^8 + 688p_1^9 - 2205c^7p_1p_2 - 315c^6p_1^2p_2 - 28350c^5p_1^3p_2 \\
& - 31752c^2p_1^6p_2 + 20016cp_1^7p_2 - 3096p_1^8p_2 + 2205c^7p_2^2 - 315c^6p_1p_2^2 + 42525c^5p_1^2p_2^2 \\
& - 7812c^4p_1^3p_2^2 + 4356c^3p_1^4p_2^2 + 40824c^2p_1^5p_2^2 - 40320cp_1^6p_2^2 + 4464p_1^7p_2^2 \\
& + 210c^6p_2^3 - 28350c^5p_1p_2^3 - 7812c^4p_1^2p_2^3 - 8592c^3p_1^3p_2^3 - 22680c^2p_1^4p_2^3 \\
& + 50904cp_1^5p_2^3 - 1176p_1^6p_2^3 + 19530c^4p_1^4p_2 - 72c^3p_1^5p_2 \\
& + 14175c^5p_2^4 + 19530c^4p_1p_2^4 + 4356c^3p_1^2p_2^4 - 22680c^2p_1^3p_2^4 - 56196cp_1^4p_2^4 \\
& - 1224p_1^5p_2^4 - 7812c^4p_2^5 - 72c^3p_1p_2^5 + 40824c^2p_1^2p_2^5 + 50904cp_1^3p_2^5 - 1224p_1^4p_2^5 \\
& + 24c^3p_2^6 - 31752c^2p_1p_2^6 - 40320cp_1^2p_2^6 - 1176p_1^3p_2^6 + 9072c^2p_2^7 + 20016cp_1p_2^7 \\
& + 4464p_1^2p_2^7 - 5004cp_2^8 - 3096p_1p_2^8 + 688p_2^9
\end{aligned}$$

BIBLIOGRAPHY

- [Abr01] Miguel Abreu, *Kähler metrics on toric orbifolds*, J. Differ. Geom. **58** (2001), no. 1, 151–187. ↑125
- [Abr03] ———, *Kähler geometry of toric manifolds in symplectic coordinates*, Symplectic and contact topology: Interactions and perspectives, AMS Fields Institute Communications, vol. 35, American Mathematical Society, 2003, pp. 1–24. ↑132
- [ACC⁺23] Caroline Araujo, Ana-Maria Castravet, Ivan Cheltsov, Kento Fujita, Anne-Sophie Kaloghiros, Jesus Martinez-Garcia, Constantin Shramov, Hendrik Süß, and Nivedita Viswanathan, *The Calabi problem for Fano threefolds*, 2023, available at <https://www.maths.ed.ac.uk/cheltsov/research.html>. ↑138
- [ACG06] Vestislav Apostolov, David M. J. Calderbank, and Paul Gauduchon, *Hamiltonian 2-Forms in Kähler Geometry, I General Theory*, J. Differ. Geom. **73** (2006), no. 3, 359–412. ↑128, 131, 132
- [ACGTF04] Vestislav Apostolov, David M. J. Calderbank, Paul Gauduchon, and Christina W. Tønnesen-Friedman, *Hamiltonian 2-forms in Kähler geometry. II: Global classification*, J. Differ. Geom. **68** (2004), no. 2, 277–345. ↑127, 128, 132
- [ACGTF08] ———, *Hamiltonian 2-forms in Kähler geometry. III: Extremal metrics and stability*, Invent. Math. **173** (2008), no. 3, 547–601. ↑118, 128, 138, 140, 141
- [ACGTF11] ———, *Extremal Kähler metrics on projective bundles over a curve*, Adv. Math. **227** (2011), no. 6, 2385–2424. ↑127, 130, 131, 133, 139
- [AJL22] Vestislav Apostolov, Simon Jubert, and Abdellah Lahdili, *Weighted K-stability and coercivity with applications to extremal Kahler and Sasaki metrics*, <https://arxiv.org/abs/2104.09709v2>, 2022. ↑126, 127, 128, 131, 133, 134, 136, 137

- [Apo22] Vestislav Apostolov, *The Kähler geometry of toric manifolds*, Lecture Notes of CIRM winter school 2019, <https://arxiv.org/abs/2208.12493>, 2022. ↑119, 126
- [BB13] Robert J. Berman and Bo Berndtsson, *Real Monge–Ampère equations and Kähler–Ricci solitons on toric log Fano varieties*, Ann. Fac. Sci. Toulouse, Math. **22** (2013), no. 4, 649–711. ↑126
- [Cal82] Eugenio Calabi, *Extremal Kähler metrics*, Seminar on Differential Geometry, Annals of Mathematics Studies, vol. 102, Princeton University Press, 1982, pp. 259–290. ↑118
- [Cal85] ———, *Extremal Kähler metrics, II*, Differential Geometry and Complex Analysis (I. Chavel and H. M. Farkas, eds.), Differential geometry and complex analysis, vol. dedic. H. E. Rauch, Springer, 1985, pp. 95–114. ↑130
- [CC18] Xiuxiong Chen and Jingrui Chen, *On the constant scalar curvature Kähler metrics (III) – general automorphism group*, <https://arxiv.org/abs/1801.05907>, 2018. ↑118, 126, 131, 132
- [CC21a] Xiuxiong Chen and Jingrui Cheng, *On the constant scalar curvature Kähler metrics. I: A priori estimates*, J. Am. Math. Soc. **34** (2021), no. 4, 909–936. ↑131, 132
- [CC21b] ———, *On the constant scalar curvature Kähler metrics. II: Existence results*, J. Am. Math. Soc. **34** (2021), no. 4, 937–1009. ↑131, 132
- [Del88] Thomas Delzant, *Hamiltoniens périodiques et image convexe de l’application moment*, Bull. Soc. Math. Fr. **116** (1988), no. 3, 315–339. ↑125, 127, 129, 132, 134
- [Del22] Thibaut Delcroix, *Uniform K-stability of polarized spherical varieties*, <https://arxiv.org/abs/2009.06463>, 2022. ↑119, 137
- [DGSW18] Ved Datar, Bin Guo, Jian Song, and Xiaowei Wang, *Connecting toric manifolds by conical Kähler–Einstein metrics*, Adv. Math. **323** (2018), 38–83. ↑125
- [Don02] Simon K. Donaldson, *Scalar curvature and stability of toric varieties*, J. Differ. Geom. **62** (2002), no. 2, 289–349. ↑121, 122, 124, 126, 132
- [Don08] ———, *Kähler geometry on toric manifolds, and some other manifolds with large symmetry*, Handbook of geometric analysis. No. 1, Advanced Lectures in Mathematics (ALM), vol. 7, International Press, 2008, pp. 29–75. ↑119
- [FM95] Akito Futaki and Toshiki Mabuchi, *Bilinear forms and extremal Kähler vector fields associated with Kähler classes*, Math. Ann. **301** (1995), no. 2, 199–210. ↑130
- [Gau10] Paul Gauduchon, *Calabi’s extremal metrics: An elementary introduction*, 2010, Lecture Notes, Preprint, p. 7. ↑119
- [GS82] Victor W. Guillemin and Shlomo Sternberg, *Convexity properties of the moment mapping*, Invent. Math. **67** (1982), 491–513. ↑130
- [Gua99] Daniel Guan, *On modified Mabuchi functional and Mabuchi moduli space of Kähler metrics on toric bundle*, Math. Res. Lett. **6** (1999), no. 5-6, 547–555. ↑132
- [Gui94] Victor W. Guillemin, *Kähler structures on toric varieties*, J. Differ. Geom. **40** (1994), no. 2, 285–309. ↑132
- [He19] Weiyong He, *On Calabi’s extremal metrics and properness*, Trans. Am. Math. Soc. **372** (2019), no. 8, 5595–5619. ↑126, 131, 132
- [His20] Tomoyuki Hisamoto, *Stability and coercivity for toric polarizations*, <https://arxiv.org/abs/1610.07998v3>, 2020. ↑121, 126
- [HS02] Andrew D. Hwang and Michael A. Singer, *A momentum construction for circle-invariant Kähler metrics*, Trans. Am. Math. Soc. **354** (2002), no. 6, 2285–2325. ↑118
- [Jub21] Simon Jubert, *A Yau–Tian–Donaldson correspondance on a class of toric fibration*, <https://arxiv.org/abs/2108.12297v3>, 2021. ↑118, 125, 126, 131, 132, 133, 134, 137

- [Lah19] Abdellah Lahdili, *Kähler metrics with weighted constant scalar curvature and weighted K -stability*, Proc. Lond. Math. Soc. **119** (2019), no. 4, 1065–1114. ↑121, 125, 130, 131, 132
- [Leg16] Eveline Legendre, *Toric Kähler–Einstein metrics and convex compact polytopes*, J. Geom. Anal. **26** (2016), no. 1, 399–427. ↑125
- [Leg19] ———, *A note on extremal toric almost Kähler metrics*, Moduli of K -stable varieties, Springer INdAM Series, vol. 31, Springer, 2019, pp. 53–74. ↑126
- [LH20] Chi Li and Jiyuan Han, *On the Yau–Tian–Donaldson conjecture for generalized Kähler–Ricci soliton equations*, accepted by *Communications on Pure and Applied Mathematics*, <https://arxiv.org/abs/2006.00903>, 2020. ↑126, 137
- [Li20] Chi Li, *Geodesic rays and stability in the cscK problem*, <https://arxiv.org/abs/2001.01366>, 2020. ↑126
- [LLS16] An-Min Li, Zhao Lian, and Li Sheng, *Some Estimates for a Generalized Abreu’s Equation*, Differ. Geom. Appl. **48** (2016), 87–103. ↑121, 125, 126
- [LLS17] ———, *Interior regularity for a Generalized Abreu Equation*, Int. J. Math. **28** (2017), no. 7, article no. 1750053. ↑126
- [LLS21] ———, *Extremal metrics on toric manifolds and homogeneous toric bundles*, <https://arxiv.org/abs/2110.08491>, 2021. ↑126, 140
- [LS93] Claude LeBrun and Santiago R. Simanca, *On the Kähler classes of extremal metrics*, Geometry and global analysis (Sendai, 1993), Tohoku University, Sendai, 1993, pp. 255–271. ↑137
- [Mab86] Toshiki Mabuchi, *K -energy Maps Integrating Futaki Invariants*, Tôhoku Math. J. **38** (1986), no. 1-2, 575–593. ↑132
- [Mat57] Yozô Matsushima, *Sur la structure du groupe d’homéomorphismes analytiques d’une certaine variété kählérienne*, Nagoya Math. J. **11** (1957), 145–150. ↑118
- [NS21] Yasufumi Nitta and Shunsuke Saito, *A uniform version of the Yau–Tian–Donaldson correspondence for extremal Kähler metrics on polarized toric manifolds*, <https://arxiv.org/abs/2110.10386v1>, 2021. ↑121, 122, 124, 126
- [Szé07] Gábor Székelyhidi, *Extremal metrics and K -stability*, Bull. Lond. Math. Soc. **39** (2007), no. 1, 76–84. ↑141
- [Szé14] ———, *An introduction to extremal Kähler metrics*, Graduate Studies in Mathematics, vol. 152, American Mathematical Society, 2014. ↑119
- [WZ04] Xu-Jia Wang and Xiaohua Zhu, *Kähler–Ricci solitons on toric manifolds with positive first Chern class*, Adv. Math. **188** (2004), no. 1, 87–103. ↑126
- [ZZ08] Bin Zhou and Xiaohua Zhu, *Relative K -stability and modified K -energy on toric manifolds*, Adv. Math. **219** (2008), no. 4, 1327–1362. ↑119, 122, 126, 132, 133

Manuscript received on 11th February 2022,
revised on 11th July 2022,
accepted on 10th October 2022.

Recommended by Editors X. Caruso and X. Chu.
Published under license CC BY 4.0.



eISSN: 2644–9463

This journal is a member of Centre Mersenne.



Thibaut DELCROIX
IMAG, Univ Montpellier,
CNRS, Montpellier (France)
thibaut.delcroix@umontpellier.fr

Simon JUBERT
Département de Mathématiques,
UQAM, C.P. 8888,
Succursale Centre-ville,
Montréal (Quebec),
H3C 3P8 (Canada)
Institut de Mathématiques
de Toulouse
Université Paul Sabatier
118 route de Narbonne
31062 Toulouse (France)
simonjubert@gmail.com