



ANNALES
HENRI LEBESGUE

JÉRÉMI DARDÉ

ARMAND KOENIG

JULIEN ROYER

NULL-CONTROLLABILITY
PROPERTIES OF THE
GENERALIZED TWO-DIMENSIONAL
BAOUENDI-GRUSHIN EQUATION
WITH NON-RECTANGULAR
CONTROL SETS

PROPRIÉTÉS DE CONTRÔLABILITÉ À
ZÉRO DE L'ÉQUATION DE
BAOUENDI-GRUSHIN GÉNÉRALISÉE
BIDIMENSIONNELLE AVEC DOMAINES DE
CONTRÔLE NON RECTANGULAIRES

Keywords: null-controllability, observability, degenerate parabolic equations, resolvent estimates.
2020 *Mathematics Subject Classification:* 35K65, 93B05, 47B28, 47A10.

DOI: <https://doi.org/10.5802/ahl.193>

(*) This work has been partially supported by the ANR LabEx CIMI (under grant ANR-11-LABX-0040) within the French State Programme “Investissements d’Avenir”.

ABSTRACT. — We consider the null-controllability problem for the *generalized Baouendi–Grushin equation* $(\partial_t - \partial_x^2 - q(x)^2 \partial_y^2)f = \mathbb{1}_\omega u$ on a rectangular domain. Sharp controllability results already exist when the control domain ω is a vertical strip, or when $q(x) = x$. In this article, we provide upper and lower bounds for the minimal time of null-controllability for general q and non-rectangular control region ω . In some geometries for ω , the upper bound and the lower bound are equal, in which case, we know the exact value of the minimal time of null-controllability.

Our proof relies on several tools: known results when ω is a vertical strip and cutoff arguments for the upper bound of the minimal time of null-controllability; spectral analysis of the Schrödinger operator $-\partial_x^2 + \nu^2 q(x)^2$ when $\operatorname{Re}(\nu) > 0$, pseudo-differential-type operators on polynomials and Runge’s theorem for the lower bound.

RÉSUMÉ. — Nous considérons le problème de la contrôlabilité à zéro de l’équation de Baouendi–Grushin généralisée $(\partial_t - \partial_x^2 - q(x)^2 \partial_y^2)f = \mathbb{1}_\omega u$ sur un domaine rectangulaire. On connaît déjà des résultats précis de contrôlabilité lorsque le domaine de contrôle ω est une bande verticale, ou lorsque $q(x) = x$. Dans cet article, nous démontrons des bornes supérieures et inférieures du temps minimal de contrôlabilité à zéro, pour une classe générale de potentiels q et de domaines de contrôle ω possiblement non rectangulaires. Pour certaines zones de contrôle, la borne supérieure et la borne inférieure coïncident, auquel cas nous connaissons la valeur exacte du temps minimal de contrôlabilité à zéro.

Notre démonstration s’appuie sur plusieurs outils. Pour la borne supérieure du temps minimal de contrôlabilité, nous utilisons des résultats connus lorsque ω est une bande verticale et des arguments de troncature. Pour la borne inférieure, nous utilisons une analyse spectrale de l’opérateur de Schrödinger $-\partial_x^2 + \nu^2 q(x)^2$ lorsque $\operatorname{Re}(\nu) > 0$, des opérateurs de type pseudo-différentiel sur les polynômes et le théorème de Runge.

1. Introduction and statements of the main results

1.1. The Baouendi–Grushin equation

In this article, we study some controllability properties of the two-dimensional *generalized Baouendi–Grushin equation*.

Let L_-, L_+ be positive, and $I = (-L_-, L_+)$. Let $q \in C^0(\bar{I})$ such that $q(0) = 0$ and $q(x) \neq 0$ for all $x \in \bar{I} \setminus \{0\}$. We denote by \mathbb{T} the one-dimensional torus $\mathbb{R}/2\pi\mathbb{Z}$. For $T > 0$, $f_0 \in L^2(I \times \mathbb{T})$ and $F \in L^2((0, T); L^2(I \times \mathbb{T}))$ we consider the generalized Baouendi–Grushin equation

$$(1.1) \quad \begin{cases} (\partial_t - \partial_x^2 - q(x)^2 \partial_y^2) f(t, x, y) = F(t, x, y), & t \in (0, T), x \in I, y \in \mathbb{T} \\ f(t, x, y) = 0, & t \in (0, T), x \in \partial I, y \in \mathbb{T}, \\ f(0, x, y) = f_0, & x \in I, y \in \mathbb{T}. \end{cases}$$

Note that, because $q(0) = 0$, the equation degenerates on the vertical axis $\{0\} \times \mathbb{T}$. Nevertheless, the equation is well posed. Precisely, the Friedrichs extension (see [Hel13, Section 4.3]) of the operator

$$f \in C_c^\infty(I \times \mathbb{T}) \mapsto -\partial_x^2 f - q(x)^2 \partial_y^2 f,$$

generates an analytic semigroup, which allows to define a solution of the generalized Baouendi–Grushin equation (1.1) in the sense of semigroups [Bre11]. In our case, this solution is smooth in the following sense:

PROPOSITION 1.1. — *For any source term $F \in L^2((0, T); L^2(I \times \mathbb{T}))$ and any initial condition $f_0 \in L^2(I \times \mathbb{T})$, there exists a unique $f \in C^0([0, T]; L^2(I \times \mathbb{T})) \cap L^2((0, T); V)$ solution of the generalized Baouendi–Grushin equation (1.1), with*

$$V = \left\{ f \in L^2(I \times \mathbb{T}), \partial_x f \in L^2(I \times \mathbb{T}), q \partial_y f \in L^2(I \times \mathbb{T}) \right\}.$$

This result is proved in [BCG14] in the case $q(x) = |x|^\gamma$ with $\gamma > 0$. The proof is easily generalized to our case of interest.

1.2. Control problem for the Baouendi–Grushin equation

Our study focuses on internal null-controllability of the Baouendi–Grushin equation. More precisely, let $\omega \subset I \times \mathbb{T}$ be a non-empty open set and $u \in L^2((0, T); L^2(\omega))$. The controlled Baouendi–Grushin equation reads

$$(1.2) \quad \begin{cases} (\partial_t - \partial_x^2 - q(x)^2 \partial_y^2) f(t, x, y) = \mathbb{1}_\omega u(t, x, y), & t \in (0, T), x \in I, y \in \mathbb{T}, \\ f(t, x, y) = 0, & t \in (0, T), x \in \partial I, y \in \mathbb{T}, \\ f(0, x, y) = f_0, & x \in I, y \in \mathbb{T}, \end{cases}$$

where f is the state of the system, and $\mathbb{1}_\omega u$ is the control supported in ω .

DEFINITION 1.2 (Null-controllability). — *Let $T > 0$. The Baouendi–Grushin equation (1.2) is said to be null-controllable on ω in time T if, for any initial condition $f_0 \in L^2(I \times \mathbb{T})$, there exists $u \in L^2((0, T); L^2(\omega))$ such that the solution f of eq. (1.2) satisfies $f(T, \cdot, \cdot) = 0$ in $I \times \mathbb{T}$.*

It is known that, contrary to usual non-degenerate parabolic equations like the heat equation, due to the degeneracy of q on $\{0\} \times \mathbb{T}$, the null-controllability properties of (1.2) strongly depend on the control set ω and the time horizon T . More precisely, for certain control sets ω , there is no time $T > 0$ such that eq. (1.2) is null-controllable, whereas for other control sets ω a minimal time of null-controllability appears. We refer to the bibliographical comments, Section 1.4 below, for a detailed description of the known results on the subject.

In the present paper, we aim to give precise null-controllability results for equation (1.2), for a large class of control sets ω and a large class of functions q .

1.3. Main results

We are interested in the case where the equation is degenerate on $\{x = 0\}$. Thus, we assume that $q(x) = 0$ only when $x = 0$, and we assume without loss of generality that $q'(0) > 0$.

Before presenting the main results of our study, we introduce the so-called *Agmon distance* of a point $x \in I$ to the origin, defined by:⁽¹⁾

$$(1.3) \quad d_{\text{Agm}} : x \in I \mapsto \int_0^x q(s) \, ds.$$

This quantity appears naturally in the computation of the minimal time of null-controllability for the generalized Baouendi–Grushin equation.

⁽¹⁾It corresponds to the usual definition of the Agmon distance given for example in [DS99, Eq. (6.3)], with $V = q^2$ and $E = 0$.

1.3.1. Lack of null-controllability in small time for a class of control sets ω

Our main result is a negative result of null-controllability for small times. We show that if the control set ω stays at positive distance from a horizontal segment of the form $(a, b) \times \{y_0\}$, with $-L_- \leq a < 0 < b \leq L_+$ and $y_0 \in \mathbb{T}$, then equation (1.2) is not null-controllable on ω for time T smaller than a precisely given critical time.

To properly state the result, we need to introduce a modified version of the Agmon distance. We set $\delta(x) = d_{\text{Agm}}(x)$ for $x \in I$, and $\delta(-L_-) = \delta(L_+) = +\infty$.

THEOREM 1.3. — *Assume that $q \in C^2(\bar{I})$ is such that $q(0) = 0$, $q'(0) > 0$ and $q(x) \neq 0$ whenever $x \neq 0$. Let ω be an open subset of $I \times \mathbb{T}$. Assume that there exist $a \in [-L_-, 0)$, $b \in (0, L_+]$ and $y_0 \in \mathbb{T}$ such that*

$$\text{distance}\left((a, b) \times \{y_0\}, \omega\right) > 0.$$

(See Figure 1.1.) Then, the generalized Baouendi–Grushin equation (1.2) is not null-controllable on ω in time T such that

$$T < \frac{1}{q'(0)} \min(\delta(a), \delta(b)).$$




Figure 1.1. In green, an example of a domain ω with, in thick black, an example of a horizontal segment that stays at positive distance of ω . In this example, Theorem 1.3 implies that the generalised Baouendi–Grushin equation is not null-controllable in time $T < d_{\text{Agm}}(a)$.

This theorem is a generalization of [DK20, Theorem 3.3], where the result is proved for I symmetric with respect to the origin and, more restrictively, for $q(x) = x$. A key step in our proof of Theorem 1.3 is the study of spectral properties of the family of operators defined on $L^2(I)$ by

$$(1.4) \quad P_\nu: -\partial_x^2 + \nu^2 q(x)^2, \quad \text{Dom}(P_\nu) = H^2(I) \cap H_0^1(I).$$

Because of technical reasons, we have to consider P_ν for every $\text{Re}(\nu) > 0$, which makes P_ν non self-adjoint.

In the previous article [DK20], the corresponding results were obtained using explicit solutions of particular ordinary differential equations. These explicit formulae are not available in our general setting.

In our case, we obtain a localization of an eigenvalue of P_ν , as well as precise Agmon type estimates for an associated eigenfunction,⁽²⁾ uniformly in $\nu = |\nu|e^{i\theta}$ with $|\nu|$ large enough and $0 \leq |\theta| \leq \theta_0$ for some $\theta_0 \in [0, \pi/2)$. To that end, we compare P_ν with the non-selfadjoint harmonic oscillator $H_{q'(0)\nu} := -\partial_x^2 + (q'(0)\nu)^2 x^2$. Theorem 1.3 is proved in Section 3, with the spectral analysis done in Section 4.

Note that [HSS05] contains closely related spectral asymptotics. However, we cannot apply them directly since our domain has a boundary and we need uniform estimates with respect to the parameter ν .

1.3.2. Precise critical time of null-controllability for a class of control sets ω

With Theorem 1.3 we can deduce in some settings the critical time for the null-controllability of (1.2).

We first mention a natural adaptation of [DK20, Theorem 3.1]. It was actually claimed in [DK20, Remark 3.2], but the statement was imprecise if q is not odd. We take the opportunity to correct the statement:

THEOREM 1.4. — *Assume that $q \in C^3(\bar{I})$ is such that $q(0) = 0$ and $\min_{\bar{I}} q' > 0$. Let ω be an open subset of $I \times \mathbb{T}$. Assume that there exists a closed path $\gamma = (\gamma_x, \gamma_y) \in C^0(\mathbb{T}; \omega)$ such that $\{-L_-\} \times \mathbb{T}$ and $\{L_+\} \times \mathbb{T}$ are included in different connected components of $(\bar{I} \times \mathbb{T}) \setminus \gamma(\mathbb{T})$ (see Figure 1.2).*

The generalized Baouendi–Grushin equation (1.2) is null-controllable on ω in time T such that

$$T > \frac{1}{q'(0)} \max_{\mathbb{T}} d_{\text{Agm}}(\gamma_x).$$




Figure 1.2. In green, an example of a domain ω , with, in blue, a corresponding path γ that satisfies the hypotheses of Theorem 1.4.

Remark 1.5. — In this theorem, we can replace the hypothesis “ $\{-L_-\} \times \mathbb{T}$ and $\{L_+\} \times \mathbb{T}$ are included in different connected components of $(\bar{I} \times \mathbb{T}) \setminus \gamma(\mathbb{T})$ ” by “ γ is not homotopic to a constant path”. These two conditions are essentially equivalent. We discuss this in Propositions B.2, B.3 and Remark B.4.

⁽²⁾ For $\nu \in \mathbb{R}_+^*$, we localize the smallest eigenvalue of P_ν , and for ν complex, we localize its analytic continuation.

Theorem 1.4 is proved in Section 2. Combining Theorems 1.3 and 1.4, we obtain the following result which gives the precise critical time of null-controllability of the generalized Baouendi–Grushin for a large class of control sets ω and functions q :

THEOREM 1.6. — Assume that $q \in C^3(\bar{I})$ is such that $q(0) = 0$ and $\min_{\bar{I}} q' > 0$. Let $\gamma_1, \gamma_2 \in C^0(\mathbb{T}; I)$ such that for every $y \in \mathbb{T}$, $\gamma_1(y) < \gamma_2(y)$. Let $\omega = \{(x, y) \in I \times \mathbb{T} : \gamma_1(y) < x < \gamma_2(y)\}$ (see Figure 1.3) and⁽³⁾

$$T_* := \frac{1}{q'(0)} \max \left(d_{\text{Agm}} \left(-\max_{\mathbb{T}} (\gamma_2^-) \right), d_{\text{Agm}} \left(\max_{\mathbb{T}} (\gamma_1^+) \right) \right).$$

Then the generalized Baouendi–Grushin equation (1.2) is null-controllable on ω in any time $T > T_*$, but it is not null-controllable on ω in time $T < T_*$.

This theorem is proved in Section 5.




Figure 1.3. In green, an example of a domain ω that satisfies the hypotheses of Theorem 1.6.

1.3.3. Comments

Before proceeding further, we make some additional comments on our results.

- The assumptions regarding the function q in Theorem 1.3 are slightly more general than in Theorem 1.4. They seem also more natural in the context of our study. Therefore, we conjecture that Theorem 1.4 holds for functions $q \in C^3(\bar{I})$ satisfying

$$q(0) = 0, \quad q'(0) > 0, \quad q(x) \neq 0 \text{ for all } x \in \bar{I}.$$

But up to our knowledge, this is still an open question.

- There are still numerous geometrical configurations not included in Theorem 1.6. Nevertheless, in many situations, Theorems 1.4 and 1.3 give information about null-controllability properties. As an example, in the geometrical configuration described in Figure 1.4, combining Theorems 1.4 and 1.3, we obtain the existence of a critical time

$$T_* \in \left(\frac{d_{\text{Agm}}(-a)}{q'(0)}, \frac{d_{\text{Agm}}(-b)}{q'(0)} \right)$$

⁽³⁾When f is a real valued function, we denote $f^+ = \max(f, 0)$ and $f^- = \max(-f, 0)$ its positive and negative part respectively.

such that the Baouendi–Grushin equation is null-controllable on ω in time $T > T_*$, and is not null-controllable on ω in time $T < T_*$.




Figure 1.4. In this configuration, we obtain lower and upper bounds of the critical time of null-controllability

In the two geometrical configurations presented in Figure 1.5, Theorem 1.3 implies that the Baouendi–Grushin equation is not null-controllable on ω in time $T < d_{\text{Agm}}(a)/q'(0)$. Note that in these two configurations, the question of the null-controllability of (1.2) in ω for some time T large enough is still open.




Figure 1.5. In these configurations, we obtain a lower bound on the critical time of null-controllability.

- These results are stated for the generalized Baouendi–Grushin equation posed on $I \times \mathbb{T}$. They can be adapted to the equation posed on $I \times (0, \pi)$ with Dirichlet boundary conditions, with very similar proofs. We refer to Appendix A for details on the statements and the corresponding proofs.

1.4. Bibliographical comments

1.4.1. On the Baouendi–Grushin equation

The study of controllability properties of system (1.2) began with the pioneering work [BCG14], where the authors study the null-controllability of the equation

$$(1.5) \quad \begin{cases} (\partial_t - \partial_x^2 - |x|^{2\gamma} \partial_y^2) f = 1_\omega u, & t \in (0, T), \ x \in (-1, 1), \ y \in \mathbb{T}, \\ f(t, x, y) = 0, & t \in (0, T), \ x = \pm 1, \ y \in \mathbb{T}. \end{cases}$$

They prove that in the case $\gamma \in (0, 1)$ (weak degeneracy), the Baouendi–Grushin equation (1.2) is null-controllable for any control set ω and any time $T > 0$, whereas in the case $\gamma > 1$ (strong degeneracy), it is not null-controllable for any control set ω

and any time $T > 0$, except if ω contains $\{0\} \times \mathbb{T}$ in which case it is null-controllable in any positive time T .

More surprisingly, in the case $\gamma = 1$, which corresponds to the Baouendi–Grushin equation (1.2) with $q(x) = x$, and for $\omega = (a, b) \times \mathbb{T}$, with $0 < a < b$, there exists a critical time $T_* \geq \frac{a^2}{2}$ such that the Baouendi–Grushin equation (1.2) is null-controllable on ω in time T , for every $T > T_*$, and is not null controllable on ω in time T , for every $T < T_*$. It is also proved that if $\gamma = 1$ and ω contains the vertical line $\{0\} \times \mathbb{T}$, equation (1.5) is null controllable in any time $T > 0$. Such a minimal time of null-controllability would not be surprising for equations with finite speed of propagation, such as the wave equation [BLR92], but the Baouendi–Grushin equation has a *infinite* speed of propagation.

Many works followed, trying to characterize precisely the critical time T_* , and to generalize the result to different geometrical settings and different functions q . The first exact characterization of T_* is given in [BMM15] in the case $q(x) = x$ and with two symmetric vertical strips as control set, that is $\omega = (-1, -a) \times (a, 1)$, $a \in (0, 1)$. Using the transmutation method and sideways energy estimates, the authors prove that eq. (1.2) is null-controllable in ω in any time $T > \frac{a^2}{2}$, and is not null-controllable in ω in any time $T < \frac{a^2}{2}$.

When ω is a vertical strip of the form $(a, b) \times \mathbb{T}$, with $a > 0$, as in [BCG14], the precise value of the critical time T_* was obtained independently in the works [ABM21, BDE20, LL23]. More precisely, in [ABM21], using new estimates for biorthogonal sequences to real exponentials and the moments method, the authors prove that in the case $q(x) = x$, the critical time is $\frac{a^2}{2}$. In [BDE20], with a function q satisfying the assumptions of Theorem 1.6, the authors use a Carleman strategy to obtain that eq. (1.2) is null controllable on ω in any time $T > T_*$, and not null-controllable on ω in any time $T < T_*$, with

$$T_* = \frac{d_{\text{Agm}}(a)}{q'(0)}.$$

Very recently, this result was obtained in [LL23] using the moments method, with a stronger smoothness assumption on q (see [LL23, Remark 1.12 and Proposition 1.13]).

All the strategies developed in [ABM21, BCG14, BDE20, BMM15], although very different, rely on a Fourier expansion of system (1.2) with respect to the y -variable and the study of the obtained family of one dimensional parabolic equations in the variables t, x . As a consequence, the control set ω has to contain a vertical strip, which seems to be an important restriction of the proposed methods.

Nevertheless, in [DK20], the authors generalize the positive null-controllability results obtained in [BDE20] to a large class of control sets: in the setting of Theorem 1.6 and with the additional assumptions that I is symmetric and q is odd, system (1.2) is null controllable in any time $T > T_*$, with

$$T_* = \frac{1}{q'(0)} \max \left(d_{\text{Agm}}(-\max(\gamma_2^-)), d_{\text{Agm}}(\max(\gamma_1^+)) \right).$$

In the specific case $q(x) = x$, they also prove that if there exist $a, b \in I$, $a < 0 < b$, and $y_0 \in \mathbb{T}$ such that $\text{distance}((a, b) \times \{y_0\} \cap \omega) > 0$, then (1.2) is not null-controllable in time $T < \min(a^2, b^2)/2$, whereas if there is $y_0 \in \mathbb{T}$ such that $\text{distance}(I \times \{y_0\} \cap \omega) > 0$,

then (1.2) is not null controllable on ω in any positive time T . Theorem 1.6 is the generalization of this result to a wider class of functions q .

To end this overview on controllability issues for the parabolic Baouendi–Grushin equation, we point out that partial controllability results are known in some multidimensional configurations [BDE20] while precise results are known for cascade systems of two-dimensional Baouendi–Grushin equations with one control, in the case $q(x) = x$ [ABM21].

1.4.2. Some related problems

Let us briefly mention the literature on related problems, in several directions: other degenerate parabolic equations, minimal time of null controllability for parabolic systems, and other type of degenerate equations.

Since the pioneering works [FR71, Yeg63] on the null-controllability of the one-dimensional heat equation, the null-controllability of non-degenerate parabolic equations has been extensively studied. The null-controllability of degenerate parabolic equations is a more recent subject of study. The case of a degeneracy at the boundary of the domain is now well-understood [CMV16] (see also the references therein).

When the degeneracy occurs in the domain, we lack for the moment a general theory, and equations are studied case by case. The two-dimensional Baouendi–Grushin equations is arguably the simplest and best understood equation of that type. Very similar results, including a minimal time of null-controllability for quadratic degeneracy, have been observed for the heat equation on the Heisenberg group [BC17, BDE20], and the Kolmogorov equation [Bea14, BHHR15, BZ09, DR21, Koe20].

The related problem of *approximate controllability* for degenerate parabolic equations has been studied in a somewhat general framework [LL22].

A minimal time of null-controllability might also appear for the heat equation with punctual control [Dol73] and for systems of parabolic equations, degenerate or not [AKBGBT16, BBM20].

Finally, let us mention that the subelliptic wave equation is not controllable [Let23], and that the Grushin–Schrödinger equation has a minimal time of controllability [BS22, FKL21, LS23].

Acknowledgements

We express our gratitude to our colleagues Joan Bellier-Millès, Paulo Carrillo-Rouse and Joost Nuiten, for giving us the key arguments in algebraic topology to prove Proposition B.3.

2. Null-controllability in large time

In this section, we prove Theorem 1.4. The idea of the proof is to use known controllability results for equation (1.2) when the control set is a vertical strip combined with a cutoff argument. More precisely, we recall the following result [BDE20, Theorem 1.4].⁽⁴⁾

PROPOSITION 2.1. — *Assume that q satisfies the assumptions of Theorem 1.4. Let $\omega = (a, b) \times \mathbb{T}$, with $-L_- \leq a < b \leq L_+$. Then*

- if $0 < a$, the Baouendi–Grushin equation is null-controllable on ω in time $T > d_{\text{Agm}}(a)/q'(0)$,
- if $b < 0$, the Baouendi–Grushin equation is null-controllable on ω in time $T > d_{\text{Agm}}(b)/q'(0)$,
- if $a < 0 < b$, the Baouendi–Grushin equation is null-controllable on ω in time $T > 0$.

Proof of Theorem 1.4. — We set $\omega_- = (-L_-, \min_{\mathbb{T}}(\gamma_x)) \times \mathbb{T}$ and $\omega_+ = (\max_{\mathbb{T}}(\gamma_x), L_+) \times \mathbb{T}$ (see Figure 2.1).




Figure 2.1. Definition of ω_- (red) and ω_+ (blue).

Proposition 2.1 implies that the Baouendi–Grushin equation is null-controllable either on ω_- or on ω_+ in any time T such that

$$T > \frac{1}{q'(0)} \max \left(d_{\text{Agm}} \left(\min_{\mathbb{T}}(\gamma_x) \right), d_{\text{Agm}} \left(\max_{\mathbb{T}}(\gamma_x) \right) \right).$$

That is, for any initial condition $f_0 \in L^2(I \times \mathbb{T})$, there exist $u_- \in L^2((0, T); L^2(\omega_-))$ and $u_+ \in L^2((0, T); L^2(\omega_+))$ such that f_- and f_+ solutions of

$$\begin{cases} (\partial_t - \partial_x^2 - q(x)^2 \partial_y^2) f_{\pm}(t, x, y) = \mathbb{1}_{\omega_{\pm}} u_{\pm}(t, x, y), & t \in (0, T), x \in I, y \in \mathbb{T}, \\ f_{\pm}(t, x, y) = 0, & t \in (0, T), x \in \partial I, y \in \mathbb{T}, \\ f_{\pm}(0, x, y) = f_0(x, y), & x \in I, y \in \mathbb{T}, \end{cases}$$

satisfy $f_{\pm}(T, \cdot, \cdot) = 0$ in $I \times \mathbb{T}$.

By definition of γ , ω_+ and ω_- are included in two distinct connected components of $(I \times \mathbb{T}) \setminus \gamma(\mathbb{T})$. As a consequence, we can construct $\chi \in C^\infty(I \times \mathbb{T})$ such that

⁽⁴⁾The reference [BDE20] states the result with a control on the boundary. But cutoff arguments allow to construct controls on vertical strips from boundary controls, as in [DK20, Appendix A].

$\chi \equiv 1$ in $\omega_- \setminus \omega$, $\chi \equiv 0$ in $\omega_+ \setminus \omega$ and $\text{supp}(\nabla \chi) \subset \omega$ (see Proposition B.1). Define $f = (1 - \chi)f_- + \chi f_+$. It is easily verified that f satisfies

$$\begin{cases} (\partial_t - \partial_x^2 - q(x)^2 \partial_y^2) f(t, x, y) = \mathbb{1}_\omega u(t, x, y), & t \in (0, T), x \in I, y \in \mathbb{T}, \\ f(t, x, y) = 0, & t \in (0, T), x \in \partial I, y \in \mathbb{T}, \\ f(0, x, y) = f_0(x, y), & x \in I, y \in \mathbb{T}, \\ f(T, x, y) = 0, & x \in I, y \in \mathbb{T}, \end{cases}$$

with a source term $u \in L^2((0, T); L^2(\omega))$. \square

3. Lack of null-controllability

In this section, we prove the following case of Theorem 1.3.

THEOREM 3.1. — *Assume that $q \in C^2(\bar{I})$ is such that $q(0) = 0$, $q'(0) > 0$ and $q(x) \neq 0$ whenever $x \neq 0$. Let ω be an open subset of $I \times \mathbb{T}$. Assume that there exist $a \in (-L_-, 0)$ and $y_0 \in \mathbb{T}$ such that*

$$\text{distance}((a, L_+) \times \{y_0\}, \omega) > 0.$$

Then, the generalized Baouendi–Grushin equation (1.2) is not null-controllable on ω in time $T < d_{\text{Agm}}(a)/q'(0)$.

Remark 3.2. — By changing x in $-x$, I in $-I$ and q in $-q$, this theorem implies that if $b \in (0, L_+)$ and if $\text{distance}((-L_-, b) \times \{y_0\}, \omega) > 0$, then the generalized Baouendi–Grushin equation (1.2) is not null-controllable on ω in time $T < d_{\text{Agm}}(b)/q'(0)$.

To completely prove Theorem 1.3, there are two more cases:

- $\text{distance}((a, b) \times \{y_0\}, \omega) > 0$ with $-L_- < a < 0 < b < L_+$, lack of null-controllability in time $T < \min(d_{\text{Agm}}(a), d_{\text{Agm}}(b))/q'(0)$,
- $\text{distance}((-L_-, L_+) \times \{y_0\}, \omega) > 0$, lack of null-controllability in any time $T > 0$.

The proofs of these cases are minor modifications of the one of Theorem 3.1. We mention in footnotes the most important modifications and leave the details to the reader.

Under the hypotheses of this theorem, there exists a closed interval W_0 that is a neighborhood of y_0 and such that $\omega \cap ([a, L_+) \times W_0) = \emptyset$ (see Figure 3.1). To prove Theorem 3.1, we assume without loss of generality that ω is the complement of the rectangle $[a, L_+) \times W_0$:

$$(3.1) \quad \omega = (I \times \mathbb{T}) \setminus ([a, L_+) \times W_0).$$




Figure 3.1. In green, the domain ω . If a horizontal segment stays at positive distance from ω , it can be thickened into a rectangle that is disjoint from ω .

3.1. Observability inequality

Using standard duality arguments [Cor07, Theorem 2.44], the null-controllability of the generalized Baouendi–Grushin equation (1.2) is equivalent to the following *observability inequality*: there exists $C > 0$ such that for every $g_0 \in L^2(I \times \mathbb{T})$, the solution g of

$$(3.2) \quad \begin{cases} (\partial_t - \partial_x^2 - q(x)^2 \partial_y^2) g(t, x, y) = 0, & t \in (0, T), x \in I, y \in \mathbb{T}, \\ g(t, x, y) = 0, & t \in (0, T), x \in \partial I, y \in \mathbb{T}, \\ g(0, x, y) = g_0(x, y) & (x, y) \in I \times \mathbb{T}, \end{cases}$$

satisfies

$$(3.3) \quad \|g(T, \cdot, \cdot)\|_{L^2(I \times \mathbb{T})}^2 \leq C \|g\|_{L^2((0, T) \times \omega)}^2.$$

To prove Theorem 3.1, we proceed in two steps: we prove that the observability inequality (3.3) implies an inequality on polynomials, and then we disprove this new inequality.⁽⁵⁾

3.2. Model case

We start with a model equation, that we study to showcase the main ideas of the proof of Theorem 3.1 without some of the more technical aspects. Consider the Baouendi–Grushin equation on $\mathbb{R} \times \mathbb{T}$:

$$(3.4) \quad (\partial_t - \partial_x^2 - x^2 \partial_y^2) g(t, x, y) = 0.$$

Let $\omega \subset \mathbb{R} \times \mathbb{T}$ be open. We say that the Baouendi–Grushin equation (3.4) is observable on ω in time $T > 0$ if there exists $C > 0$ such that for all g solution of (3.4), the following observability inequality holds:

$$(3.5) \quad \|g(T, \cdot, \cdot)\|_{L^2(\mathbb{R} \times \mathbb{T})}^2 \leq C \|g\|_{L^2((0, T) \times \omega)}^2.$$

We prove the following theorem.

⁽⁵⁾ Actually, we could reformulate this proof to directly construct a counterexample to the observability inequality (3.3).

THEOREM 3.3. — *Let $a > 0$, $W_0 \subset \mathbb{T}$ a closed interval with non-empty interior and*

$$(3.6) \quad \omega = (\mathbb{R} \times \mathbb{T}) \setminus ([-a, a] \times W_0).$$

Let $T > 0$ such that

$$T < \frac{a^2}{2}.$$

The Baouendi–Grushin equation (3.4) is not observable on ω in time T .

Before going into the proof, let us examine some solutions of the Baouendi–Grushin equation that are concentrated around $x = 0$. Taking the n^{th} Fourier coefficient in y of g , which we will denote by $\widehat{g}(t, x, n)$, we get

$$(3.7) \quad (\partial_t - \partial_x^2 + n^2 x^2) \widehat{g}(t, x, n) = 0.$$

Thus, the Baouendi–Grushin equation is transformed into a family of parabolic equations $(\partial_t + H_n)g_n = 0$, where H_n is the *harmonic oscillator* $-\partial_x^2 + n^2 x^2$. The spectral properties of the harmonic oscillator are well-known (see, e.g., [Hel13, § 1.3] or Appendix C), and in particular the first eigenvalue is $|n|$ with associated eigenfunction $\varphi_n(x) = (|n|/\pi)^{1/4} e^{-|n|x^2/2}$. Thus, if $(a_n)_{n>0}$ is a complex-valued sequence with only a finite number of nonzero terms, the function g defined by

$$(3.8) \quad g(t, x, y) := \sum_{n>0} a_n e^{iny - nx^2/2 - nt}$$

is a solution of the Baouendi–Grushin equation (3.4). We will look for a counterexample of the observability inequality (3.5) in this class of functions.

This solution can be written as $g(t, x, y) = g_{\text{pol}}(e^{iy-t-x^2/2})$ with

$$(3.9) \quad g_{\text{pol}}(z) := \sum_{n>0} a_n z^n.$$

We will use the fact that g is a polynomial in $z = e^{iy-t-x^2/2}$ to rewrite the observability inequality we want to disprove as an inequality on polynomials. More precisely, we have the following estimate.




Figure 3.2. The domain U .

LEMMA 3.4. — Assume that the observability inequality for the Baouendi–Grushin equation (3.5) holds. Let $U \subset \mathbb{C}$ be defined by (see Figure 3.2)

$$U = D\left(0, e^{-a^2/2}\right) \cup \{z \in \mathbb{C} : |z| < 1, \arg(z) \notin W_0\}.$$

Then, there exists $C > 0$ such that for every polynomial $p \in \mathbb{C}[X]$,

$$\|p\|_{L^2(D(0, e^{-T}))} \leq C \|p\|_{L^\infty(U)}.$$

Proof. —

Step 1: Observability inequality. Let $p(z) = \sum_{n \geq 0} a_n z^n$ a polynomial and set $g_{\text{pol}}(z) = zp(z) = \sum_{n > 0} a_{n-1} z^n$. The discussion above shows that g defined by

$$g(t, x, y) = g_{\text{pol}}\left(e^{iy-t-x^2/2}\right)$$

is a solution of the Baouendi–Grushin equation (3.4). The observability inequality on this class of functions reads

$$(3.10) \quad \int_{\mathbb{R} \times \mathbb{T}} |g(T, x, y)|^2 dx dy \leq C \int_{[0, T] \times \omega} \left| g_{\text{pol}}\left(e^{iy-t-x^2/2}\right) \right|^2 dt dx dy.$$

Step 2: Left-hand side of the observability inequality (3.10). Since the functions $\psi_n : (x, y) \mapsto e^{iny-nx^2/2}$ are orthogonal in $L^2(\mathbb{R} \times \mathbb{T})$, the left-hand side can be rewritten as

$$\begin{aligned} \int_{\mathbb{R} \times \mathbb{T}} |g(T, x, y)|^2 dx dy &= \int_{\mathbb{R} \times \mathbb{T}} \left| \sum_{n > 0} a_{n-1} e^{-nT} \psi_n(x, y) \right|^2 dx dy \\ &= \sum_{n > 0} |a_{n-1}|^2 e^{-2nT} \|\psi_n\|_{L^2(\mathbb{R} \times \mathbb{T})}^2 \\ &= \sum_{n > 0} \frac{2\pi^{3/2}}{\sqrt{n}} |a_{n-1}|^2 e^{-2nT}. \end{aligned}$$

Elementary computations in polar coordinates prove that the functions $z \mapsto z^n$ are orthogonal in $L^2(D(0, R), m)$, where m is the Lebesgue measure on $\mathbb{C} \simeq \mathbb{R}^2$, and that for $R > 0$

$$\|z^n\|_{L^2(D(0, R), m)}^2 = \frac{\pi R^{2n+2}}{n+1}.$$

Thus,

$$\begin{aligned} \|p\|_{L^2(D(0, e^{-T}))}^2 &= \sum_{n \geq 0} \frac{\pi}{n+1} |a_n|^2 e^{-2(n+1)T} \\ (3.11) \quad &\leq \sum_{n > 0} \frac{\pi}{\sqrt{n}} |a_{n-1}|^2 e^{-2nT} \\ &= \frac{1}{2\sqrt{\pi}} \int_{\mathbb{R} \times \mathbb{T}} |g(T, x, y)|^2 dx dy. \end{aligned}$$

Step 3: Right-hand side of the observability inequality (3.10). We write the right-hand side of the observability inequality by making the change of variables $(x, z) = (x, e^{-t+iy-x^2/2})$. We have $dx \, dm(z) = |z|^2 \, dt \, dx \, dy$. Thus, if we denote by $\Omega \subset \mathbb{R} \times \mathbb{C}$ the image of $(0, T) \times \omega$ by this change of variables, we have

$$(3.12) \quad \begin{aligned} \int_{(0,T) \times \omega} |g_{\text{pol}}(e^{-t+iy-x^2/2})|^2 \, dt \, dx \, dy &= \int_{\Omega} |g_{\text{pol}}(z)|^2 |z|^{-2} \, dx \, dm(z) \\ &= \int_{\Omega} |p(z)|^2 \, dx \, dm(z). \end{aligned}$$

By definition, $\Omega = \bigcup_{x \in \mathbb{R}} (\{x\} \times \mathcal{D}_x)$, where

$$\mathcal{D}_x = \left\{ e^{-t+iy-x^2/2}, \, t \in (0, T), (x, y) \in \omega \right\}.$$

We claim that for every $0 < t < T$ and $x \in \mathbb{R}$, $\mathcal{D}_x \subset U$ (this is the reason we defined U this way). Indeed, if $-a < x < a$, and $(x, y) \in \omega$, then, by definition of ω as the complement of $[-a, a] \times W_0$ (eq. (3.6)), we necessarily have $y \notin W_0$. It follows that $e^{-t+iy-x^2/2} \in U$. In the case $x \notin [-a, a]$, we have $x^2/2 > a^2/2$. Then, $|e^{-t+iy-x^2/2}| < e^{-a^2/2}$. It follows again that $e^{-t+iy-x^2/2} \in U$. Thus, using Hölder inequality in eq. (3.12),

$$\begin{aligned} \int_{(0,T) \times \omega} |g_{\text{pol}}(e^{-t+iy-x^2/2})|^2 \, dt \, dx \, dy &= \int_{x \in \mathbb{R}} \int_{\mathcal{D}_x} |p(z)|^2 \, dm(z) \, dx \leq \int_{x \in \mathbb{R}} m(\mathcal{D}_x) \|p\|_{L^\infty(U)}^2 \, dx. \end{aligned}$$

Since $\mathcal{D}_x \subset D(0, e^{-x^2/2})$, $m(\mathcal{D}_x) \leq \pi e^{-x^2}$. Hence

$$(3.13) \quad \int_{(0,T) \times \omega} |g_{\text{pol}}(e^{-t+iy-x^2/2})|^2 \, dt \, dx \, dy \leq \pi^{3/2} \|p\|_{L^\infty(U)}^2.$$

Now, plugging the lower-bound of the left-hand side (3.11) and the upper bound of the right-hand side (3.13) into the observability inequality (3.10), we obtain a constant $C > 0$ such that

$$\|p\|_{L^2(D(0, e^{-T}))}^2 \leq C \|p\|_{L^\infty(U)}^2. \quad \square$$

Proof of Theorem 3.3. — To disprove the inequality, we only have to disprove the inequality on polynomials given by the previous Lemma 3.4. If $T < a^2/2$, the disk $D(0, e^{-T})$ is not included in U (see Figure 3.3). For instance, if $y_1 \in \dot{W}_0$ and $\varepsilon > 0$ is small enough, $z_0 = e^{iy_1 - T - \varepsilon}$ is not in U . In fact, the half-line $z_0[1, +\infty)$ stays at positive distance from U (see Figure 3.3). Then, according to Runge's theorem [Rud86, Theorem 13.9], there exists a sequence of polynomials (p_k) that converges uniformly on every compact of $\mathbb{C} \setminus z_0[1, +\infty)$ to $z \mapsto (z - z_0)^{-1}$.

Since \overline{U} is a compact subset of $\mathbb{C} \setminus z_0[1, +\infty)$, the sequence p_k is uniformly bounded on U , i.e., $\sup_k \|p_k\|_{L^\infty(U)} < +\infty$. Recall that $z_0 \in D(0, e^{-T})$, which implies that $\|(z - z_0)^{-1}\|_{L^2(D(0, e^{-T}))} = +\infty$. Thanks to Fatou's lemma, this proves that $\|p_k\|_{L^2(D(0, e^{-T}))} \rightarrow +\infty$ as $k \rightarrow +\infty$.

We have proved that (p_k) is a counterexample to the inequality of Lemma 3.4, which concludes the proof of Theorem 3.3. \square




Figure 3.3. When the disk $D(0, e^{-T})$ (in red) is not included in U , we can find holomorphic functions that are small in U but arbitrarily large in $D(0, e^{-T})$. For instance, we can construct with Runge's theorem a sequence of polynomials that converges to $z \mapsto (z - z_0)^{-1}$ away from the blue line.

3.3. From the model case to the generalized Baouendi–Grushin equation

Now, our goal is to adapt the strategy used in the model case to the generalized Baouendi–Grushin equation (1.2). In the generalized Baouendi–Grushin equation, if we take the n^{th} Fourier coefficient in y of g , we get

$$(3.14) \quad (\partial_t - \partial_x^2 + n^2 q(x)^2) \hat{g}(t, x, n) = 0.$$

Recall that for $n \geq 0$, P_n is the unbounded operator $-\partial_x^2 + n^2 q^2$ on $L^2(I)$ with Dirichlet boundary conditions. We will denote by λ_n the first eigenvalue of P_n and by φ_n a corresponding eigenfunction. Notice that φ_n is not required to be normalized in $L^2(I)$. Then, we will look for a counterexample of the observability inequality (3.3) with solutions of the generalized Baouendi–Grushin equation (3.2) of the form

$$(3.15) \quad g(t, x, y) := \sum_{n \geq 0} a_n \varphi_n(x) e^{iny - \lambda_n t}.$$

Heuristically, this should work because we expect the eigenfunction φ_n to be localized around $x = 0$ as $n \rightarrow +\infty$, in which case the operator $-\partial_x^2 + n^2 q^2$ looks like $-\partial_x^2 + n^2 q'(0)^2 x^2$, and the eigenvalue and eigenfunction look like $\lambda_n \approx n q'(0)$ and $\varphi_n(x) \approx n^{1/4} e^{-n q'(0) x^2 / 2}$. So the solutions g defined above look like the solutions used to treat the model case (eq. (3.8)), up to a factor $q'(0)$.

In fact, a better approximation of φ_n would be the so-called WKB approximation⁽⁶⁾

⁽⁶⁾We write here the first term in the WKB expansion of eigenfunctions, and only in dimension 1, because it is enough for our purposes. But such a construction can be refined with more terms and in higher dimension [DS99, Chapter 3 & Chapter 6, Theorem A.3].

Also, in the differential equation that defines c_0 , we divide by $d'_{\text{Agm}}(x)$, which is equal to 0 at $x = 0$. But the numerator is also 0 at $x = 0$, and simple Taylor expansions at $x = 0$ prove that the quotient appearing in the differential equation for c_0 is actually well-defined at $x = 0$.

$$\begin{aligned}
 \varphi_n(x) &\approx n^{1/4} c_0(x) e^{-n d_{\text{Agm}}(x)}, \\
 c'_0(x) &= \frac{q'(0) - d''_{\text{Agm}}(x)}{2 d'_{\text{Agm}}(x)} c_0(x), \\
 c_0(0) &= 1.
 \end{aligned}
 \tag{3.16}$$

Thus, we have

$$g(t, x, y) \approx c_0(x) \sum_n a_n e^{n(iy - q'(0)t - d_{\text{Agm}}(x))}, \tag{3.17}$$

i.e., g can almost be written as $g(t, x, y) \approx c_0(x) g_{\text{pol}}(e^{iny - tq'(0) - d_{\text{Agm}}(x)})$, where g_{pol} is the polynomial

$$g_{\text{pol}}(z) := \sum_n a_n z^n.$$

Let us write this in an exact way. Consider $\tilde{\varphi}_n(x) := n^{1/4} e^{-nq'(0)x^2/2}$ and let Π_n be the spectral projection associated to the first eigenvalue λ_n of P_n . We define⁽⁷⁾

$$\varphi_n := \Pi_n \tilde{\varphi}_n. \tag{3.18}$$

We will see later that $\varphi_n \neq 0$, at least if n is large enough. Let $\varepsilon \in (0, 1)$, that we need for technical reason, and that we will later choose close to 0. We define $\gamma_{t,x}(n)$ by

$$\gamma_{t,x}(n-1) := e^{-t(\lambda_n - q'(0)n)} \varphi_n(x) e^{n d_{\text{Agm}}(x)(1-\varepsilon)}. \tag{3.19}$$

The shift of n in the definition is linked to the fact that we will consider $p(z) = g_{\text{pol}}(z)/z$, as we did in the model case. Then, the solution g defined in eq. (3.15) can be written as

$$g(t, x, y) = \sum_n a_n \gamma_{t,x}(n-1) e^{n(iy - q'(0)t - (1-\varepsilon) d_{\text{Agm}}(x))}. \tag{3.20}$$

In some sense, this formula tells us that g can be written as “pseudo-differential-type” operator applied to the “model solution” $g_{\text{pol}}(e^{iy - q'(0)t - (1-\varepsilon) d_{\text{Agm}}(x)})$. To successfully adapt the strategy used for the model Baouendi–Grushin equation, we need some continuity estimates for these “pseudo-differential-type” operators. We claim that the following estimate holds.

LEMMA 3.5. — *Let $T > 0$ and $\varepsilon > 0$. Define $\gamma_{t,x}$ as in eq. (3.19). Let $\gamma_{t,x}(z\partial_z)$ be the operator on polynomials defined by*

$$\gamma_{t,x}(z\partial_z) \left(\sum a_n z^n \right) = \sum \gamma_{t,x}(n) a_n z^n.$$

Let X be a compact subset of \mathbb{C} . Let V be an open neighborhood of X that is star-shaped with respect to 0. There exist $C > 0$ and $N \in \mathbb{N}$ such that for every polynomial $p \in \mathbb{C}[X]$ with a zero of order N at 0 and for every $0 < t < T$ and $x \in I$,

$$\|\gamma_{t,x}(z\partial_z)(p)\|_{L^\infty(X)} \leq C \|p\|_{L^\infty(V)}.$$

⁽⁷⁾We could also have chosen $\tilde{\varphi}_n$ to be the WKB expansion defined previously, which would be a better approximation of the eigenfunction. But since we are projecting on the actual eigenfunction afterwards, this is not necessary.




Figure 3.4. The domains U and V .

As $\gamma_{t,x}(n)$ is related to the eigenvalues and eigenfunctions of P_n , proving this lemma requires a spectral analysis of this operator. What is more surprising is that we actually need a spectral analysis of P_ν when ν is not necessarily real, meaning we have to do some *nonselfadjoint* spectral analysis. We will prove Lemma 3.5 in Section 4.3 with the spectral analysis done in the rest of Section 4 and a general estimate on operators on polynomials [Koe17, Theorem 18].

We will also use the relatively elementary bounds on λ_n and $\|\varphi_n\|_{L^2(I)}$ given by the following proposition:

PROPOSITION 3.6. — *In the limit $n \rightarrow +\infty$, $\lambda_n = nq'(0) + o(n)$. Moreover, there exist $c > 0$ and $N \geq 0$ such that for every $n \geq N$, $\|\varphi_n\|_{L^2(I)} \geq c$.*

This proposition is standard (see, e.g., [DS99, Theorem 4.23 & Eq. (4.20)]), nevertheless, for the reader convenience, we provide a proof in Section 4.2.

With these two estimates, we prove the following version of Lemma 3.4 adapted for the generalized Baouendi–Grushin equation.

LEMMA 3.7. — *Assume that the observability inequality (3.3) for the generalized Baouendi–Grushin equation holds. Let $\varepsilon > 0$ and let $U \subset \mathbb{C}$ be defined by (see Figure 3.4)⁽⁸⁾*

$$U = D\left(0, e^{-(1-\varepsilon) d_{\text{Agm}}(a)}\right) \cup \{z \in \mathbb{C} : |z| < 1, \arg(z) \notin W_0\}.$$

Let V be a neighborhood of \overline{U} that is star-shaped with respect to 0. Then, there exist $C > 0$ and $N \in \mathbb{N}$ such that for every polynomial $p \in \mathbb{C}[X]$ with a zero of order N at 0, we have

$$\|p\|_{L^2(D(0, e^{-q'(0)T(1+\varepsilon)})} \leq C \|p\|_{L^\infty(V)}.$$

⁽⁸⁾In the variant of Theorem 3.1 where $\text{distance}((a, b) \times \{y_0\}, \omega) > 0$ mentioned in Remark 3.2, we have to add $D(0, e^{-(1-\varepsilon) d_{\text{Agm}}(b)})$ to U . In the variant where $\text{distance}(I \times \{y_0\}, \omega) > 0$, U is only the pacman $\{z \in \mathbb{C} : \arg(z) \notin W_0\}$. Their proofs are minor adaptations and are left to the reader.

Proof. — The proof mostly follows the one of Lemma 3.4, but with the error term $\gamma_{t,x}$ which will be handled by Lemma 3.5. Let $N > 0$ as in Lemma 3.5 and Proposition 3.6. Let $p(z) = \sum_{n \geq N} a_n z^n$ a polynomial and $g_{\text{pol}}(z) = zp(z)$. The discussion above shows that g defined by

$$(3.21) \quad \begin{aligned} g(t, x, y) &= \sum_{n > N} a_{n-1} \varphi_n(x) e^{-\lambda_n t + i n y} \\ &= \sum_{n > N} a_{n-1} \gamma_{t,x}(n-1) e^{n(iy - q'(0)t - (1-\varepsilon) d_{\text{Agm}}(x))} \end{aligned}$$

is a solution of the Baouendi–Grushin equation (3.2).

Step 1: Left-hand side of the observability inequality (3.3). Since the functions $\psi_n: (x, y) \mapsto \varphi_n(x) e^{i n y}$ are orthogonal, the left-hand side can be rewritten as

$$\begin{aligned} \int_{I \times \mathbb{T}} |g(T, x, y)|^2 dx dy &= \int_{I \times \mathbb{T}} \left| \sum_{n \geq N} a_{n-1} e^{-\lambda_n T} \psi_n(x, y) \right|^2 dx dy \\ &= \sum_{n \geq N} |a_{n-1}|^2 e^{-2\lambda_n T} \|\psi_n\|_{L^2(I \times \mathbb{T})}^2, \end{aligned}$$

using the lower bounds on $\|\varphi_n\|_{L^2(I)}$ given by Proposition 3.6, we get $\|\psi_n\|_{L^2(I \times \mathbb{T})} \geq c > 0$ for $n \geq N$. Thus,

$$\int_{I \times \mathbb{T}} |g(T, x, y)|^2 dx dy \geq c \sum_{n \geq N} |a_{n-1}|^2 e^{-2\lambda_n T}.$$

Now, thanks to the asymptotics for λ_n given by Proposition 3.6, there exists $C_\varepsilon > 0$ such that $\lambda_n \leq nq'(0)(1 + \varepsilon) + C_\varepsilon$. Thus,

$$\int_{I \times \mathbb{T}} |g(T, x, y)|^2 dx dy \geq c e^{-2TC_\varepsilon} \sum_{n \geq N} |a_{n-1}|^2 e^{-2nq'(0)T(1+\varepsilon)}.$$

As in the proof of Lemma 3.4, we denote by m the Lebesgue measure on $\mathbb{C} \simeq \mathbb{R}^2$, the functions $z \mapsto z^n$ are orthogonal on $L^2(D(0, R), m)$ and $\|z^n\|_{L^2(D(0, R), m)}^2 = \pi R^{2n+2}/(n+1)$. Thus,

$$(3.22) \quad \begin{aligned} \|p\|_{L^2(D(0, e^{-q'(0)T(1+\varepsilon)}))}^2 &= \sum_{n \geq N} \frac{\pi}{n+1} |a_n|^2 e^{-2(n+1)q'(0)T(1+\varepsilon)} \\ &\leq \sum_{n > N} \pi |a_{n-1}|^2 e^{-2nq'(0)T(1+\varepsilon)} \\ &\leq C \int_{I \times \mathbb{T}} |g(T, x, y)|^2 dx dy. \end{aligned}$$

Step 2: Right-hand side of the observability inequality (3.3). We make the analogous change of variables as in the model case, but adapted to our case, i.e., $(x, z) = (x, e^{-q'(0)t + i y - (1-\varepsilon) d_{\text{Agm}}(x)})$. We have $dx dm(z) = q'(0)|z|^2 dt dx dy$. Thus, if we denote by $\Omega \subset I \times \mathbb{C}$ the image of $(0, T) \times \omega$ by this change of variables, which is a subset of $I \times D(0, 1)$, we have

$$\begin{aligned}
(3.23) \quad & \int_{(0,T) \times \omega} |g(t, x, y)|^2 dt dx dy \\
&= \frac{1}{q'(0)} \int_{\Omega} \left| \sum_{n > N} a_{n-1} \gamma_{t,x}(n-1) z^n \right|^2 |z|^{-2} dx dm(z) \\
&= \frac{1}{q'(0)} \int_{\Omega} |\gamma_{t,x}(z \partial_z)(p)(z)|^2 dx dm(z).
\end{aligned}$$

We keep for simplicity the notation $\gamma_{t,x}$ but of course t is now a function of (x, z) . As in the model case, if $(x, z) \in \Omega$, then $z \in U$. Indeed, let $(x, z) \in \Omega$, i.e., $z = e^{-t+iy-d_{\text{Agm}}(x)(1-\varepsilon)}$ with $(x, y) \in \omega$. If $a \leq x < L_+$, then, by definition of ω as the complement of $[a, L_+) \times W_0$ (eq. (3.1)), we necessarily have $y \notin W_0$. It follows that $z \in U$. In the case $x < a < 0$, since d_{Agm} is decreasing on $[-L_-, 0]$, we have $|z| = e^{-t-d_{\text{Agm}}(x)(1-\varepsilon)} < e^{-d_{\text{Agm}}(a)(1-\varepsilon)}$. It follows again that $z \in U$. Thus,

$$\int_{(0,T) \times \omega} |g(t, x, y)|^2 dt dx dy \leq C \sup_{(t,x) \in (0,T) \times I} \|\gamma_{t,x}(z \partial_z)(p)\|_{L^\infty(U)}^2.$$

Now, we use the operator estimate of Lemma 3.5, which gives

$$(3.24) \quad \int_{(0,T) \times \omega} |g(t, x, y)|^2 dt dx dy \leq C \|p\|_{L^\infty(V)}^2.$$

Step 3: Conclusion. Now, plugging the lower-bound of the left-hand side (3.22) and the upper bound of the right-hand side (3.24) into the observability inequality (3.3), we get

$$\|p\|_{L^2(D(0, e^{-q'(0)T(1+\varepsilon)}))}^2 \leq C \|p\|_{L^\infty(V)}^2. \quad \square$$




Figure 3.5. When the disk $D(0, e^{-q'(0)T(1+\varepsilon)})$ (in red) is not included in U , we can find holomorphic functions that are small in U but arbitrarily big in $D(0, e^{-q'(0)T(1+\varepsilon)})$. For instance, we can construct with Runge's theorem a sequence of polynomials that converges to $z \mapsto z^{N+1}(z - z_0)^{-1}$ away from the blue line.

Proof of Theorem 3.1. — As in the model case, we end the proof of non-null controllability by disproving the inequality on polynomials given by the previous Lemma 3.7. If

$$q'(0)T(1+\varepsilon) < d_{\text{Agm}}(a)(1-\varepsilon),$$

the disk $D(0, e^{-q'(0)T(1+\varepsilon)})$ is not included⁽⁹⁾ in \overline{U} , and we can choose a compact neighborhood V of \overline{U} that is star-shaped with respect to 0 and such that $D(0, e^{-q'(0)T(1+\varepsilon)})$ is not included in \overline{V} (see Figure 3.5). Choose some $z_0 \in D(0, e^{-q'(0)T(1+\varepsilon)})$ that is not in \overline{V} . Since V is star-shaped with respect to 0, the half-line $z_0[1, +\infty)$ stays at positive distance from V . Then, according to Runge's theorem [Rud86, Theorem 13.9], there exists a sequence of polynomials $(\tilde{p}_k)_k$ that converges uniformly on every compact of $\mathbb{C} \setminus z_0[1, +\infty)$ to $z \mapsto (z - z_0)^{-1}$.

Set $p_k(z) := z^{N+1}\tilde{p}_k$. We prove that (p_k) is a counterexample to the inequality of Lemma 3.7 with the same method as in the model case. Since \overline{V} is a compact subset of $\mathbb{C} \setminus z_0[1, +\infty)$, p_k stays uniformly bounded on V , i.e., $\sup_k \|p_k\|_{L^\infty(V)} < +\infty$. But $z_0 \in D(0, e^{-q'(0)T(1+\varepsilon)})$, and therefore $\|z^{N+1}(z - z_0)^{-1}\|_{L^2(D(0, e^{-q'(0)T(1+\varepsilon)}))} = +\infty$. Thanks to Fatou's lemma, this proves that $\|p_k\|_{L^2(D(0, e^{-q'(0)T(1+\varepsilon)}))} \rightarrow +\infty$ as $k \rightarrow +\infty$.

We have proved that the inequality of Lemma 3.7 does not hold, which implies that the observability inequality (3.3) does not hold either, which in turn implies that the generalized Baouendi–Grushin equation is not null-controllable. This holds for any $\varepsilon > 0$ and any T such that $(1+\varepsilon)q'(0)T < d_{\text{Agm}}(a)(1-\varepsilon)$. Thus, the generalized Baouendi–Grushin equation is not null controllable if $T < d_{\text{Agm}}(a)/q'(0)$. \square

4. Spectral Analysis

As explained in Section 3.3, we need some spectral properties on the operator $P_\nu = -\partial_x^2 + \nu^2 q(x)^2$ with Dirichlet boundary conditions on I (defined precisely in eq. (1.4)). We start with an asymptotic of the first eigenvalue, and in following subsection, we prove some Agmon-type upper bound for the associated eigenfunctions.

For $\theta_0 \in [0, \frac{\pi}{2})$, we set

$$(4.1) \quad \Sigma_{\theta_0} := \{\nu \in \mathbb{C} : |\nu| \geq 1, |\arg(\nu)| \leq \theta_0\}.$$

4.1. The first eigenvalue and corresponding spectral projection

For $\beta \in \mathbb{C}$ with $\text{Re}(\beta) > 0$, we denote by H_β the non-selfadjoint harmonic oscillator $-\partial_x^2 + \beta^2 x^2$ on \mathbb{R} . We refer to Appendix C for the precise definition and properties of H_β .

In this paragraph we prove that the operator P_ν has an eigenvalue close to the eigenvalue $q'(0)\nu$ of the model operator $H_{q'(0)\nu}$, and that the corresponding spectral

⁽⁹⁾ In the variant of Theorem 3.1 where $\text{distance}((a, b) \times \{y_0\}, \omega) > 0$ mentioned in Remark 3.2, taking into account Footnote 8, the condition becomes $q'(0)T(1+\varepsilon) < \min(d_{\text{Agm}}(a), d_{\text{Agm}}(b))(1-\varepsilon)$. In the variant where $\text{distance}(I \times \{y_0\}, \omega) > 0$, again taking into account Footnote 8, $D(0, e^{-q'(0)T(1+\varepsilon)})$ is never included in U .

projection is also a perturbation of the spectral projection of $H_{q'(0)\nu}$. See Proposition 4.2.

For this we first prove that the resolvent of P_ν is a perturbation of the resolvent of $H_{q'(0)\nu}$, in the sense that the difference between these two resolvents is smaller than the resolvent of $H_{q'(0)\nu}$.

Notice that the resolvents of P_ν and $H_{q'(0)\nu}$ are not defined on the same space. We denote by $\mathbb{1}_I$ the operator that maps a function $v \in L^2(I)$ to its extension by 0 on \mathbb{R} . Then $\mathbb{1}_I^*$ is the operator which maps $u \in L^2(\mathbb{R})$ to its restriction on I : $\mathbb{1}_I^*u = u|_I \in L^2(I)$.

PROPOSITION 4.1. — *Let $\theta_0 \in [0, \frac{\pi}{2})$. Let $\gamma > 0$ and $\varepsilon > 0$. For $\nu \in \Sigma_{\theta_0}$ we set (see eq. (C.2))*

$$(4.2) \quad \begin{aligned} \tilde{\mathcal{Z}}_\nu &= \mathcal{Z}_{q'(0)\nu, \varepsilon, \gamma} \\ &= \left\{ z \in \mathbb{C} : |z| \leq \gamma q'(0) |\nu|, \text{ distance}(z, \sigma(H_{q'(0)\nu})) \geq \varepsilon q'(0) |\nu| \right\}. \end{aligned}$$

Then there exists $\nu_0 \geq 1$ such that for $\nu \in \Sigma_{\theta_0}$ with $|\nu| \geq \nu_0$ and $z \in \tilde{\mathcal{Z}}_\nu$ we have $z \in \rho(P_\nu)$, and

$$\sup_{z \in \tilde{\mathcal{Z}}_\nu} \left\| (P_\nu - z)^{-1} - \mathbb{1}_I^* (H_{q'(0)\nu} - z)^{-1} \mathbb{1}_I \right\|_{\mathcal{L}(L^2(I))} = \underset{\substack{|\nu| \rightarrow +\infty \\ \nu \in \Sigma_{\theta_0}}}{o} \left(\frac{1}{|\nu|} \right).$$

Proof. — For $\nu \in \Sigma_{\theta_0}$ and $z \in \tilde{\mathcal{Z}}_\nu$ we set

$$R_\nu(z) = \mathbb{1}_I^* (H_{q'(0)\nu} - z)^{-1} \mathbb{1}_I \in \mathcal{L}(L^2(I)).$$

Let

$$\rho \in \left] \frac{1}{3}, \frac{1}{2} \right[.$$

We consider a cut-off function $\chi \in C_0^\infty(\mathbb{R}, [0, 1])$ supported in $[-2, 2]$ and equal to 1 on $[-1, 1]$. Then for $\nu \in \Sigma_{\theta_0}$ and $x \in \bar{I}$ we set

$$\chi_\nu(x) = \chi(|\nu|^\rho x).$$

Step 1: Approximation close to $x = 0$. We first prove that if $|\nu|$ is large enough then $R_\nu(z)\chi_\nu(P_\nu - z) - \chi_\nu$ extends to a bounded operator on $L^2(I)$ for all $z \in \tilde{\mathcal{Z}}_\nu$, and

$$(4.3) \quad \sup_{z \in \tilde{\mathcal{Z}}_\nu} \|R_\nu(z)\chi_\nu(P_\nu - z) - \chi_\nu\|_{\mathcal{L}(L^2(I))} \xrightarrow{|\nu| \rightarrow +\infty} 0.$$

Here and everywhere below it is implicitly understood that ν always belongs to Σ_{θ_0} .

Let $u \in \text{Dom}(P_\nu)$. We have $\chi_\nu u \in \text{Dom}(P_\nu)$ and, if $|\nu|$ is large enough, $\mathbb{1}_I \chi_\nu u$ belongs to $\text{Dom}(H_{q'(0)\nu})$. For $x \in \bar{I}$ we set

$$r(x) = q(x)^2 - q'(0)^2 x^2,$$

so that, for $|\nu|$ large enough,

$$R_\nu(z)(P_\nu - z)\chi_\nu u = \chi_\nu u + \nu^2 R_\nu(z)r\chi_\nu u.$$

The commutator $[\chi_\nu, P_\nu]$ of χ_ν and P_ν is equal to $[\chi_\nu, P_\nu] = [\partial_x^2, \chi_\nu] = \chi_\nu'' + 2\chi_\nu' \partial_x$, hence

$$(4.4) \quad \begin{aligned} R_\nu(z) \chi_\nu (P_\nu - z) u &= R_\nu(z) (P_\nu - z) \chi_\nu u + R_\nu(z) [\chi_\nu, P_\nu] u \\ &= \chi_\nu u + \nu^2 R_\nu(z) r \chi_\nu u + R_\nu(z) \chi_\nu'' u + 2R_\nu(z) (\chi_\nu' u)' . \end{aligned}$$

By the resolvent estimate (C.3), we have

$$(4.5) \quad \forall \nu \in \Sigma_{\theta_0}, \forall z \in \tilde{\mathcal{Z}}_\nu, \quad \left\| \left(H_{q'(0)\nu} - z \right)^{-1} \right\|_{\mathcal{L}(L^2(\mathbb{R}))} \leq \frac{C}{|\nu|}.$$

Since $|r(x)\chi_\nu(x)| \lesssim |\nu|^{-3\rho}$, this gives

$$\left\| \nu^2 R_\nu(z) r \chi_\nu \right\|_{\mathcal{L}(L^2(I))} \lesssim |\nu|^{2-1-3\rho} \xrightarrow{|\nu| \rightarrow +\infty} 0.$$

Similarly,

$$\left\| R_\nu(z) \chi_\nu'' \right\|_{\mathcal{L}(L^2(I))} \lesssim |\nu|^{2\rho-1} \xrightarrow{|\nu| \rightarrow +\infty} 0.$$

Considering the last term in eq. (4.4), we have for $v \in L^2(\mathbb{R})$

$$(4.6) \quad \begin{aligned} \left\| \partial_x \left(H_{q'(0)\bar{\nu}} - \bar{z} \right)^{-1} v \right\|_{L^2(\mathbb{R})}^2 + \bar{\nu}^2 q'(0)^2 \left\| x \left(H_{q'(0)\bar{\nu}} - \bar{z} \right)^{-1} v \right\|_{L^2(\mathbb{R})}^2 \\ = \left\langle H_{q'(0)\bar{\nu}} \left(H_{q'(0)\bar{\nu}} - \bar{z} \right)^{-1} v, \left(H_{q'(0)\bar{\nu}} - \bar{z} \right)^{-1} v \right\rangle_{L^2(\mathbb{R})} \\ = \left\langle v, \left(H_{q'(0)\bar{\nu}} - \bar{z} \right)^{-1} v \right\rangle_{L^2(\mathbb{R})} + \bar{z} \left\| \left(H_{q'(0)\bar{\nu}} - \bar{z} \right)^{-1} v \right\|_{L^2(\mathbb{R})}^2 . \end{aligned}$$

We multiply by $e^{i\theta}$ and take the real part. This gives, uniformly in $\nu \in \Sigma_{\theta_0}$ and $z \in \tilde{\mathcal{Z}}_\nu$,

$$(4.7) \quad \left\| \partial_x \left(H_{q'(0)\bar{\nu}} - \bar{z} \right)^{-1} v \right\|_{\mathcal{L}(L^2(\mathbb{R}))}^2 \lesssim \frac{1}{|\nu|} \quad \text{and} \quad \left\| x \left(H_{q'(0)\bar{\nu}} - \bar{z} \right)^{-1} v \right\|_{\mathcal{L}(L^2(\mathbb{R}))}^2 \lesssim \frac{1}{|\nu|^3}.$$

Taking the adjoint in the first inequality gives, for $|\nu|$ large enough,

$$\begin{aligned} \left\| R_\nu(z) \partial_x (\chi_\nu' u) \right\|_{L^2(I)} &\leq \left\| \left(H_{q'(0)\nu} - z \right)^{-1} \partial_x (\mathbf{1}_I \chi_\nu' u) \right\|_{L^2(\mathbb{R})} \\ &\lesssim |\nu|^{-\frac{1}{2}} \left\| \mathbf{1}_I \chi_\nu' u \right\|_{L^2(\mathbb{R})} \\ &\lesssim |\nu|^{\rho-\frac{1}{2}} \|u\|_{L^2(I)}, \end{aligned}$$

and eq. (4.3) follows.

Step 2: Approximation away from $x = 0$. There exists $c_0 \in (0, 1]$ such that for all $x \in I$ we have

$$|q(x)| \geq c_0 |x|.$$

On $L^2(I)$ we consider the operator

$$\tilde{P}_\nu = P_\nu + \nu^2 \mathbf{1}_{[-|\nu|^{-\rho}, |\nu|^{-\rho}]},$$

with domain $\text{Dom}(\tilde{P}_\nu) = H^2(I) \cap H_0^1(I)$. It has compact resolvent, so its spectrum consists of eigenvalues. We have $q(x)^2 + \mathbb{1}_{[-|\nu|^{-\rho}, |\nu|^{-\rho}]} \geq c_0^2 |\nu|^{-2\rho}$ on I . Then, for $u \in \text{Dom}(\tilde{P}_\nu)$ and $z \in \tilde{\mathcal{Z}}_\nu$,

$$\text{Re} \left(e^{-i\theta} \langle (\tilde{P}_\nu - z) u, u \rangle \right) \geq \left(\cos(\theta) c_0^2 |\nu|^{2-2\rho} - \gamma q'(0) |\nu| \right) \|u\|_{L^2(I)}^2.$$

Thus, when $|\nu|$ is so large that $\gamma q'(0) |\nu| \leq \cos(\theta_0) c_0^2 |\nu|^{2-2\rho} / 2$ we have $\tilde{\mathcal{Z}}_\nu \subset \rho(\tilde{P}_\nu)$ and, for $z \in \tilde{\mathcal{Z}}_\nu$,

$$(4.8) \quad \left\| (\tilde{P}_\nu - z)^{-1} \right\|_{\mathcal{L}(L^2(I))} \leq \frac{2 |\nu|^{2\rho-2}}{\cos(\theta) c_0^2}.$$

Then we have

$$\begin{aligned} & (\tilde{P}_\nu - z)^{-1} (1 - \chi_\nu) (P_\nu - z) u \\ &= (\tilde{P}_\nu - z)^{-1} (1 - \chi_\nu) (\tilde{P}_\nu - z) u \\ &= (\tilde{P}_\nu - z)^{-1} (\tilde{P}_\nu - z) (1 - \chi_\nu) u + (\tilde{P}_\nu - z)^{-1} [1 - \chi_\nu, \tilde{P}_\nu] u \\ &= (1 - \chi_\nu) u - 2 (\tilde{P}_\nu - z)^{-1} (\chi'_\nu u)' + (\tilde{P}_\nu - z)^{-1} \chi''_\nu u. \end{aligned}$$

As above we estimate

$$\left\| (\tilde{P}_\nu - z)^{-1} \chi''_\nu \right\|_{\mathcal{L}(L^2(I))} \lesssim |\nu|^{4\rho-2} \xrightarrow{|\nu| \rightarrow +\infty} 0$$

and

$$\left\| (\tilde{P}_\nu - z)^{-1} (\chi'_\nu u)' \right\|_{L^2(I)} \lesssim |\nu|^{\rho-1} \|\chi'_\nu u\|_{L^2(I)} \lesssim |\nu|^{2\rho-1} \|u\|_{L^2(I)}.$$

This proves that

$$(4.9) \quad \sup_{z \in \tilde{\mathcal{Z}}_\nu} \left\| (\tilde{P}_\nu - z)^{-1} (1 - \chi_\nu) (P_\nu - z) - (1 - \chi_\nu) \right\|_{\mathcal{L}(L^2(I))} \xrightarrow{|\nu| \rightarrow +\infty} 0.$$

Step 3: Conclusion. For $\nu \in \Sigma_{\theta_0}$ and $z \in \tilde{\mathcal{Z}}_\nu$ we set

$$Q_\nu(z) = R_\nu(z) \chi_\nu + (\tilde{P}_\nu - z)^{-1} (1 - \chi_\nu).$$

We have for $u \in \text{Dom}(P_\nu)$

$$\begin{aligned} \|Q_\nu(z)(P_\nu - z)u - u\|_{L^2(I)} &\leq \|R_\nu(z) \chi_\nu (P_\nu - z)u - \chi_\nu u\|_{L^2(I)} \\ &\quad + \left\| (\tilde{P}_\nu - z)^{-1} (1 - \chi_\nu) (P_\nu - z)u - (1 - \chi_\nu)u \right\|_{L^2(I)}, \end{aligned}$$

hence by eq. (4.3) and eq. (4.9) we obtain

$$\|Q_\nu(z)(P_\nu - z)u - u\|_{L^2(I)} = o_{|\nu| \rightarrow +\infty}(1) \|u\|_{L^2(I)}.$$

This proves that for $|\nu|$ large enough, we can extend the operator $Q_\nu(z)(P_\nu - z) = 1 + (Q_\nu(z)(P_\nu - z) - 1)$ to a bounded operator on $L^2(I)$, which is invertible with inverse bounded uniformly in ν , and

$$(4.10) \quad \left\| \left(Q_\nu(z)(P_\nu - z) \right)^{-1} - 1 \right\|_{\mathcal{L}(L^2(I))} \xrightarrow{|\nu| \rightarrow +\infty} 0.$$

In particular, $(P_\nu - z)$ is injective. Since it has compact resolvent, it is boundedly invertible, and

$$(P_\nu - z)^{-1} = \left(Q_\nu(z)(P_\nu - z) \right)^{-1} Q_\nu(z).$$

We get

$$(4.11) \quad \begin{aligned} (P_\nu - z)^{-1} - R_\nu(z) &= \left(Q_\nu(z)(P_\nu - z) \right)^{-1} \left(\tilde{P}_\nu - z \right)^{-1} (1 - \chi_\nu) \\ &\quad - \left(Q_\nu(z)(P_\nu - z) \right)^{-1} R_\nu(z) (1 - \chi_\nu) \\ &\quad + \left(\left(Q_\nu(z)(P_\nu - z) \right)^{-1} - 1 \right) R_\nu(z). \end{aligned}$$

We prove that each term of the right-hand side is of size $O(|\nu|^{-1})$. For the first term we use eq. (4.8). For the third we use eq. (4.10) and eq. (4.5). Finally, for the second term we observe that on $\text{supp}(1 - \chi_\nu)$ we have $|x| \gtrsim |\nu|^{-\rho}$ so for $u \in L^2(I)$ we have by the second inequality of eq. (4.7)

$$\begin{aligned} \left\| (1 - \chi_\nu) \mathbb{1}_I^* \left(H_{q'(0)\bar{\nu}} - \bar{z} \right)^{-1} \mathbb{1}_I u \right\|_{L^2(I)}^2 &\lesssim |\nu|^{2\rho} \left\| x \left(H_{q'(0)\bar{\nu}} - \bar{z} \right)^{-1} \mathbb{1}_I u \right\|_{L^2(\mathbb{R})}^2 \\ &\lesssim |\nu|^{2\rho-3} \|u\|_{L^2(I)}^2. \end{aligned}$$

Taking the adjoint gives

$$\left\| \mathbb{1}_I^* \left(H_{q'(0)\nu} - z \right)^{-1} \mathbb{1}_I (1 - \chi_\nu) \right\|_{\mathcal{L}(L^2(I))} = \underset{\nu \rightarrow +\infty}{o} \left(\frac{1}{|\nu|} \right),$$

and the conclusion follows from eq. (4.11). \square

PROPOSITION 4.2. — *Let $\theta_0 \in [0, \frac{\pi}{2})$. There exists $\nu_{\theta_0} \geq 0$ such that for $\nu \in \Sigma_{\theta_0}$ with $|\nu| \geq \nu_{\theta_0}$ the operator P_ν has a unique eigenvalue λ_ν which satisfies $|\lambda_\nu - q'(0)\nu| \leq q'(0)|\nu|$. Moreover, λ_ν is algebraically simple and*

$$|\lambda_\nu - q'(0)\nu| = \underset{\substack{|\nu| \rightarrow +\infty \\ \nu \in \Sigma_{\theta_0}}}{o} (|\nu|).$$

If we denote by $\Pi_\nu \in \mathcal{L}(L^2(I))$ the associated spectral projection, and by $\Pi_{q'(0)\nu}^H \in \mathcal{L}(L^2(\mathbb{R}))$ the spectral projection associated to the eigenvalue $q'(0)\nu$ of $H_{q'(0)\nu}$, then

$$\left\| \Pi_\nu - \mathbb{1}_I^* \Pi_{q'(0)\nu}^H \mathbb{1}_I \right\|_{\mathcal{L}(L^2(I))} \xrightarrow[\nu \in \Sigma_{\theta_0}]{|\nu| \rightarrow +\infty} 0.$$

Proof. — We recall that for $\nu \in \Sigma_{\theta_0}$ we have

$$\Pi_{q'(0)\nu}^H = \frac{1}{2i\pi} \int_{|z - q'(0)\nu| = q'(0)|\nu|} (H_{q'(0)\nu} - z)^{-1} dz.$$

Let ν_0 be given by Proposition 4.1 for $\varepsilon = 1$ and $\gamma = \frac{5}{2}$. If $|\nu| \geq \nu_0$ we can set

$$B_\nu = \frac{1}{2i\pi} \int_{|z - q'(0)\nu| = q'(0)|\nu|} (P_\nu - z)^{-1} dz.$$

This is a projection of $L^2(I)$ whose range is the sum of the generalized eigenspaces of P_ν corresponding to the eigenvalues in the disk $D(q'(0)\nu, q'(0)|\nu|)$. In particular the dimension of $\text{Ran}(B_\nu)$ does not depend on $\nu \in \Sigma_{\theta_0}$ and is finite. We denote by m this dimension and prove that $m = 1$ by computing the trace $\text{Tr}(B_\nu)$ of B_ν .

By Proposition 4.1 we have

$$\left\| B_\nu - \mathbb{1}_I^* \Pi_{q'(0)\nu}^H \mathbb{1}_I \right\|_{\mathcal{L}(L^2(I))} \xrightarrow[\nu \in \Sigma_{\theta_0}]{|\nu| \rightarrow +\infty} 0.$$

Let F_ν be a subspace of $L^2(I)$ of dimension $m + 1$ which contains $\text{Ran}(B_\nu)$ and $\mathbb{1}_I^* \varphi_{q'(0)\nu}^H$. We consider an orthonormal basis $(e_{0,\nu}, \dots, e_{m,\nu})$ of F_ν . Then

$$\begin{aligned} \left| \text{Tr} \left(B_\nu - \mathbb{1}_I^* \Pi_{q'(0)\nu}^H \mathbb{1}_I \right) \right| &\leq \sum_{j=0}^m \left| \left\langle \left(B_\nu - \mathbb{1}_I^* \Pi_{q'(0)\nu}^H \mathbb{1}_I \right) e_{j,\nu}, e_{j,\nu} \right\rangle \right| \\ &\leq (m + 1) \left\| B_\nu - \mathbb{1}_I^* \Pi_{q'(0)\nu}^H \mathbb{1}_I \right\|_{\mathcal{L}(L^2(I))} \xrightarrow[\nu \in \Sigma_{\theta_0}]{|\nu| \rightarrow +\infty} 0. \end{aligned}$$

Moreover, according to eq. (C.2), we have

$$\text{Tr} \left(\mathbb{1}_I^* \Pi_{q'(0)\nu}^H \mathbb{1}_I \right) \xrightarrow[\nu \in \Sigma_{\theta_0}]{|\nu| \rightarrow +\infty} 1.$$

Thus,

$$\text{Rank}(B_\nu) = \text{Tr}(B_\nu) \xrightarrow[\nu \in \Sigma_{\theta_0}]{|\nu| \rightarrow +\infty} 1,$$

and hence $m = 1$. This means that for $\nu \in \Sigma_{\theta_0}$ large enough the operator P_ν has a unique eigenvalue λ_ν in the disk $D(q'(0)\nu, q'(0)|\nu|)$, and this eigenvalue is algebraically simple.

It remains to prove the estimate on λ_ν . For this we reproduce the same argument with any $\varepsilon \in]0, 1]$. Then, given $\varepsilon \in]0, 1]$, there exists $\nu_\varepsilon \geq \nu_{\theta_0}$ such that the projection

$$B_{\nu,\varepsilon} = \frac{1}{2i\pi} \int_{|z - q'(0)\nu| = \varepsilon q'(0)|\nu|} (P_\nu - z)^{-1} dz$$

is well defined and has rank 1. This implies that λ_ν belongs to $D(q'(0)\nu, \varepsilon q'(0)|\nu|)$ when $|\nu| \geq \nu_\varepsilon$ and concludes the proof. \square

4.2. Agmon Estimates of Eigenfunctions

PROPOSITION 4.3. — *Let $\nu \in \mathbb{C}$ such that $\text{Re}(\nu) > 0$. Let λ be an eigenvalue of P_ν with associated eigenfunction φ . Set*

$$\mu := \frac{\text{Re} \left(e^{-i \arg(\nu)} \lambda \right)}{\cos(\arg(\nu))}.$$

Let $\kappa \in C^2(I; \mathbb{R})$ and $w := e^{\nu \kappa} \varphi$. The following Agmon equality holds:

$$\|w'\|_{L^2(I)}^2 + \int_I \left(|\nu|^2 \left(q(x)^2 - \kappa'^2(x) \right) - \mu \right) |w(x)|^2 dx = 0.$$

Proof. — Let $\nu = |\nu|e^{i\theta}$. The proof follows the usual Agmon’s equality strategy. We have $\varphi = e^{-\nu\kappa}w$. Thus, using Leibniz’ formula,

$$\begin{aligned}(P_\nu - \lambda)\varphi &= -\left(e^{-\nu\kappa}\right)''w - 2\left(e^{-\nu\kappa}\right)'w' - e^{-\nu\kappa}w'' + \left(\nu^2q^2 - \lambda\right)e^{-\nu\kappa}w \\ &= \left(-w'' + 2\nu\kappa'w' + \left(-\nu^2\kappa'^2 + \nu\kappa'' + \nu^2q^2 - \lambda\right)w\right)e^{-\nu\kappa}.\end{aligned}$$

Since $(P_\nu - \lambda_\nu)\varphi = 0$,

$$(4.12) \quad -w'' + 2\nu\kappa'w' + \left(-\nu^2\kappa'^2 + \nu\kappa'' + \nu^2q^2 - \lambda\right)w = 0.$$

Multiplying by $e^{-i\theta}\bar{w}$, integrating and taking the real part, we get

$$\begin{aligned}0 &= -\underbrace{\operatorname{Re} \int_I e^{-i\theta} w'' \bar{w}}_{I_1} \\ &\quad + \underbrace{|\nu| \operatorname{Re} \int_I \left(2\kappa' w' \bar{w} + \kappa'' |w|^2\right)}_{I_2} \\ &\quad + \underbrace{\int_I \left(\cos(\theta)|\nu|^2 (q^2 - \kappa'^2) - \operatorname{Re}(e^{-i\theta}\lambda)\right) |w|^2}_{I_3}.\end{aligned}$$

Integrating by parts in I_1 , we have $I_1 = \cos(\theta)\|w'\|_{L^2(I)}^2$. Considering I_2 , we get

$$I_2 = |\nu| \int_I \left(\kappa' |w|^2\right)' = 0.$$

Thus,

$$0 = I_1 + I_3 = \cos(\theta)\|w'\|_{L^2(I)}^2 + \int_I \left(\cos(\theta)|\nu|^2 (q^2 - \kappa'^2) - \operatorname{Re}(e^{-i\theta}\lambda)\right) |w|^2,$$

which is the claimed estimate multiplied by $\cos(\theta)$. \square

We will use Proposition 4.3 with $\kappa = (1 - \varepsilon) d_{\text{Agm}}$, where d_{Agm} defined in eq. (1.3). Up to this point, we assumed φ to be an eigenfunction of P_ν , but we did not specified which one, neither how it is normalized. We do this in the following definition, which is the natural extension of the definition of φ_n when $n \in \mathbb{N}$ (eq. (3.18)). For $\operatorname{Re}(\nu) > 0$ that satisfies the hypotheses of Proposition 4.2, let $\tilde{\varphi}_\nu \in L^2(I)$ be defined by

$$\tilde{\varphi}_\nu(x) := \nu^{1/4} e^{-q'(0)\nu x^2/2},$$

and

$$(4.13) \quad \varphi_\nu := \Pi_\nu(\tilde{\varphi}_\nu),$$

where Π_ν is the spectral projection for P_ν associated with λ_ν , as defined in Proposition 4.2.

PROPOSITION 4.4. — *Let φ_ν as in eq. (4.13). Let $\theta_0 \in \left[0, \frac{\pi}{2}\right)$. There exists $C > 0$ such that for every $\nu \in \Sigma_{\theta_0}$ with $|\nu| > \nu_0$,*

$$\begin{aligned}\|\tilde{\varphi}_\nu\|_{L^2(I)} &\leq C, \\ \|\varphi_\nu\|_{L^2(I)} &\leq C.\end{aligned}$$

Proof. —

Step 1: Estimation on $\tilde{\varphi}_\nu$. Since $\tilde{\varphi}_\nu$ is a restriction of $\nu^{1/4}e^{-\nu q'(0)x^2/2}$ on \mathbb{R} , we have

$$(4.14) \quad \|\tilde{\varphi}_\nu\|_{L^2(I)}^2 \leq |\nu|^{1/2} \int_{\mathbb{R}} e^{-2(\nu)q'(0)x^2/2} dx = \sqrt{\frac{\pi|\nu|}{q'(0)\operatorname{Re}(\nu)}} \leq \sqrt{\frac{\pi}{q'(0)\cos(\theta_0)}}.$$

Step 2: Estimation on φ_ν . Using the notations of Proposition 4.2, we have

$$(4.15) \quad \|\varphi_\nu\|_{L^2(I)} \leq \left(\left\| \mathbf{1}_I^* \Pi_{q'(0)\nu}^H \mathbf{1}_I \right\|_{\mathcal{L}(L^2(I))} + \left\| \Pi_\nu - \mathbf{1}_I^* \Pi_{q'(0)\nu}^H \mathbf{1}_I \right\|_{\mathcal{L}(L^2(I))} \right) \|\tilde{\varphi}_\nu\|_{L^2(I)}.$$

Moreover, according to Proposition 4.2, for $|\nu|$ large enough in Σ_{θ_0} ,

$$\left\| \Pi_\nu - \mathbf{1}_I^* \Pi_{q'(0)\nu}^H \mathbf{1}_I \right\|_{\mathcal{L}(L^2(I))} \leq 1.$$

Since the left-hand side is continuous in ν , we have that for $|\nu| > \nu_0$ and $\nu \in \Sigma_{\theta_0}$,

$$\left\| \Pi_\nu - \mathbf{1}_I^* \Pi_{q'(0)\nu}^H \mathbf{1}_I \right\|_{\mathcal{L}(L^2(I))} \leq C.$$

Finally, according to Proposition C.1, $\|\Pi_{q'(0)\nu}^H\|_{\mathcal{L}(L^2)}$ stays bounded for $\nu \in \Sigma_{\theta_0}$, thus

$$\left\| \mathbf{1}_I^* \Pi_{q'(0)\nu}^H \mathbf{1}_I \right\|_{\mathcal{L}(L^2(I))} + \left\| \Pi_\nu - \mathbf{1}_I^* \Pi_{q'(0)\nu}^H \mathbf{1}_I \right\|_{\mathcal{L}(L^2(I))} \leq C.$$

Plugging this fact into eq. (4.15) and combined with the fact that $\|\tilde{\varphi}_\nu\|_{L^2(I)}$ is bounded (eq. (4.14)), we get the claimed estimate. \square

COROLLARY 4.5. — *Let φ_ν as in eq. (4.13). Let $\theta_0 \in [0, \frac{\pi}{2})$ and $\varepsilon \in (0, 1)$. There exists $C > 0$ such that for every $\nu \in \Sigma_{\theta_0}$,*

$$\begin{aligned} \int_I \left| \varphi_\nu(x) e^{\nu(1-\varepsilon) d_{\text{Agm}}(x)} \right|^2 dx &\leq C|\nu|, \\ \left\| \varphi_\nu e^{\nu(1-\varepsilon) d_{\text{Agm}}(x)} \right\|_{L^\infty(I)} &\leq C|\nu|. \end{aligned}$$

Proof. — Let $w_\nu(x) := \varphi_\nu(x) e^{\nu(1-\varepsilon) d_{\text{Agm}}(x)}$. According to the Agmon equality of Proposition 4.3 with $\varphi = \varphi_\nu$ and $\lambda = \lambda_\nu$, and denoting the corresponding μ by μ_ν , we have

$$(4.16) \quad \|w'_\nu\|_{L^2(I)}^2 + \int_I \left(|\nu|^2 q(x)^2 (1 - (1 - \varepsilon)^2) - \mu_\nu \right) |w_\nu(x)|^2 dx = 0.$$

Step 1: First inequality. Let $K > 0$. We claim that if $x \in I$, $\nu \in \Sigma_{\theta_0}$ and $|\nu| d_{\text{Agm}}(x) > K$, then $|\nu|^2 q(x)^2 > cK|\nu|$ for some c depending on q , but not on $x \in I$, $\nu \in \Sigma_{\theta_0}$ nor K .

Indeed, routine computations show that in the limit $x \rightarrow 0$, $q(x)^2 \sim q'(0)^2 x^2$ and $d_{\text{Agm}}(x) \sim q'(0)x^2/2$. Thus, $q(x)^2/d_{\text{Agm}}(x) \rightarrow 2q'(0)$ as $x \rightarrow 0$. Moreover, we assumed that for every $x \neq 0$, $q(x) \neq 0$, thus for every $x \neq 0$, $q(x)^2/d_{\text{Agm}}(x) \neq 0$. This proves the claim.

Thus, if K is large enough, then for every $x \in I$ such that $d_{\text{Agm}}(x)|\nu| > K$

$$|\nu|^2 q(x)^2 (1 - (1 - \varepsilon)^2) - \mu_\nu \geq c|\nu|,$$

where we used the fact that $\mu_\nu \sim q'(0)|\nu|/\cos(\arg(\nu))$ because $\lambda_\nu = q'(0)\nu + o(\nu)$. Then, splitting the integral in the Agmon equality (4.16) into a part for $|\nu| d_{\text{Agm}}(x) < K$ and $|\nu| d_{\text{Agm}}(x) > K$, we get

$$\begin{aligned} c|\nu| \|w_\nu\|_{L^2(|\nu| d_{\text{Agm}}(x) > K)}^2 &\leq \int_{|\nu| d_{\text{Agm}}(x) < K} \left(\mu_\nu - |\nu|^2 q(x)^2 (1 - (1 - \varepsilon)^2) \right) |w_\nu(x)|^2 dx \\ &\leq C|\nu|^2 \|w_\nu\|_{L^2(|\nu| d_{\text{Agm}}(x) < K)}^2. \end{aligned}$$

We rewrite this as

$$\|w_\nu\|_{L^2(|\nu| d_{\text{Agm}}(x) > K)}^2 \leq C|\nu| \|w_\nu\|_{L^2(|\nu| d_{\text{Agm}}(x) < K)}^2.$$

Adding $\|w_\nu\|_{L^2(|\nu| d_{\text{Agm}}(x) < K)}^2$ on each side, this proves that

$$\|w_\nu\|_{L^2(I)}^2 \leq C|\nu| \|w_\nu\|_{L^2(|\nu| d_{\text{Agm}}(x) < K)}^2.$$

Using the definition of w_ν , we see that for $|\nu| d_{\text{Agm}}(x) < K$, $|w_\nu(x)| \leq e^K |\varphi_\nu(x)|$. Thus, using also the property $\|\varphi_\nu\|_{L^2(I)} \leq C$ (Proposition 4.4)

$$\|w_\nu\|_{L^2(I)}^2 \leq C|\nu| \|\varphi_\nu\|_{L^2(|\nu| d_{\text{Agm}}(x) < K)}^2 \leq C|\nu|.$$

Step 2: Second inequality. We again use Agmon's equality (4.16) to get

$$\|w'_\nu\|_{L^2(I)}^2 \leq \mu_\nu \|w_\nu\|_{L^2(I)}^2 \leq C|\nu| \|w_\nu\|_{L^2(I)}^2 \leq C|\nu|^2.$$

The claimed estimate then follows from Sobolev's embedding of $H^1(I)$ into $L^\infty(I)$. \square

We also prove the lower bound of Proposition 3.6 for φ_n when $n \geq 0$ is large enough.⁽¹⁰⁾

Proof of Proposition 3.6. — The part about λ_n was already proved in Proposition 4.2. By definition of φ_n , we have

$$\varphi_n = \Pi_n \tilde{\varphi}_n = \mathbb{1}_I^* \Pi_{q'(0)n}^H \mathbb{1}_I \tilde{\varphi}_n + (\Pi_n - \mathbb{1}_I^* \Pi_{q'(0)n}^H \mathbb{1}_I) \tilde{\varphi}_n.$$

According to Proposition 4.2, we have

$$\Pi_n - \mathbb{1}_I^* \Pi_{q'(0)n}^H \mathbb{1}_I \xrightarrow{n \rightarrow +\infty} 0.$$

Moreover, denoting by $\varphi_{\beta,1}^H(x) = (\text{Re}(\beta)/\pi)^{1/4} e^{-\beta x^2/2}$ the first eigenvector of H_β , we have for $\beta > 0$, $\Pi_\beta^H = \langle \varphi_{\beta,1}^H, \cdot \rangle \varphi_{\beta,1}^H$. Thus,

$$\langle \varphi_{q'(0)n,1}^H, \mathbb{1}_I^* \tilde{\varphi}_n \rangle = \sqrt{n} \left(\frac{q'(0)}{\pi} \right)^{1/4} \int_I e^{-nq'(0)x^2} dx.$$

The integral above is on I , but if we integrate on \mathbb{R} instead, we only add a small error term. Thus,

$$\langle \varphi_{q'(0)n,1}^H, \mathbb{1}_I^* \tilde{\varphi}_n \rangle = \left(\frac{\pi}{q'(0)} \right)^{1/4} + o_{n \rightarrow +\infty}(1).$$

⁽¹⁰⁾ This theorem actually holds if ν ranges over Σ_{θ_0} by using the expression of Π_β^H . We don't need this, so we refrain from doing so.

Hence,

$$\varphi_n = \left(\frac{\pi}{q'(0)} \right)^{1/4} \mathbb{1}_I^* \varphi_{q'(0)n,1}^H + o_{n \rightarrow +\infty} \left(\|\tilde{\varphi}_n\|_{L^2(I)} + 1 \right).$$

Since $\|\tilde{\varphi}_n\|_{L^2(I)}$ is bounded (Proposition 4.4) and since $\|\mathbb{1}_I^* \varphi_{q'(0)n,1}^H\|_{L^2(I)} = 1 + o(1)$ (thanks to similar computations as above), this proves the claimed lower bound. \square

4.3. Estimate for some pseudo-differential type operators on polynomials

In this section, we use the spectral analysis of the operator P_ν to deduce the operator estimate of Lemma 3.5. In order to do that, we need some definitions and theorems about a general class of operators on polynomials. The following comes from [Koe17, Definition 9 & Theorem 18].

DEFINITION 4.6. — *Let Ω be an open subset of \mathbb{C} . Assume that there exists $(r_\theta)_{0 \leq \theta < \pi/2}$ with $r_\theta \geq 0$ such that $\bigcup_{0 \leq \theta < \pi/2} \Sigma_\theta \setminus D(0, r_\theta) \subset \Omega$ (see Figure 4.1).*

We denote by $\mathcal{S}(\Omega)$ the set of functions γ holomorphic on Ω that have sub-exponential growth on each $\Sigma_\theta \cap \Omega$, i.e., for each $\theta \in [0, \pi/2)$ and $\delta > 0$, we have

$$p_{\theta,\delta}(\gamma) := \sup_{\nu \in \Sigma_\theta \cap \Omega} |\gamma(z) e^{-\delta|\nu|}| < +\infty.$$

We endow $\mathcal{S}(\Omega)$ with the topology defined by the seminorms $p_{\theta,\delta}$ for all $\theta \in [0, \pi/2)$ and $\delta > 0$.




Figure 4.1. An example of a set $\Sigma_\theta \setminus D(0, r_\theta) \subset \Omega$. The angle θ is allowed to be arbitrarily close to $\pi/2$, but then, the radius r_θ of the disk we avoid may blow up arbitrarily fast. For the Ω we will consider, the corresponding r_θ does blow up when $\theta \rightarrow \pi/2$ (at least, we cannot exclude that it blows up).

For the next theorem, if U is an open subset of \mathbb{C} , we denote the set of bounded holomorphic functions on U that have a zero of order n_0 at 0 by $\mathcal{O}_{n_0}^\infty(U)$. We endow $\mathcal{O}_{n_0}^\infty(U)$ with the L^∞ -norm.

THEOREM 4.7. — *Let $\Omega \subset \mathbb{C}$ as in Definition 4.6 and set $n_0 = \min\{n \in \mathbb{N} : [n, +\infty) \subset \Omega\}$. Let γ in $\mathcal{S}(\Omega)$ and $\gamma(z\partial_z)$ be the operator on polynomials with a zero of order n_0 at 0, defined by:*

$$\gamma(z\partial_z) \left(\sum_{n \geq n_0} a_n z^n \right) = \sum_{n \geq n_0} \gamma(n) a_n z^n.$$

Let U be a bounded open subset of \mathbb{C} . Let V be a neighborhood of \bar{U} that is star shaped with respect to 0. Then there exists $C > 0$ such that for all polynomials p with a zero of order n_0 at 0,

$$\|\gamma(z\partial_z)(p)\|_{L^\infty(U)} \leq C \|p\|_{L^\infty(V)}.$$

Moreover, the constant C above can be chosen continuously in $\gamma \in \mathcal{S}(\Omega)$: the map $\gamma \in \mathcal{S}(\Omega) \mapsto \gamma(z\partial_z)$ is continuous from $\mathcal{S}(\Omega)$ to $\mathcal{L}(\mathcal{O}_{n_0}^\infty(V), \mathcal{O}_{n_0}^\infty(U))$.

Remark 4.8. — While the operator $\gamma(z\partial_z)$ is initially defined only on polynomials with a zero of order n_0 at 0 (which is enough for our purposes), the continuity inequality of Theorem 4.7 allows us to extend $\gamma(z\partial_z)$ by continuity on the closure of the polynomials (with a zero of order n_0 at 0) for the $L^\infty(V)$ norm. If V is star-shaped with respect to 0, it is simply connected. If in addition it is bounded, Runge's theorem [Rud86, Theorem 13.9] implies that this closure is all bounded holomorphic functions on V with a zero of order n_0 at 0: $\gamma(z\partial_z)$ is a linear continuous operator on $\mathcal{O}_{n_0}^\infty(V)$.

We now have all the pieces needed to prove Lemma 3.5.

Proof of Lemma 3.5. — Let $\theta_0 \in [0, \frac{\pi}{2})$. According to Proposition 4.2 there exists r_{θ_0} such that if $|\nu| > r_{\theta_0}$ and $|\arg(\nu)| < \theta_0$, then there exists a unique eigenvalue λ_ν of P_ν close to $q'(0)\nu$. Moreover, this eigenvalue is algebraically and geometrically simple.

Set $\Omega = \bigcup_{\theta_0 \in [0, \pi/2)} \Sigma_{\theta_0} \setminus D(0, r_{\theta_0})$. Notice that by definition, Ω satisfies the property of Definition 4.6. For $0 < t < T$, $x \in I$ and $\nu \in \Omega$, we define

$$\tilde{\gamma}_{t,x}(\nu) := e^{-t(\lambda_\nu - q'(0)\nu)} \varphi_\nu(x) e^{\nu \operatorname{d}_{\text{Agm}}(x)(1-\varepsilon)},$$

and

$$(4.17) \quad \gamma_{t,x}(\nu) = \tilde{\gamma}_{t,x}(\nu + 1)$$

which is the natural extension of the definition of $\gamma_{t,x}(n)$ when $n \in \mathbb{N}$.

Step 1: The family $(\tilde{\gamma}_{t,x})_{0 < t < T, x \in I}$ is a bounded family⁽¹¹⁾ of $\mathcal{S}(\Omega)$. According to Proposition 4.2, λ_ν is algebraically simple on Ω . Thus, according to analytic perturbation theory (see, e.g., [Kat95, Chapter VII, § 1]), λ_ν and the associated spectral projection are holomorphic in $\nu \in \Omega$. Since $\varphi_\nu = \Pi_\nu(\nu^{1/4} e^{-\nu q'(0)x^2/2})$, φ_ν is holomorphic in $\nu \in \Omega$. We conclude that $\tilde{\gamma}_{t,x}(\nu)$ is holomorphic in $\nu \in \Omega$.

⁽¹¹⁾ Let us recall that if E is locally convex vector space whose topology is generated by a family $(p_\iota)_\iota$ of seminorms, a subset X of E is bounded if and only if for every ι , the set $\{p_\iota(x), x \in X\}$ is a bounded subset of \mathbb{R} .

We still have to prove that $(\tilde{\gamma}_{t,x})_{0 < t < T, x \in I}$ is a bounded family of $\mathcal{S}(\Omega)$. Let us set

$$\begin{aligned}\eta_t(\nu) &:= e^{-t(\lambda_\nu - q'(0)\nu)}, \\ \zeta_x(\nu) &:= \varphi_\nu(x) e^{\nu \operatorname{d}_{\operatorname{Agm}}(x)(1-\varepsilon)},\end{aligned}$$

and prove that both of the families $(\eta_t)_{0 < t < T}$ and $(\zeta_x)_{x \in I}$ are bounded in $\mathcal{S}(\Omega)$. It is easy to see that $(\gamma_1, \gamma_2) \in (\mathcal{S}(\Omega))^2 \mapsto \gamma_1 \gamma_2 \in \mathcal{S}(\Omega)$ is bounded, so this will prove the claim.

Let $\theta_0 \in [0, \frac{\pi}{2})$ and $\delta > 0$. According to Proposition 4.2, we have in the limit $|\nu| \rightarrow +\infty$, $\nu \in \Sigma_{\theta_0}$, $\lambda_\nu - \nu q'(0) = o(\nu)$. Thus, for ν large enough in Σ_{θ_0} ,

$$|\eta_t(\nu) e^{-\delta|\nu|}| = |e^{to(|\nu|) - \delta|\nu|}| < e^{TC_{\theta_0, \delta}}.$$

Thus, $(\eta_t)_{0 < t < T}$ is a bounded family of $\mathcal{S}(\Omega)$.

Similarly, according to Corollary 4.5, we have for any $x \in I$, and ν large enough in Σ_{θ_0} ,

$$|\zeta_x(\nu) e^{-\delta|\nu|}| \leq C.$$

This proves that $(\zeta_x)_{x \in I}$ is a bounded family of $\mathcal{S}(\Omega)$.

Step 2: The family $(\gamma_{t,x})_{0 < t < T, x \in I}$ is a bounded family of $\mathcal{S}(\Omega)$. According to the definition of Ω as a union of domains that look like the one of Figure 4.1, Ω is stable by $\nu \mapsto \nu + 1$. Then, the map $\gamma \in \mathcal{S} \mapsto \gamma(\cdot + 1) \in \mathcal{S}$ is well-defined and continuous. Thus, according to the first step and the definition of γ (eq. 4.17), the family $(\gamma_{t,x})_{0 < t < T, x \in I}$ is indeed a bounded family of $\mathcal{S}(\Omega)$.

Step 3: Conclusion. Let $n_0 = \min\{n \in \mathbb{N} : [n, +\infty) \subset \Omega\}$.⁽¹²⁾ Let U be a bounded open neighborhood of X such that $\bar{U} \subset V$. Then, the sets U and V satisfy the hypotheses of Theorem 4.7. Recall that linear continuous mappings between topological vector spaces maps every bounded subset to a bounded subset [Rud91, Theorem 1.32]. Hence, according to Theorem 4.7, the family of operators $(\gamma_{t,x}(z\partial_z))_{0 < t < T, x \in I}$ is bounded in $\mathcal{L}(\mathcal{O}_{n_0}^\infty(V), \mathcal{O}_{n_0}^\infty(U))$. Thus, there exists $C > 0$ such that for every polynomials p with a zero of order n_0 at 0, and for every $x \in I$ and $0 < t < T$,

$$\|\gamma_{t,x}(z\partial_z)(p)\|_{L^\infty(U)} \leq C\|p\|_{L^\infty(V)}.$$

□

⁽¹²⁾ In fact, we can be more precise in the construction of Ω and ensure that $\mathbb{R}_+ \subset \Omega$, in which case $n_0 = 0$. Indeed, the spectral theory of compact operators proves that for every $\nu \in \mathbb{C}$, the spectrum of P_ν is a discrete sequence of eigenvalues. The uniqueness of the solution of Cauchy problems for ODEs proves that when $\nu > 0$, these eigenvalues are actually geometrically and algebraically simple. Finally, perturbation theory proves that the first eigenvalue is holomorphic on the neighborhood of \mathbb{R}_+ . We do not need this, so we do not detail this.

5. Critical time of null-controllability for some domains

In this section, we prove Theorem 1.6.

Proof of Theorem 1.6. — Set $a_- = -\max(\gamma_2^-)$ and $a_+ = \max(\gamma_1^+)$. Denote by $y_- \in \mathbb{T}$ and $y_+ \in \mathbb{T}$ points where these maxima are reached.

Step 1: Lower bound of the minimal time. For this step, we only have to treat the case $T_* > 0$. In this case, either $a_- < 0$ or $a_+ > 0$. If $a_- < 0$, for any $a_- < a < 0$, the segment $[a, L_+] \times \{y_-\}$ stays at positive distance of ω , and thanks to Theorem 3.1, the generalized Baouendi–Grushin equation (3.2) is not null-controllable on ω in time $T < d_{\text{Agm}}(a)/q'(0)$. Similarly, if $a_+ > 0$, for any $0 < a < a_+$, the segment $[-L_-, a] \times \{y_+\}$ stays at positive distance from ω , and the generalized Baouendi–Grushin equation is not null-controllable in time $T < d_{\text{Agm}}(a)/q'(0)$.

This holds for any $a_- < a < 0$ and $0 < a < a_+$, thus the generalized Baouendi–Grushin equation (3.2) is not null-controllable in time $T < T_*$.

Step 2: Upper bound of the minimal time. Let $\varepsilon > 0$ small enough so that $\gamma_2 - \gamma_1 > \varepsilon$. Let $\tilde{\gamma}_1 = \max(\gamma_1, a_- - \varepsilon)$ and $\tilde{\gamma}_2 = \min(\gamma_2, a_+ + \varepsilon)$. By using the information $\gamma_2 - \gamma_1 > \varepsilon$, $a_- \leq \gamma_2$, $\gamma_1 \leq a_+$ and by looking at the different cases, we readily get $\tilde{\gamma}_2 - \tilde{\gamma}_1 \geq \varepsilon$. Then, we define the path

$$\gamma = (\gamma_x, \gamma_y): s \in \mathbb{T} \mapsto \left(\frac{\tilde{\gamma}_1(s) + \tilde{\gamma}_2(s)}{2}, s \right).$$

This path satisfies $\gamma_1 + \varepsilon/2 \leq \gamma_x \leq \gamma_2 - \varepsilon/2$, hence $\gamma(\mathbb{T}) \subset \omega$. Moreover, we see that it satisfies the hypotheses of Theorem 1.4, because the connected components of $(I \times \mathbb{T}) \setminus \gamma(\mathbb{T})$ are $\{(x, y): x < \gamma_x(y)\}$ and $\{(x, y): x > \gamma_x(y)\}$. Moreover, this path satisfies

$$a_- - \frac{\varepsilon}{2} \leq \gamma_x \leq a_+ + \frac{\varepsilon}{2}.$$

Thus, Theorem 1.4 implies that the generalized Baouendi–Grushin equation (3.2) is null-controllable in time $T > \max(d_{\text{Agm}}(a_+ + \varepsilon/2), d_{\text{Agm}}(a_- - \varepsilon/2))/q'(0)$. As this holds for every $\varepsilon > 0$ small enough, the result follows. \square

Appendix A. Baouendi–Grushin equation on $I \times (0, \pi)$

In the article, we stated and proved results on the Baouendi–Grushin posed on $I \times \mathbb{T}$. These results have a version for the Baouendi–Grushin equation posed on $I \times (0, \pi)$:

$$(A.1) \quad \begin{cases} (\partial_t - \partial_x^2 - q(x)^2 \partial_y^2) f(t, x, y) = \mathbb{1}_\omega u(t, x, y), & t \in (0, T), x \in I, y \in (0, \pi) \\ f(t, x, y) = 0, & t \in (0, T), (x, y) \in \partial(I \times (0, \pi)), \\ f(0, x, y) = f_0, & x \in I, y \in (0, \pi). \end{cases}$$

Here, we precisely state them and explain what are the differences, if any, in their proofs. The precise definition of the operator, especially its domain, is again the




Figure 5.1. In green, the domain ω . At $y = y_-$, the function γ_2^- takes its maximum a_- . Then, the interval $(a_-, L_+) \times \{y_-\}$ is disjoint from $\bar{\omega}$. So, the Grushin equation is not null-controllable in time $T < d_{\text{Agm}}(a_-)/q'(0)$. Similarly, the interval $(-L_-, a_+) \times \{y_+\}$ is disjoint from $\bar{\omega}$. So, the Grushin equation is not null-controllable in time $T < d_{\text{Agm}}(a_+)/q'(0)$. Also, if we take a path γ (here in blue) that is close to the boundary of ω around $y = y_-$ and $y = y_+$, then, we can apply Theorem 1.4, and the Grushin equation is null-controllable in time $T > \max(d_{\text{Agm}}(a_-), d_{\text{Agm}}(a_+))/q'(0)$.

Friedrichs' extension. That it generates an analytic semigroup is again proved with Hille–Yosida's theorem. We again refer to [BCG14, Bre11, Hel13] for the details.

The adaptation of the positive controllability result is:

THEOREM A.1. — Assume that $q \in C^3(\bar{I})$ is such that $q(0) = 0$ and $\min_{\bar{I}} q' > 0$. Let ω be an open subset of $I \times [0, \pi]$. Assume that there exists $\gamma = (\gamma_x, \gamma_y) \in C^0([0, 1], I \times [0, \pi])$ such that $\gamma((0, 1)) \subset \omega$, $\gamma_y(0) = 0$ and $\gamma_y(1) = \pi$.

The generalized Baouendi–Grushin equation (A.1) is null-controllable on ω in time T such that

$$T > \frac{1}{q'(0)} \max \left(d_{\text{Agm}} \left(\min_{\mathbb{T}}(\gamma_x) \right), d_{\text{Agm}} \left(\max_{\mathbb{T}}(\gamma_x) \right) \right).$$

The proof is mostly the same, the only small difference being the construction of the cutoff function, which is done thanks to Proposition B.5 and the natural adaptation of Proposition B.1.

The adaptation of the negative result Theorem 1.3 is straightforward. We use the same notation δ as in Theorem 1.3:

THEOREM A.2. — Assume that $q \in C^2(\bar{I})$ is such that $q(0) = 0$, $q'(0) > 0$ and $q(x) \neq 0$ whenever $x \neq 0$. Let ω be an open subset of $I \times (0, \pi)$. Assume that there exist $a \in [-L_-, 0)$, $b \in (0, L_+]$ and $y_0 \in (0, \pi)$ such that

$$\text{distance} \left((a, b) \times \{y_0\}, \omega \right) > 0.$$

Then, the generalized Baouendi–Grushin equation (A.1) is not null-controllable on ω in time T such that

$$T < \frac{1}{q'(0)} \min(\delta(a), \delta(b)).$$

We prove this theorem with the following lemma:

LEMMA A.3. — Let $T > 0$ and let ω be an open subset of $I \times (0, \pi)$. Denote by $S(\omega)$ the symmetric of ω with respect to $\{y = 0\}$. Let λ_n and φ_n as in eq. (3.18).

Assume that the generalized Baouendi–Grushin equation with Dirichlet boundary conditions (A.1) is null controllable on ω in time T , then for every complex sequence (a_n) with a finite number of nonzero terms,

$$\sum_{n \geq 0} |a_n|^2 \|\varphi_n\|_{L^2(I)}^2 e^{-2\lambda_n T} \leq C \int_{[0, T] \times (\omega \cup S(\omega))} \left| \sum_{n \geq 0} a_n \varphi_n(x) e^{iny - \lambda_n t} \right|^2 dt dx dy.$$

Sketch of the proof. — This lemma is proved by testing the associated observability inequality on the function $g(t, x, y) = \sum_{n \geq 0} a_n \varphi_n(x) \sin(ny) e^{-\lambda_n t}$, and writing $\sin(ny) = (e^{iny} - e^{-iny})/(2i)$. Thus, with $\tilde{g}(t, x, y) = \sum_{n \geq 0} a_n \varphi_n(x) e^{iny - \lambda_n t}$, $g(t, x, y) = (\tilde{g}(t, x, y) - \tilde{g}(t, x, -y))/(2i)$, the right-hand side of the observability inequality satisfies

$$\|g\|_{L^2([0, T] \times \omega)}^2 \leq \frac{1}{2} \left(\|\tilde{g}\|_{L^2([0, T] \times \omega)}^2 + \|\tilde{g}\|_{L^2([0, T] \times S(\omega))}^2 \right).$$

The right-hand side of this inequality is the right-hand side of the claimed estimate. \square

Theorem A.2 is then proved by remarking that we already disproved such an inequality in Section 3.3.

Appendix B. Existence of the cutoff function and homotopy

We begin with the construction of the cutoff function used in the proof of Theorem 1.4.

PROPOSITION B.1. — Let $a < b$ and let ω be an open subset of $(a, b) \times \mathbb{T}$. Assume that there exists a closed path $\gamma = (\gamma_x, \gamma_y) \in C^0(\mathbb{T}; \omega)$ such that $\{a\} \times \mathbb{T}$ and $\{b\} \times \mathbb{T}$ are included in different connected components of $([a, b] \times \mathbb{T}) \setminus \gamma(\mathbb{T})$. Let $\omega_- = [a, \min \gamma_x] \times \mathbb{T}$ and $\omega_+ = [\max \gamma_x, b] \times \mathbb{T}$. There exists a function $\chi \in C^\infty([a, b] \times \mathbb{T})$ such that:

- $\chi = 0$ on $\omega_+ \setminus \omega$;
- $\chi = 1$ on $\omega_- \setminus \omega$;
- $\text{supp}(\nabla \chi) \subset \omega$.

Proof of Proposition B.1. —

Step 1: Defining χ . Let $\varepsilon > 0$ small enough so that $\text{distance}(\gamma(\mathbb{T}), \{a, b\} \times \mathbb{T}) > \varepsilon$ and such that for any $z \in \gamma(\mathbb{T})$, $B(z, \varepsilon) \subset \omega$. We set

$$\omega_0 := \{z \in [a, b] \times \mathbb{T} : \text{distance}(z, \gamma(\mathbb{T})) < \varepsilon\}.$$

Let $\rho \in C_c^\infty(B(0, \varepsilon/2))$ with $\int_{B(0, \varepsilon/2)} \rho(z) dz = 1$. Consider Ω the connected component of $\{a\} \times \mathbb{T}$ in $(\mathbb{R} \times \mathbb{T}) \setminus \gamma(\mathbb{T})$. Set $\chi = \rho * \mathbb{1}_\Omega$ (initially defined on $\mathbb{R} \times \mathbb{T}$ and then restricted on $[a, b] \times \mathbb{T}$).

Step 2: $\text{supp}(\nabla \chi) \subset \omega_0$. According to the definition of Ω , $\mathbb{1}_\Omega$ is locally constant outside of $\gamma(\mathbb{T})$. This implies that χ is locally constant around each z such that

$\text{distance}(z, \gamma(\mathbb{T})) > \varepsilon/2$. Hence $\text{supp}(\nabla\chi) \subset \{z: \text{distance}(z, \gamma(\mathbb{T})) \leq \varepsilon/2\}$. According to our choice of ε , this proves the claim that $\text{supp}(\nabla\chi) \subset \omega_0 \subset \omega$.

Step 3: Value of χ on $\omega_- \setminus \omega$. According to the definition of χ and the fact that $\text{distance}(\{a\} \times \mathbb{T}, \gamma(\mathbb{T})) > \varepsilon$, for any $y_0 \in \mathbb{T}$, $\chi(a, y_0) = 1$. Moreover, $\omega_- \setminus \omega_0$ is connected (according to the definition of ω_- , we can connect every $(x, y) \in \omega_- \setminus \omega_0$ to the left boundary $\{a\} \times \mathbb{T}$ with the horizontal segment $[a, x] \times \{y\}$). According to the previous step, χ is locally constant outside ω_0 . Hence, χ is constant in $\omega_- \setminus \omega_0$.

Step 4: Value of χ on $\omega_+ \setminus \omega$. According to the definition of χ and the hypothesis, $\chi = 0$ on $\{b\} \times \mathbb{T}$. The rest of this step is a copy-paste of the previous step. \square

Now, we justify Remark 1.5, with the following two propositions:

PROPOSITION B.2. — *Let $a < b$ and let $\gamma \in C^0(\mathbb{T}, (a, b) \times \mathbb{T})$ be a closed path that is not homotopic to a constant path. Then $\{a\} \times \mathbb{T}$ and $\{b\} \times \mathbb{T}$ are included in different connected components of $([a, b] \times \mathbb{T}) \setminus \gamma(\mathbb{T})$.*

Proof. — Assume that for some $y_0, y_1 \in \mathbb{T}$, there exists a continuous path c_1 in $(\bar{I} \times \mathbb{T}) \setminus \gamma(\mathbb{T})$ from (a, y_0) to (b, y_1) . Since $\bar{I} \times \mathbb{T}$ is Hausdorff, we may assume that c_1 is simple. We can also assume that c_1 touches the boundaries $\{a, b\} \times \mathbb{T}$ only at the start and end.

Now, consider the universal cover $[a, b] \times \mathbb{R}$ of $[a, b] \times \mathbb{T}$. Consider \tilde{c}_1 a lift of c_1 to $[a, b] \times \mathbb{R}$, that starts at (a, \tilde{y}_0) and ends at (b, \tilde{y}_1) . Let c_2 the simple closed loop formed by concatenating c_1 , the vertical segment $\{b\} \times [\tilde{y}_1, \tilde{y}_1 + 2\pi]$, the reverse of the path $\tilde{c}_1 + (0, 2\pi)$, and finally the vertical segment $\{a\} \times [\tilde{y}_0, \tilde{y}_0 + 2\pi]$ from top to bottom (see Figure B.1).




Figure B.1. Illustration of the path c_2 defined in the proof of Proposition B.2.

If we see this path c_2 as a path on \mathbb{R}^2 , according to Jordan's theorem, $\mathbb{R}^2 \setminus c_2$ has two path-connected components, one of them bounded. Let us denote by Ω_1 this bounded component, which, according to Jordan–Schoenflies' theorem, is simply connected. One of the lift of γ lies in Ω_1 , let us call it $\tilde{\gamma}$. But γ is not homotopic to a constant path, which contradicts the simple connectedness of Ω_1 . \square

PROPOSITION B.3. — *Let ω be a connected open subset of $[a, b] \times \mathbb{T}$ such that $\{a\} \times \mathbb{T}$ and $\{b\} \times \mathbb{T}$ are included in different connected components of $([a, b] \times \mathbb{T}) \setminus \omega$. Let $\tilde{\omega}$ a connected open subset of $[a, b] \times \mathbb{T}$ such that $\bar{\omega} \subset \tilde{\omega}$.*

There exists a closed path $\gamma \in C^0(\mathbb{T}, \tilde{\omega})$ that is not homotopic in $[a, b] \times \mathbb{T}$ to a constant path.

Remark B.4. — Let γ be a closed path in $(a, b) \times \mathbb{T}$ such that $\{a\} \times \mathbb{T}$ and $\{b\} \times \mathbb{T}$ are included in different connected components of $([a, b] \times \mathbb{T}) \setminus \gamma(\mathbb{T})$. It is possible this path γ is homotopic to a constant path, but Proposition B.3 applied with $\omega := \{z: \text{distance}(z, \gamma(\mathbb{T})) < \varepsilon\}$ and $\tilde{\omega} := \{z: \text{distance}(z, \gamma(\mathbb{T})) < 2\varepsilon\}$ tells us that for any $\varepsilon > 0$ there exists a path $\tilde{\gamma}$ that stays at distance at most 2ε from $\gamma(\mathbb{T})$ and that is not homotopic to a constant path.

Proof. — The proof uses some basic tools of algebraic topology, in particular van Kampen’s theorem (see for instance Hatcher’s “Algebraic Topology” [Hat02, § 0.1, § 1.1–1.2]).

Let C_- be the connected component of $\{a\} \times \mathbb{T}$ in $([a, b] \times \mathbb{T}) \setminus \omega$. Set

$$(B.1) \quad A_- := C_- \cup \tilde{\omega}, \quad A_+ := (([a, b] \times \mathbb{T}) \setminus C_-) \cup \tilde{\omega}.$$

Step 1: Every connected component C of $([a, b] \times \mathbb{T}) \setminus \omega$ is closed and satisfies $\partial C \subset \tilde{\omega}$. Here, ∂C is the boundary of C as a subset of $[a, b] \times \mathbb{T}$.

Indeed, connected components of a topological space X are closed in X , hence C is closed in $([a, b] \times \mathbb{T}) \setminus \omega$, which is itself closed in $[a, b] \times \mathbb{T}$.

If $x \in \partial C$ is such that $x \notin \partial \omega$, then there exists $\varepsilon > 0$ such that $B(x, \varepsilon) \subset ([a, b] \times \mathbb{T}) \setminus \omega$. But then, $B(x, \varepsilon)$ is included in the connected component of x , i.e., C . By contradiction, we see that every $x \in \partial C$ is in $\partial \omega \subset \tilde{\omega}$.

Step 2: A_+ and A_- are open and connected. We begin with the openness of A_- . If $x \in A_-$, there are three cases:

- If x is in the interior of C_- , it is in the interior of A_- by definition.
- If x is in $\tilde{\omega}$, since $\tilde{\omega}$ is open, x is also in the interior of A_- .
- If $x \in \partial C_-$, according to Step 1, $x \in \tilde{\omega}$ which implies that x is in the interior of A_- .

The subset A_+ is open because it is the union of the open subsets $([a, b] \times \mathbb{T}) \setminus C_-$ and $\tilde{\omega}$.

Since A_- is the union of two connected subset that have a non-empty intersection (we saw in Step 1 that $\partial C_- \subset C_- \cap \tilde{\omega}$), A_- is connected. Finally, A_+ is connected because it is the union of the connected subset $\tilde{\omega}$ and of the connected components of $[a, b] \times \mathbb{T} \setminus \omega$ other than C_- , which all have a non-empty intersection with $\tilde{\omega}$.

Step 3: Conclusion using van Kampen’s theorem. If α is a closed path in a topological space X , we will denote its homotopy class by $[\alpha]_X$. We will denote the fundamental group of X by $\pi_1(X)$. We will denote by p_- (respectively p_+) the canonical injection of $\pi_1(A_-)$ (respectively $\pi_1(A_+)$) into the free product $\pi_1(A_-) * \pi_1(A_+)$.

According to van Kampen’s theorem [Hat02, Theorem 1.20], the map $k: \pi_1(A_-) * \pi_1(A_+) \rightarrow \pi_1([a, b] \times \mathbb{T})$ defined by

$$[\alpha]_{A_{\pm}} \in \pi_1(A_{\pm}) \mapsto [\alpha]_{[a, b] \times \mathbb{T}} \in \pi_1([a, b] \times \mathbb{T})$$

is surjective. Moreover, its kernel is generated by $p_+([\alpha]_{A_+})p_-([\alpha]_{A_-})^{-1}$ for all closed paths α in $A_- \cap A_+ = \tilde{\omega}$.

If we denote the closed path $s \in \mathbb{T} \mapsto (a, s) \in [a, b] \times \mathbb{T}$ (respectively $s \mapsto (b, s)$) by β_- (respectively β_+), the definition of k implies that

$$k\left(p_-([\beta_-]_{A_-})\right) = [\beta_-]_{[a,b] \times \mathbb{T}} = [\beta_+]_{[a,b] \times \mathbb{T}} = k\left(p_+([\beta_+]_{A_+})\right).$$

Thus, $\xi := p_-([\beta_-]_{A_-})p_+([\beta_+]_{A_+})^{-1} \in \ker(k)$. According to the previous discussion, ξ is a product of terms of the form $p_{\pm}([\alpha_k]_{A_{\pm}})p_{\mp}([\alpha_k]_{A_{\mp}})^{-1}$ for a finite number of paths α_k in $\tilde{\omega}$. Reducing the word ξ in the free product $\pi_1(A_-) * \pi_1(A_+)$, ξ can be written in the form

$$(B.2) \quad p_-([\beta_-]_{A_-})p_+([\beta_+]_{A_+})^{-1} = \xi = p_{\pm}([\tilde{\alpha}_1]_{A_{\pm}})p_{\mp}([\tilde{\alpha}_2]_{A_{\mp}})p_{\pm}([\tilde{\alpha}_3]_{A_{\pm}}) \cdots$$

where none of the terms in the right-hand side are the neutral element of $\pi_1(A_{\pm})$. Since the left-hand side is already a reduced word, by definition of the free product of groups, the two words on the left and right-hand side of this equality are the same. Thus there are exactly two factors in the right-hand side of eq. (B.2) and

$$[\beta_-]_{A_-} = [\tilde{\alpha}_1]_{A_-}, \quad [\beta_+]_{A_+}^{-1} = [\tilde{\alpha}_2]_{A_+}.$$

The first equality tells us that $\tilde{\alpha}_1$ is homotopic in A_- to β_- . Since $\tilde{\alpha}_1$ is in $\tilde{\omega}$ and since β_- is not homotopic to a constant path in $[a, b] \times \mathbb{T}$, this proves the proposition. \square

To end this appendix, we mention that Proposition B.2 has a variant when the domain is $(a, b) \times [0, \pi]$ instead of $(a, b) \times \mathbb{T}$:

PROPOSITION B.5. — *Let $a < b$ and let $\gamma = (\gamma_x, \gamma_y) \in C^0([0, 1], (a, b) \times [0, \pi])$ be a closed path such that $\gamma_y(0) = 0$ and $\gamma_y(1) = \pi$. Then $\{a\} \times [0, \pi]$ and $\{b\} \times [0, \pi]$ are included in different connected components of $[a, b] \times [0, \pi] \setminus (\gamma([0, 1]))$.*

The proof also uses Jordan's theorem, but in a simpler way than Proposition B.2, and is left to the reader.

Appendix C. Non-selfadjoint harmonic oscillators

Let $\beta \in \mathbb{C}$ with $\operatorname{Re}(\beta) > 0$. We discuss in this appendix the basic properties of the non-selfadjoint harmonic oscillator (or Davies operator) defined on $L^2(\mathbb{R})$ by

$$(C.1) \quad H_{\beta} = -\partial_x^2 + \beta^2 x^2.$$

More precisely, we set

$$\operatorname{Dom}(H_{\beta}) = \left\{ u \in L^2(\mathbb{R}) : -\partial_x^2 u + \beta^2 x^2 u \in L^2(\mathbb{R}) \right\},$$

where $-\partial_x^2 u + \beta^2 x^2 u$ is understood in the sense of distributions, and we define H_{β} by eq. (C.1) on $\operatorname{Dom}(H_{\beta})$. This defines an unbounded operator on $L^2(\mathbb{R})$. When $\beta = 1$ we recover the usual harmonic oscillator.

The spectral properties of the Davies operator has been studied (see, among others, [Dav07, § 14.5], [Hel13, § 14.4], [KSTV15], and the references therein), and the properties stated in this appendix are standard, at least in spirit. Nevertheless, for the reader convenience, we collect and prove the properties needed in our study.

PROPOSITION C.1. — Let $\beta \in \mathbb{C}$ with $\operatorname{Re}(\beta) > 0$.

- (i) The operator H_β is closed and has compact resolvent.
- (ii) The adjoint of H_β is $H_\beta^* = H_{\bar{\beta}}$.
- (iii) The spectrum of H_β is given by the sequence of (geometrically and algebraically) simple eigenvalues $(\lambda_{\beta,k}^H)_{k \in \mathbb{N}^*}$ where

$$\lambda_{\beta,k}^H = (2k - 1)\beta.$$

An eigenfunction associated to $\lambda_{\beta,k}^H$ is given by

$$\varphi_{\beta,k}^H(x) = h_{k-1}(\sqrt{\beta}x) e^{-\beta x^2/2}, \quad \text{where } h_{k-1}(x) = (-1)^{k-1} e^{x^2} \partial_x^{k-1} (e^{-x^2})$$

is the $(k - 1)^{\text{th}}$ Hermite polynomial.

- (iv) For $\gamma > 0$ and $\varepsilon > 0$ we set

$$(C.2) \quad \mathcal{Z}_{\beta,\varepsilon,\gamma} := \{z \in \mathbb{C} : |z| \leq \gamma |\beta|, \text{ distance}(z, \sigma(H_\beta)) \geq \varepsilon |\beta|\}.$$

Let $\theta_0 \in [0, \frac{\pi}{2})$. There exists $C > 0$ such that if $\beta \in \Sigma_{\theta_0}$ (see eq. (4.1)) then for $z \in \mathcal{Z}_{\beta,\varepsilon,\gamma}$ we have

$$(C.3) \quad \|(H_\beta - z)^{-1}\|_{\mathcal{L}(L^2(\mathbb{R}))} \leq \frac{C}{|\beta|}.$$

- (v) We denote by Π_β^H the spectral projection of H_β associated to the eigenvalue $\lambda_{\beta,1}^H = \beta$. Then we have (note the $\bar{\beta}$ instead of β in the first argument of the inner products)

$$(C.4) \quad \Pi_\beta^H u = \frac{\langle \varphi_{\bar{\beta},1}^H, u \rangle}{\langle \varphi_{\bar{\beta},1}^H, \varphi_{\beta,1}^H \rangle} \varphi_{\beta,1}^H.$$

and

$$(C.5) \quad \|\Pi_\beta^H\|_{\mathcal{L}(L^2(\mathbb{R}))} = \sqrt{\frac{|\beta|}{\operatorname{Re}(\beta)}} = \frac{1}{\sqrt{\cos(\arg(\beta))}}.$$

Proof. —

Step 1: H_β has compact resolvent. The closedness of H_β is clear. Let $\theta = \arg(\beta) \in (-\frac{\pi}{2}, \frac{\pi}{2})$. For $u \in \operatorname{Dom}(H_\beta)$ we have

$$(C.6) \quad \langle e^{-i\theta} H_\beta u, u \rangle_{L^2(\mathbb{R})} = e^{-i\theta} \|u'\|_{L^2(\mathbb{R})}^2 + |\beta|^2 e^{i\theta} \|xu\|_{L^2(\mathbb{R})}^2,$$

so the numerical range of $e^{-i\theta} H_\beta$ is included in Σ_θ . In particular, $(e^{-i\theta} H_\beta + 1)$ is injective. Now let

$$B^1(\mathbb{R}) = \{u \in H^1(\mathbb{R}) : xu \in L^2(\mathbb{R})\}.$$

This is a Hilbert space for the natural norm

$$\|u\|_{B^1(\mathbb{R})}^2 = \|u\|_{H^1(\mathbb{R})}^2 + \|xu\|_{L^2(\mathbb{R})}^2.$$

For $u, v \in B^1(\mathbb{R})$ we set

$$Q_\beta(u, v) = e^{-i\theta} \langle u', v' \rangle_{L^2(\mathbb{R})} + |\beta|^2 e^{i\theta} \langle xu, xv \rangle_{L^2(\mathbb{R})} + \langle u, v \rangle_{L^2(\mathbb{R})}.$$

Let $f \in L^2(\mathbb{R})$. By the Lax–Milgram Theorem, there exists a unique $u \in B^1(\mathbb{R})$ such that $Q_\beta(u, v) = \langle f, v \rangle_{L^2(\mathbb{R})}$ for all $v \in B^1(\mathbb{R})$. In the sense of distributions we have $e^{-i\theta}(-u'' + \beta^2 x^2 u) + u = f \in L^2(\mathbb{R})$, so $u \in \text{Dom}(H_\beta)$ and $(e^{-i\theta}H_\beta + 1)u = f$. This proves that $-e^{i\theta}$ belongs to the resolvent set of H_β .

Finally, taking the real part of eq. (C.6) gives for $u \in \text{Dom}(H_\beta)$

$$\|u'\|_{L^2(\mathbb{R})}^2 + |\beta|^2 \|xu\|_{L^2(\mathbb{R})}^2 \leq \frac{\text{Re} \langle e^{-i\theta} H_\beta u, u \rangle}{\cos(\theta)} \leq \frac{1}{2 \cos(\theta)} (\|H_\beta u\|_{L^2(\mathbb{R})}^2 + \|u\|_{L^2(\mathbb{R})}^2).$$

We deduce that $\text{Dom}(H_\beta)$ is compactly embedded in $L^2(\mathbb{R})$. Since H_β has nonempty resolvent set, it has compact resolvent. In particular, its spectrum consists of isolated eigenvalues of finite multiplicities.

Step 2: Computation of $(H_\beta)^$.* Let $v \in \text{Dom}(H_{\bar{\beta}})$. For $u \in \text{Dom}(H_\beta)$ we have $\langle H_\beta u, v \rangle = \langle u, H_{\bar{\beta}} v \rangle$, so $\text{Dom}(H_{\bar{\beta}}) \subset \text{Dom}(H_\beta^*)$ and $H_\beta^* = H_{\bar{\beta}}$ on $\text{Dom}(H_{\bar{\beta}})$. Now let $v \in \text{Dom}(H_\beta^*)$ and $f = H_\beta^* v$. Then $v \in L^2(\mathbb{R})$ and in the sense of distributions we have $-v'' + \bar{\beta}^2 x^2 v = f \in L^2(\mathbb{R})$, so $v \in \text{Dom}(H_{\bar{\beta}})$. This proves that $\text{Dom}(H_\beta^*) \subset \text{Dom}(H_{\bar{\beta}})$, and hence $H_\beta^* = H_{\bar{\beta}}$.

Step 3: Eigenvalues and eigenfunctions of H_β , completeness of the eigenfunctions. For $k \in \mathbb{N}^*$ we have $\varphi_{\beta,k}^H \in \text{Dom}(H_\beta)$ and it is classical computation that for $\beta = 1$, (see, e.g., [Hel13, §1.3])

$$H_1 \varphi_{1,k}^H = \lambda_{1,k}^H \varphi_{1,k}^H.$$

Noticing that $\varphi_{\beta,k}^H(x) = \varphi_{1,k}(\sqrt{\beta}x)$, routine computations using the scaling $x' = \sqrt{\beta}x$ show that

$$H_\beta \varphi_{\beta,k}^H = \lambda_{\beta,k}^H \varphi_{\beta,k}^H.$$

Then $\lambda_{\beta,k}^H$ is an eigenvalue of H_β and $\varphi_{\beta,k}^H$ is a corresponding eigenfunction.

Let $u \in \text{span}(\varphi_{\beta,k}^H)_{k \in \mathbb{N}^*}^\perp$. For all polynomial p we have $\int_{\mathbb{R}} u(x) p(\sqrt{\beta}x) e^{-\beta x^2/2} dx = 0$. For $\xi \in \mathbb{R}$ we set $F(\xi) = \int_{\mathbb{R}} e^{i\xi x} u(x) e^{-\beta x^2/2} dx$. Then F is analytic and $F^{(n)}(0) = 0$ for all $n \in \mathbb{N}$. This implies that $u(x) = 0$ for almost all $x \in \mathbb{R}$, so the family $(\varphi_{\beta,k}^H)_{k \in \mathbb{N}^*}$ is complete.

Step 4: Resolvent estimate. The map

$$(\theta, \zeta) \mapsto (H_{e^{i\theta}} - e^{i\theta} \zeta)^{-1}$$

is continuous and hence bounded on the compact $[-\theta_0, \theta_0] \times \mathcal{Z}_{1,\varepsilon,\gamma}$, so eq. (C.3) holds if $|\beta| = 1$.

For $\rho > 0$ we consider on $L^2(\mathbb{R})$ the unitary operator Θ_ρ such that for $u \in L^2(\mathbb{R})$ and $x \in \mathbb{R}$ we have

$$(\Theta_\rho u)(x) = \rho^{\frac{1}{2}} u(\rho x).$$

We observe that

$$\Theta_{|\beta|^{\frac{1}{2}}}^{-1} H_\beta \Theta_{|\beta|^{\frac{1}{2}}} = |\beta| H_{\frac{\beta}{|\beta|}},$$

so for $z \in \mathcal{Z}_{\beta,\varepsilon,\gamma}$ we have

$$\|(H_\beta - z)^{-1}\|_{\mathcal{L}(L^2(\mathbb{R}))} = \frac{1}{|\beta|} \left\| \left(H_{\frac{\beta}{|\beta|}} - \frac{z}{|\beta|} \right)^{-1} \right\|_{\mathcal{L}(L^2(\mathbb{R}))}.$$

Since $|\beta|^{-1} \mathcal{Z}_{\beta, \varepsilon, \gamma} = \mathcal{Z}_{\beta/|\beta|, \varepsilon, \gamma}$, we deduce eq. (C.3) in the general case.

Step 5: Spectral projection. In the integral $\langle \varphi_{\beta, k}^H, \varphi_{\beta, 1}^H \rangle = \int_{\mathbb{R}} h_{k-1}(\sqrt{\beta}x) e^{-\beta x^2} dx$, we make the change of variables and integration path $x' = \sqrt{\beta}x$, which can be justified thanks to the gaussian decay of the integrand, and we find

$$\langle \varphi_{\beta, k}^H, \varphi_{\beta, 1}^H \rangle = \frac{1}{\sqrt{\beta}} \int_{\mathbb{R}} h_{k-1}(x) e^{-x^2} dx = \frac{1}{\sqrt{\beta}} \langle \varphi_{1, k}^H, \varphi_{1, 1}^H \rangle.$$

Since the functions $(\varphi_{1, k}^H)$ are eigenfunctions of the self-adjoint operator H_1 associated to different eigenvalues, we have $\langle \varphi_{1, k}^H, \varphi_{1, 1}^H \rangle = 0$ for $k \neq 1$. Since $\langle \varphi_{1, 1}^H, \varphi_{1, 1}^H \rangle = \sqrt{\pi}$, we finally have

$$(C.7) \quad \langle \varphi_{\beta, k}^H, \varphi_{\beta, 1}^H \rangle = \begin{cases} \sqrt{\frac{\pi}{\beta}} & \text{if } k = 1 \\ 0 & \text{if } k \neq 1. \end{cases}$$

Thus, we can define

$$\tilde{\Pi}_{\beta} u = \frac{\langle \varphi_{\beta, 1}^H, u \rangle}{\langle \varphi_{\beta, 1}^H, \varphi_{\beta, 1}^H \rangle} \varphi_{\beta, 1}^H.$$

Then, for $k \in \mathbb{N}^*$

$$\tilde{\Pi}_{\beta} \varphi_{\beta, k}^H = \begin{cases} \varphi_{\beta, 1}^H & \text{if } k = 1 \\ 0 & \text{if } k \neq 1. \end{cases}$$

According to Step 3, the family $(\varphi_{\beta, k}^H)_k$ is complete, hence, by density, $\tilde{\Pi}_{\beta}$ is indeed the spectral projection Π_{β}^H .

Since $\varphi_{\beta, 1}^H(x) = e^{-\beta x^2/2}$, we can compute

$$\|\varphi_{\beta, 1}^H\|_{L^2(\mathbb{R})}^2 = \|\varphi_{\beta, 1}^H\|_{L^2(\mathbb{R})}^2 = \sqrt{\frac{\pi}{\operatorname{Re}(\beta)}},$$

and eq. (C.5) follows with eq. (C.7). \square

COROLLARY C.2. — *Let Π_{β}^H as in Proposition C.1. Let $I \subset \mathbb{R}$ be an open interval that contains 0. Let $\theta_0 \in [0, \pi/2)$. Then*

$$\operatorname{Tr} \left(\mathbf{1}_I^* \Pi_{\beta}^H \mathbf{1}_I \right) \xrightarrow[|\arg(\beta)| \leq \theta_0]{|\beta| \rightarrow \infty} 1.$$

Proof. — The reader who is not familiar with the trace of operators in infinite dimensional space may read, for instance, [Kat95, Chapter 10, §1.3–1.4]. We have $\operatorname{Tr}(\mathbf{1}_I^* \Pi_{\beta}^H \mathbf{1}_I) = \operatorname{Tr}(\mathbf{1}_I \mathbf{1}_I^* \Pi_{\beta}^H)$. Let $(\psi_k)_k$ be an orthonormal basis of $L^2(\mathbb{R})$ such that $\psi_1 = \|\mathbf{1}_I \mathbf{1}_I^* \varphi_{\beta, 1}^H\|^{-1} \mathbf{1}_I \mathbf{1}_I^* \varphi_{\beta, 1}^H$. Then,

$$\begin{aligned}
\mathrm{Tr} \left(\mathbb{1}_I \mathbb{1}_1^* \Pi_\beta^H \right) &= \sum_k \left\langle \mathbb{1}_I \mathbb{1}_I^* \Pi_\beta^H \psi_k, \psi_k \right\rangle = \left\langle \mathbb{1}_I \mathbb{1}_I^* \Pi_\beta^H \psi_1, \psi_1 \right\rangle \\
&= \left\| \mathbb{1}_I \mathbb{1}_I^* \varphi_{\beta,1}^H \right\|^{-2} \frac{\left\langle \varphi_{\beta,1}^H, \mathbb{1}_I \mathbb{1}_I^* \varphi_{\beta,1}^H \right\rangle}{\left\langle \varphi_{\beta,1}^H, \varphi_{\beta,1}^H \right\rangle} \left\langle \varphi_{\beta,1}^H, \mathbb{1}_I \mathbb{1}_I^* \varphi_{\beta,1}^H \right\rangle \\
&= \frac{\left\langle \varphi_{\beta,1}^H, \mathbb{1}_I \mathbb{1}_I^* \varphi_{\beta,1}^H \right\rangle}{\left\langle \varphi_{\beta,1}^H, \varphi_{\beta,1}^H \right\rangle} = \frac{\int_I \varphi_{\beta,1}^H(x)^2 dx}{\int_{\mathbb{R}} \varphi_{\beta,1}^H(x)^2 dx}
\end{aligned}$$

(note the $\bar{\beta}$ instead of β in the first argument of the inner products). Since $0 \in I$, the saddle point method proves that the right-hand side tends to 1 as $|\beta| \rightarrow \infty$ and $|\arg(\beta)| \leq \theta_0$. \square

BIBLIOGRAPHY

- [ABM21] Damien Allonsius, Franck Boyer, and Morgan Morancey, *Analysis of the Null Controllability of Degenerate Parabolic Systems of Grushin Type via the Moments Method*, J. Evol. Equ. **21** (2021), no. 4, 4799–4843. \uparrow 1486, 1487
- [AKBGBT16] Farid Ammar Khodja, Assia Benabdallah, Manuel González-Burgos, and Luz de Teresa, *New Phenomena for the Null Controllability of Parabolic Systems: Minimal Time and Geometrical Dependence*, J. Math. Anal. Appl. **444** (2016), no. 2, 1071–1113. \uparrow 1487
- [BBM20] Assia Benabdallah, Franck Boyer, and Morgan Morancey, *A Block Moment Method to Handle Spectral Condensation Phenomenon in Parabolic Control Problems*, Ann. Henri Lebesgue **3** (2020), 717–793. \uparrow 1487
- [BC17] Karine Beauchard and Piermarco Cannarsa, *Heat Equation on the Heisenberg Group: Observability and Applications*, J. Differ. Equations **262** (2017), no. 8, 4475–4521. \uparrow 1487
- [BCG14] Karine Beauchard, Piermarco Cannarsa, and Roberto Guglielmi, *Null Controllability of Grushin-type Operators in Dimension Two*, J. Eur. Math. Soc. **16** (2014), no. 1, 67–101. \uparrow 1481, 1485, 1486, 1512
- [BDE20] Karine Beauchard, Jérémie Dardé, and Sylvain Ervedoza, *Minimal Time Issues for the Observability of Grushin-type Equations*, Ann. Inst. Fourier **70** (2020), no. 1, 247–312. \uparrow 1486, 1487, 1488
- [Bea14] Karine Beauchard, *Null Controllability of Kolmogorov-type Equations*, Math. Control Signals Syst. **26** (2014), no. 1, 145–176. \uparrow 1487
- [BHHR15] Karine Beauchard, Bernard Helffer, Raphael Henry, and Luc Robbiano, *Degenerate Parabolic Operators of Kolmogorov Type with a Geometric Control Condition*, ESAIM, Control Optim. Calc. Var. **21** (2015), no. 2, 487–512. \uparrow 1487
- [BLR92] Claude Bardos, Gilles Lebeau, and Jeffrey Rauch, *Sharp Sufficient Conditions for the Observation, Control, and Stabilization of Waves from the Boundary*, SIAM J. Control Optim. **30** (1992), no. 5, 1024–1065. \uparrow 1486
- [BMM15] Karine Beauchard, Luc Miller, and Morgan Morancey, *2d Grushin-type Equations: Minimal Time and Null Controllable Data*, J. Differ. Equations **259** (2015), no. 11, 5813–5845. \uparrow 1486
- [Bre11] Haim Brezis, *Functional Analysis, Sobolev Spaces and Partial Differential Equations*, Universitext, Springer, 2011. \uparrow 1480, 1512

- [BS22] Nicolas Burq and Chenmin Sun, *Time Optimal Observability for Grushin Schrödinger Equation*, Anal. PDE **15** (2022), no. 6, 1487–1530. ↑1487
- [BZ09] Karine Beauchard and Enrique Zuazua, *Some Controllability Results for the 2D Kolmogorov Equation*, Ann. Inst. Henri Poincaré, Anal. Non Linéaire **26** (2009), no. 5, 1793–1815. ↑1487
- [CMV16] Piermarco Cannarsa, Patrick Martinez, and Judith Vancostenoble, *Global Carleman Estimates for Degenerate Parabolic Operators with Applications*, vol. 239, Memoirs of the American Mathematical Society, no. 1133, American Mathematical Society, 2016. ↑1487
- [Cor07] Jean-Michel Coron, *Control and Nonlinearity*, Mathematical Surveys and Monographs, vol. 143, American Mathematical Society, 2007. ↑1490
- [Dav07] Edward Brian Davies, *Linear Operators and Their Spectra*, Cambridge Studies in Advanced Mathematics, vol. 106, Cambridge University Press, 2007. ↑1516
- [DK20] Michel Duprez and Armand Koenig, *Control of the Grushin Equation: Non-Rectangular Control Region and Minimal Time*, ESAIM, Control Optim. Calc. Var. **26** (2020), article no. 3. ↑1482, 1483, 1486, 1488
- [Dol73] Szymon Dolecki, *Observability for the One-Dimensional Heat Equation*, Stud. Math. **48** (1973), 291–305. ↑1487
- [DR21] Jérémie Dardé and Julien Royer, *Critical Time for the Observability of Kolmogorov-type Equations*, J. Éc. Polytech., Math. **8** (2021), 859–894. ↑1487
- [DS99] Mouez Dimassi and Johannes Sjöstrand, *Spectral Asymptotics in the Semi-Classical Limit*, London Mathematical Society Lecture Note Series, no. 268, Cambridge University Press, 1999. ↑1481, 1494, 1496
- [FKL21] Clotilde Fermanian Kammerer and Cyril Letrouit, *Observability and Controllability for the Schrödinger Equation on Quotients of Groups of Heisenberg Type*, J. Éc. Polytech., Math. **8** (2021), 1459–1513. ↑1487
- [FR71] Hector O. Fattorini and David Lewis Russell, *Exact Controllability Theorems for Linear Parabolic Equations in One Space Dimension*, Arch. Ration. Mech. Anal. **43** (1971), no. 4, 272–292. ↑1487
- [Hat02] Allen Hatcher, *Algebraic Topology*, Cambridge University Press, 2002. ↑1515
- [Hel13] Bernard Helffer, *Spectral Theory and Its Applications*, Cambridge Studies in Advanced Mathematics, vol. 139, Cambridge University Press, 2013. ↑1480, 1491, 1512, 1516, 1518
- [HSS05] Frédéric Hérau, Johannes Sjöstrand, and Christiaan C. Stolk, *Semiclassical Analysis for the Kramers–Fokker–Planck Equation*, Commun. Partial Differ. Equations **30** (2005), no. 5-6, 689–760. ↑1483
- [Kat95] Tosio Kato, *Perturbation Theory for Linear Operators*, Classics in Mathematics, no. 132, Springer, 1995. ↑1509, 1519
- [Koe17] Armand Koenig, *Non-Null-Controllability of the Grushin Operator in 2D*, C. R. Math. Acad. Sci. Paris **355** (2017), no. 12, 1215–1235. ↑1496, 1508
- [Koe20] ———, *Lack of Null-Controllability for the Fractional Heat Equation and Related Equations*, SIAM J. Control Optim. **58** (2020), no. 6, 3130–3160. ↑1487
- [KSTV15] David Krejčířík, Petr Siegl, Miloš Tater, and Joe Viola, *Pseudospectra in Non-Hermitian Quantum Mechanics*, J. Math. Phys. **56** (2015), article no. 103513. ↑1516
- [Let23] Cyril Letrouit, *Subelliptic Wave Equations Are Never Observable*, Anal. PDE **16** (2023), no. 3, 643–678. ↑1487
- [LL22] Camille Laurent and Matthieu Léautaud, *Tunneling Estimates and Approximate Controllability for Hypoelliptic Equations*, Memoirs of the American Mathematical Society, vol. 276, American Mathematical Society, 2022. ↑1487

- [LL23] ———, *On uniform controllability of 1D transport equations in the vanishing viscosity limit*, C. R. Math. Acad. Sci. Paris **361** (2023), 265–312. ↑1486
- [LS23] Cyril Letrouit and Chenmin Sun, *Observability of Baouendi–Grushin-type Equations through Resolvent Estimates*, J. Inst. Math. Jussieu **22** (2023), no. 2, 541–579. ↑1487
- [Rud86] Walter Rudin, *Real and Complex Analysis*, third ed., McGraw Hill Education, 1986. ↑1493, 1499, 1509
- [Rud91] ———, *Functional Analysis*, 2nd ed., McGraw-Hill, 1991. ↑1510
- [Yeg63] Yu. V. Yegorov, *Some problems in the theory of optimal control*, Zh. Vychisl. Mat. Mat. Fiz. **3** (1963), 887–904. ↑1487

Manuscript received on 2nd July 2022,
revised on 9th May 2023,
accepted on 18th June 2023.

Recommended by Editors P. Gérard and F. Hérau.
Published under license CC BY 4.0.



eISSN: 2644–9463

This journal is a member of Centre Mersenne.



Jérémi DARDÉ
Institut de Mathématiques
de Toulouse, UMR 5219,
Université de Toulouse,
CNRS, UPS,
F-31062 Toulouse Cedex 9, France
jeremi.darde@math.univ-toulouse.fr

Armand KOENIG
Institut de Mathématiques
de Toulouse, UMR 5219,
Université de Toulouse,
CNRS, UPS,
F-31062 Toulouse Cedex 9, France
armand.koenig@math.univ-toulouse.fr

Julien ROYER
Institut de Mathématiques
de Toulouse, UMR 5219,
Université de Toulouse,
CNRS, UPS,
F-31062 Toulouse Cedex 9, France
julien.royer@math.univ-toulouse.fr