



ANNALES
HENRI LEBESGUE

ANATOLE GAUDIN

HODGE DECOMPOSITIONS AND
MAXIMAL REGULARITIES FOR
HODGE LAPLACIANS IN
HOMOGENEOUS FUNCTION
SPACES ON THE HALF-SPACE

DECOMPOSITIONS DE HODGE ET
RÉGULARITÉ MAXIMALE POUR LES
LAPLACIENS DE HODGE DANS LES
ESPACES DE FONCTIONS HOMOGÈNES SUR
LE DEMI-ESPACE

ABSTRACT. — In this article, the Hodge decomposition for any degree of differential forms is investigated on the whole space \mathbb{R}^n and the half-space \mathbb{R}_+^n on different scales of function spaces namely the homogeneous and inhomogeneous Besov and Sobolev spaces, $\dot{H}^{s,p}$, $\dot{B}_{p,q}^s$, $H^{s,p}$ and $B_{p,q}^s$, for $p \in (1, +\infty)$, $s \in (-1 + \frac{1}{p}, \frac{1}{p})$. The bounded holomorphic functional calculus, and other functional analytic properties, of Hodge Laplacians is also investigated in the half-space, and yields similar results for Hodge–Stokes and other related operators via the proven Hodge decomposition. As consequences, the homogeneous operator and interpolation theory revisited

Keywords: homogeneous Sobolev spaces, homogeneous Besov spaces, differential forms, Hodge decomposition, interpolation with boundary conditions, maximal regularity, evolutionary Stokes systems, half-space.

2020 *Mathematics Subject Classification:* 35Q35, 42B37, 46B70, 46E35, 58A10.

DOI: <https://doi.org/10.5802/ahl.224>

by Danchin, Hieber, Mucha and Tolksdorf is applied to homogeneous function spaces subject to boundary conditions and leads to various maximal regularity results with global-in-time estimates that could be of use in fluid dynamics. Moreover, the bond between the Hodge Laplacian and the Hodge decomposition will even enable us to state the Hodge decomposition for higher order Sobolev and Besov spaces with additional compatibility conditions, for regularity index $s \in (-1 + \frac{1}{p}, 2 + \frac{1}{p})$. In order to make sense of all those properties in desired function spaces, we also give appropriate meaning of partial traces on the boundary in the appendix.

“La raison d’être” of this paper lies in the fact that the chosen realization of homogeneous function spaces is suitable for non-linear and boundary value problems, but requires a careful approach to reprove results that are already morally known.

RÉSUMÉ. — Dans cet article, la décomposition de Hodge pour tout degré de formes différentielles est étudiée sur l’espace entier \mathbb{R}^n et le demi-espace \mathbb{R}_+^n pour différentes familles d’espaces de fonctions, à savoir les espaces de Besov et de Sobolev homogènes et inhomogènes, $\dot{H}^{s,p}$, $\dot{B}_{p,q}^s$, $H^{s,p}$ et $B_{p,q}^s$, pour tout $p \in (1, +\infty)$ et $s \in (-1 + \frac{1}{p}, \frac{1}{p})$. Le calcul fonctionnel holomorphe borné ainsi que d’autres propriétés des Laplaciens de Hodge sont également étudiés dans le demi-espace, ce qui donne des résultats similaires pour les opérateurs de Hodge–Stokes et d’autres opérateurs qui lui sont liés, via la décomposition de Hodge qui est démontrée en amont. En conséquence, la théorie des opérateurs et la théorie de l’interpolation dans leur version homogène, revisitées par Danchin, Hieber, Mucha et Tolksdorf, sont appliquées aux espaces de fonctions homogènes soumis à des conditions au bord. Cela conduit à divers résultats de régularité maximale avec des estimations globales en temps, pouvant être utiles en dynamique des fluides. De plus, le lien entre le Laplacien de Hodge et la décomposition de Hodge nous permettra même d’énoncer la décomposition de Hodge pour des espaces de Sobolev et de Besov d’ordre supérieur avec des conditions de compatibilité supplémentaires, pour des indices de régularité $s \in (-1 + \frac{1}{p}, 2 + \frac{1}{p})$. Afin de donner un sens à toutes ces propriétés dans les espaces fonctionnels souhaités, nous donnons également une signification appropriée aux traces partielles sur le bord en annexe.

La raison d’être de cet article réside dans le fait que la réalisation choisie des espaces de fonctions homogènes est adaptée aux problèmes non linéaires et aux problèmes avec conditions au bord, mais nécessite une approche minutieuse pour redémontrer des résultats déjà moralement connus.

CONTENTS

1. Introduction	1459
1.1. Motivations and interests	1459
1.1.1. One Laplacian to rule (almost) them all: the differential form formalism and the Hodge decomposition	1459
1.1.2. Global-in-time estimates in L^q -maximal regularity: the role of homogeneous function spaces and their interpolation	1463
1.2. Notations, definitions, and usual concepts	1467
1.2.1. Smooth and measurable functions	1467
1.2.2. (Bi)Sectorial operators on Banach spaces	1468
1.2.3. Interpolation of normed vector spaces	1470
1.2.4. Sobolev and Besov spaces on \mathbb{R}^n	1470
1.2.5. Homogeneous Sobolev and Besov spaces on \mathbb{R}_+^n	1473
1.2.6. Operators on Sobolev and Besov spaces	1475
Acknowledgments	1475
2. Hodge Laplacians, Hodge decompositions and Hodge–Stokes operators	1475
2.1. Differential forms on Euclidean space, and corresponding function spaces	1476
2.2. The case of the whole space	1481
2.3. The case of the half-space \mathbb{R}_+^n	1487

2.3.1. L^2 -theory for Hodge Laplacians and the Hodge decomposition	1487
2.3.2. $\dot{H}^{s,p}$ and $\dot{B}_{p,q}^s$ -theory for Hodge Laplacians and the Hodge decomposition	1497
2.3.3. Hodge–Stokes and Hodge–Maxwell operators	1504
3. L^q -maximal regularity with global-in-time estimates	1505
3.1. Interpolation of homogeneous $\dot{H}^{s,p}$ -domains of operators	1508
3.2. Maximal regularity for Hodge Laplacians and related operators	1521
3.3. Maximal regularity for the Stokes system with Navier-slip boundary conditions	1523
Appendix A. Partial traces of differential forms	1525
References	1530

1. Introduction

1.1. Motivations and interests

1.1.1. One Laplacian to rule (almost) them all: the differential form formalism and the Hodge decomposition

The study of incompressible fluid dynamics, and in particular the treatment of Navier–Stokes equations, relies mostly on the Helmholtz decomposition of a vector field in appropriate function spaces. The Helmholtz decomposition of vector field $u : \Omega \rightarrow \mathbb{C}^n$ is given by a vector field $v : \Omega \rightarrow \mathbb{C}^n$ and a function $q : \Omega \rightarrow \mathbb{C}$, such that

$$u = v + \nabla q \text{ and } \operatorname{div} v = 0 \left(\text{with possibly } v \cdot \nu_{|\partial\Omega} = 0 \right).$$

This point is central since the incompressibility condition for the velocity of a fluid u is carried over by the condition $\operatorname{div} u = 0$.

In the interest of the Navier–Stokes and related equations, one wants the above decomposition to hold topologically in an appropriate normed vector space of functions⁽¹⁾ with uniqueness (up to a constant for q). It is indeed true in $L^2(\Omega, \mathbb{C}^n)$, since \mathbb{P} , the usual Helmholtz–Leray projector on divergence free vector fields with null normal trace at the boundary, i.e. such that

$$\mathbb{P} : L^2(\Omega, \mathbb{C}^n) \rightarrow L^2_\sigma(\Omega) = \left\{ u \in L^2(\Omega, \mathbb{C}^n) \mid \operatorname{div} u = 0, u \cdot \nu_{|\partial\Omega} = 0 \right\},$$

is well-defined, linear, bounded, and unique by construction of the orthogonal projector on a closed subspace of a Hilbert space, here $L^2_\sigma(\Omega) \subset L^2(\Omega, \mathbb{C}^n)$. It gives the classical orthogonal and topological Helmholtz decomposition, see [Soh01, Chapter 2, Section 2.5],

$$L^2(\Omega, \mathbb{C}^n) = L^2_\sigma(\Omega) \oplus \overline{\nabla H^{1,2}(\Omega, \mathbb{C})},$$

for any (bounded) Lipschitz domain Ω , see [Soh01, Lemma 2.5.3]. Here $H^{1,2}(\Omega, \mathbb{C})$ is the standard L^2 -Sobolev space of order 1 on Ω .

⁽¹⁾From here the divergence will be understood in the distributional sense.

The L^2 -theory for the Helmholtz decomposition on a domain Ω relies mostly on pure Hilbertian operator theory. However, the question about the L^p -theory, $p \neq 2$, i.e. to know if

$$(1.1) \quad L^p(\Omega, \mathbb{C}^n) = L^p_\sigma(\Omega) \oplus \overline{\nabla H^{1,p}(\Omega, \mathbb{C})},$$

(or even the Sobolev or Besov counterpart) is actually a harder question, which falls generally in the field of harmonic analysis. The underlying range of Lebesgue and Sobolev exponents for which such decomposition holds will generally depend on the regularity of the boundary and the geometry of the domain Ω .

The L^p setting has been widely studied, we mention the work of Fabes, Mendez and Mitrea, [FMM98, Theorem 12.2], where the result has been proven for bounded Lipschitz domains: (1.1) holds whenever $p \in (3/2 - \varepsilon, 3 + \varepsilon)$. The work of Sohr and Simader [SS92, Theorem 1.4] yields (1.1) for C^1 bounded and exterior domains, allowing $p \in (1, +\infty)$. For general unbounded domains, when $p \neq 2$, the decomposition (1.1) may fail: see the counterexample by Bogovskiĭ [Bog86, Section 2]. Tolksdorf has shown in his PhD dissertation [Tol17, Theorem 5.1.10] that (1.1) is true for all $p \in (\frac{2n}{2n+1} - \varepsilon, \frac{2n}{2n-1} + \varepsilon)$, provided Ω is a special Lipschitz domain, $\varepsilon > 0$ depending on Ω . We also mention the works of Farwig, Kozono and Sohr where the decomposition is investigated in a more exotic setting in [FKS05, FKS07] for general uniform C^1 unbounded domains.

Our interest here is the case of the half-space \mathbb{R}^n_+ , where the Helmholtz decomposition is mainly known to be true on $L^p(\mathbb{R}^n_+, \mathbb{C}^n)$ for all $p \in (1, +\infty)$, see [Gal11, Remark III.1.2]: we aim to generalize this result to the scale of inhomogeneous, and homogeneous Sobolev and Besov spaces on the half-space. To be more precise, we want to investigate decompositions of the type

$$(1.2) \quad \dot{H}^{s,p}(\mathbb{R}^n_+, \mathbb{C}^n) = \dot{H}^{s,p}_\sigma(\mathbb{R}^n_+, \mathbb{C}^n) \oplus \overline{\nabla \dot{H}^{s+1,p}(\mathbb{R}^n_+, \mathbb{C}^n)},$$

and similarly for Besov spaces, and their inhomogeneous counterparts, provided $s \in \mathbb{R}$, $p \in (1, +\infty)$.

In the scale of inhomogeneous and homogeneous Besov and Sobolev spaces on bounded and exterior $C^{2,1}$ domains the Helmholtz decomposition was shown by Fujiwara and Yamazaki [FY06, Theorem 3.1]: the Helmholtz decomposition holds on $H^{s,p}(\Omega, \mathbb{C}^n)$ and $B^s_{p,q}(\Omega, \mathbb{C}^n)$, $p \in (1, +\infty)$, $s \in (-1 + 1/p, 1/p)$, $q \in [1, +\infty]$, even allowing $p = 1, +\infty$ in case of Besov spaces. We also mention the work of Monniaux and Mitrea [MM08, Proposition 2.16] on bounded Lipschitz domain where the result is true for (inhomogeneous) Sobolev spaces that lie near the family $(H^{s,2})_{|s| < 1/2}$.

It has been notified in several works, e.g. see [GHT13, Introduction], [MS18, Section 4], that the following Laplace operator acting on vector fields,

$$(1.3) \quad -\Delta_{\mathcal{H}} u := -\Delta u = \text{curl curl } u - \nabla \text{div } u, \text{ and } [u \cdot \nu|_{\partial\Omega} = 0, \nu \times \text{curl } u|_{\partial\Omega} = 0]$$

called the⁽²⁾ Hodge Laplacian, has a strong bond with, and respects, the Helmholtz decomposition in the sense that for all u in the domain of above Laplacian, $\mathbb{P}u$ also lies in, and we have

⁽²⁾In fact, this is a Hodge Laplacian, the one with normal boundary conditions, we do not make the distinction here for introductory purposes.

$$-\mathbb{P}\Delta u = \operatorname{curl} \operatorname{curl} u = -\Delta \mathbb{P}u, \quad \text{and} \quad \left[\mathbb{P}u \cdot \nu|_{\partial\Omega} = 0, \nu \times \operatorname{curl} \mathbb{P}u|_{\partial\Omega} = 0 \right].$$

Therefore, since the Hodge Laplacian and the Helmholtz–Leray projector seem to copy the corresponding behavior of the whole space, it seems reasonable to infer that

$$(1.4) \quad \mathbb{P} = \mathbb{I} + \nabla \underline{\operatorname{div}} (-\Delta_{\mathcal{H}})^{-1},$$

where $\underline{\operatorname{div}}$ drives a boundary condition $\nu \cdot u|_{\partial\Omega} = 0$.

But, the above use of curl operators restricts us to the three dimensional case. We can avoid such trouble, by means of the differential forms formalism, so that (1.3) becomes

$$(1.5) \quad -\Delta_{\mathcal{H}}u := -\Delta u = (d^*d + dd^*)u = (d + d^*)^2 u, \quad \text{and} \quad \left[\nu \lrcorner u|_{\partial\Omega} = 0, \nu \lrcorner du|_{\partial\Omega} = 0 \right]$$

where $d : \Lambda^k \rightarrow \Lambda^{k+1}$ is the exterior derivative, defined on the complexified exterior algebra of \mathbb{R}^n , $\Lambda = \Lambda^0 \oplus \Lambda^1 \oplus \dots \oplus \Lambda^n$, and satisfies $d^2 = 0$. The operator $d^* : \Lambda^k \rightarrow \Lambda^{k-1}$ is the formal dual operator of d , satisfying also $(d^*)^2 = 0$ so that on \mathbb{R}^3 , we can make the identifications

$$\begin{aligned} d|_{\Lambda^1} &= \operatorname{curl}, & d|_{\Lambda^0} &= \nabla, \\ d^*|_{\Lambda^2} &= \underline{\operatorname{curl}}, & d^*|_{\Lambda^1} &= -\underline{\operatorname{div}}, \\ \nu \lrcorner ()|_{\Lambda^1} &= \nu \cdot (), & \nu \lrcorner ()|_{\Lambda^2} &= \nu \times (). \end{aligned}$$

The $\underline{\operatorname{curl}}$ operator drives a boundary condition $\nu \times u|_{\partial\Omega} = 0$.

Notice this definition still makes sense for differential forms of any degree, in arbitrary dimension. One would check that (1.5) reduce to the Neumann Laplacian in the case of 0-forms identified with scalar-valued functions.

Going back to the case of vector fields, instead of (1.4), the above formalism and the fact that d and d^* are nilpotent, and then commutes (at least formally) with $\Delta_{\mathcal{H}}$, we may infer the next formula, similar to the one mentioned in [ACDH04, Section 5]:

$$(1.6) \quad \mathbb{P} = \mathbb{I} - dd^*(-\Delta_{\mathcal{H}})^{-1} = \mathbb{I} - d(-\Delta_{\mathcal{H}})^{-1/2}d^*(-\Delta_{\mathcal{H}})^{-1/2}.$$

Under the use of the differential forms formalism, the desired Helmholtz decomposition (1.2) becomes, for $0 \leq k \leq n$ different degrees of differential forms,

$$(1.7) \quad \dot{H}^{s,p}(\mathbb{R}_+^n, \Lambda^k) = \dot{H}_{n,\sigma}^{s,p}(\mathbb{R}_+^n, \Lambda^k) \oplus \overline{d\dot{H}^{s+1,p}(\mathbb{R}_+^n, \Lambda^{k-1})}$$

which is called the *Hodge decomposition* instead of the Helmholtz decomposition. Here, the space $\dot{H}_{n,\sigma}^{s,p}(\mathbb{R}_+^n, \Lambda^k)$ stands for k -differential forms u whose coefficients lie in $\dot{H}^{s,p}(\mathbb{R}_+^n, \mathbb{C})$, and such that $d^*u = 0$, and $\nu \lrcorner u|_{\partial\mathbb{R}_+^n} = 0$.

The Hodge decomposition for differential forms is treated by Schwarz [Sch95, Theorem 2.4.2, Theorem 2.4.14] on smooth compact Riemannian manifold M with smooth boundary where the decomposition is stated on $H^{k,p}(M)$, $k \in \mathbb{N}$, $p \in (1, +\infty)$. For the case of Ω a bounded Lipschitz domain of \mathbb{R}^n , we refer to the work of Monniaux and M^cIntosh [MM18, Theorem 4.3, Theorem 7.1] where the Hodge decomposition is proved to be true on $L^p(\Omega, \Lambda)$ for all $p \in (\frac{2n}{2n+1} - \varepsilon, \frac{2n}{2n-1} + \varepsilon)$ where $\varepsilon > 0$

depends on Ω . The bounded holomorphic functional calculus of the Hodge Laplacian is also proved for the same range of indices. One may also consult the work of Mitrea and Monniaux, and Hofmann, Mitrea and Monniaux, [MM09b, HMM11], for the treatment of the Hodge Laplacian on bounded Lipschitz domains of compact Riemannian manifolds, where functional analytic properties like analyticity of the generated semigroup, or boundedness of associated Riesz transforms are investigated.

One may wonder about the superficiality of proving an identity like (1.7) for general differential forms, instead of vector fields (differential forms of degree 1, $n - 1$) only. In fact, the differential forms formalism has shown its efficiency, allowing to treat some partial differential equations initially restricted to the three-dimensional setting in arbitrary dimension. See for instance [Mon21, Den22], where the magnetohydrodynamical (MHD) system is treated, so that either the triplet $\Lambda^1, \Lambda^2, \Lambda^3$ or the triplet $\Lambda^{n-3}, \Lambda^{n-2}, \Lambda^{n-1}$ are involved. Indeed, the magnetic field is in fact not an effective vector field but a 2-form, identified, when $n = 3$, with a vector field. We also mention that reformulation using differential forms for this kind of systems allows looking at vorticity-like formulation of the Navier-Stokes (and related) equations, it is also purely intrinsic so that one can perform a similar treatment on manifolds.

To reach our goal, the idea will be to prove that the formula (1.6) holds on $L^2(\mathbb{R}_+^n, \Lambda)$, yielding an operator for which we can also prove its boundedness on Sobolev and Besov spaces, so that we are able to obtain the next theorem.

THEOREM 1.1 (see Theorem 2.33 & Corollary 3.14). — *Let $p \in (1, +\infty)$, $s \in (-1 + 1/p, 1/p)$, and let $k \in \llbracket 0, n \rrbracket$. It holds that*

- (1) *The (generalized) Helmholtz–Leray projector is well-defined and bounded as an operator*

$$\mathbb{P} : \dot{H}^{s,p}(\mathbb{R}_+^n, \Lambda^k) \longrightarrow \dot{H}_{n,\sigma}^{s,p}(\mathbb{R}_+^n, \Lambda^k).$$

Moreover, the following identity is true

$$\mathbb{P} = \text{I} - \text{d}(-\Delta_{\mathcal{H}})^{-\frac{1}{2}} \text{d}^*(-\Delta_{\mathcal{H}})^{-\frac{1}{2}}.$$

- (2) *The following Hodge decomposition holds*

$$\dot{H}^{s,p}(\mathbb{R}_+^n, \Lambda^k) = \dot{H}_{n,\sigma}^{s,p}(\mathbb{R}_+^n, \Lambda^k) \oplus \dot{H}_{\gamma}^{s,p}(\mathbb{R}_+^n, \Lambda^k).$$

Moreover, the result remains true if we replace

- $\dot{H}^{s,p}$ by $\dot{B}_{p,q}^s$, $q \in [1, +\infty]$;
- (\dot{H}, \dot{B}) by (H, B) .

The symbol X_{γ} stands for the range of $\text{I} - \mathbb{P}$ in X .⁽³⁾

The way we reach Theorem 1.1 through intermediate results and proofs is so that we recover many different properties of the Hodge Laplacian as well as its bounded holomorphic functional calculus on Sobolev and Besov spaces almost for free. This is due to the particular structure of the boundary of \mathbb{R}_+^n , and the properties of the

⁽³⁾The subscript (or exponent in case of Besov spaces) γ is a legacy of the writing of G spaces as spaces of gradients of scalar functions in the case of vector fields.

Laplacian on the whole space \mathbb{R}^n . This, above Theorem 1.1, and the fact that one can define the Hodge–Stokes operator as

$$u \in \dot{D}_p^s(\mathbb{A}_{\mathcal{H}}) = \mathbb{P}\dot{D}_p^s(\Delta_{\mathcal{H}}) \text{ and } \mathbb{A}_{\mathcal{H}}u := -\Delta_{\mathcal{H}}u = d^*du,$$

will yield automatically

THEOREM 1.2 (see Theorem 2.35). — *Let $p \in (1, +\infty)$, $s \in (-1 + 1/p, 1/p)$. For all $\mu \in (0, \pi)$, the operator $\mathbb{A}_{\mathcal{H}}$ admits a bounded $(\mathbf{H}^\infty(\Sigma_\mu)$ -)holomorphic functional calculus on $\dot{H}_{n,\sigma}^{s,p}(\mathbb{R}_+^n, \Lambda)$. Moreover, the result remains true if we replace*

- $\dot{H}^{s,p}$ by $\dot{B}_{p,q}^s$, $q \in [1, +\infty)$;
- (\dot{H}, \dot{B}) by (H, B) .

We mention that our strategy is not morally so different from the one presented in [GHT13, Beginning of Section 4], identifying some Neumann and Dirichlet boundary conditions on various components. However, the treatment of boundary values is done in a more careful way, adapted with the scales of homogeneous function spaces, thanks to a weak-strong correspondence of (partial) traces by means of appropriate results in the Appendix A.

1.1.2. Global-in-time estimates in L^q -maximal regularity: the role of homogeneous function spaces and their interpolation

Another tool which is central in the study of parabolic equations and also for a large class of fluid dynamics problems is the L^q -maximal regularity.

The general problem of global in time L^q -maximal regularity is: for a closed operator $(D(A), A)$ on a Banach space X , let us consider the evolution equation

$$(1.8) \quad \begin{cases} \partial_t u(t) + Au(t) = f(t), t \in (0, +\infty), \\ u(0) = 0. \end{cases}$$

Provided $q \in [1, +\infty]$ and $f \in L^q((0, +\infty), X)$, can we solve uniquely (1.8), with an a priori estimate

$$(1.9) \quad \|(\partial_t u, Au)\|_{L^q((0,+\infty),X)} \lesssim \|f\|_{L^q((0,+\infty),X)} \text{ ?}$$

When X is a UMD Banach space (i.e. a space such that the Hilbert transform is bounded on $L^r(\mathbb{R}, X)$ for one (or equivalently all) $r \in (1, +\infty)$), the problem has been extensively studied in [Ama95, Chapter III, Section 4], [DHP03] and [KW04]. It has been proved in this case, that the truthfulness of (1.9) for $q \in (1, +\infty)$ is equivalent to the \mathcal{R} -boundedness of the resolvent A of angle $\phi_A^{\mathcal{R}} < \frac{\pi}{2}$, i.e. if $\sigma(A) \subset \bar{\Sigma}_\phi$, and for some $\frac{\pi}{2} > \mu > \phi$, the set

$$\left\{ \lambda(\lambda I - A)^{-1} \right\}_{\lambda \in \mathbb{C} \setminus \Sigma_\mu}$$

is \mathcal{R} -bounded, and $\phi_A^{\mathcal{R}}$ is the infimum on all such μ . More precisely, for some $\mu \in (\phi, \frac{\pi}{2})$, for all $(\lambda_j)_{j \in \mathbb{N}} \subset \mathbb{C} \setminus \Sigma_\mu$, all $(f_j)_{j \in \mathbb{N}} \subset X$, we have for all $N \in \mathbb{N}$

$$\left\| \sum_{j=0}^N r_j(\cdot) \lambda_j (\lambda_j I - A)^{-1} f_j \right\|_{L^2((0,1),X)} \lesssim_{X,A,\mu} \left\| \sum_{j=0}^N r_j(\cdot) f_j \right\|_{L^2((0,1),X)},$$

where $(r_j)_{j \in \mathbb{N}}$ is the Rademacher system of functions, and the implicit constant is independent of N . This result was initially due to Weis, see [Wei01, Theorem 4.2]. For a wide review of the concept of \mathcal{R} -boundedness and its applications to maximal regularity, we refer to, e.g., [DHP03, KW04] and [DK07].

It has been shown in many cases that Stokes operators satisfy the L^q -maximal regularity on $L^p_\sigma(\Omega)$, for various classes of open sets Ω for $p, q \in (1, +\infty)$, with various boundary conditions and this has been widely used to treat various fluid dynamics problems, mainly Navier–Stokes equations. See for instance, and the list is far from being exhaustive, [GS91, HM13, HNPS16, Hie20, MM09a, Mon13, Mon21, Tol18, TW20].

Getting back to the abstract problem (1.8), when X is UMD, $q \in (1, +\infty)$ and A is invertible, i.e., $0 \in \rho(A)$, it is known that the solution u belongs to $C^0_b(\mathbb{R}_+, (X, D(A))_{1-1/q, q})$ with the estimate

$$(1.10) \quad \|u\|_{L^\infty(\mathbb{R}_+, (X, D(A))_{1-1/q, q})} \lesssim_{A, q} \|f\|_{L^q(\mathbb{R}_+, X)},$$

where $(\cdot, \cdot)_{\theta, r}$, $(\theta, r) \in (0, 1) \times [1, +\infty]$, stands for the real interpolation functor; check for instance [Ama95, Chapter III, Theorem 4.10.2].

If $0 \in \sigma(A)$, one only has $u \in C^0([0, T], (X, D(A))_{1-1/q, q})$, with for each $T < +\infty$,

$$(1.11) \quad \|u\|_{L^\infty([0, T], (X, D(A))_{1-1/q, q})} \lesssim_{A, q, T} \|f\|_{L^q(\mathbb{R}_+, X)},$$

where the implicit constant blows up as T goes to $+\infty$. Notice that this is the case for the (negative) Laplacian $A = -\Delta$ on $X = L^p(\mathbb{R}^n)$, $p \in (1, +\infty)$, which is quite inconvenient, see [Gau24a, Section 1] and the references therein for more details.

Another issue is that one cannot reach L^1 and L^∞ -maximal regularity estimates through above theory, but one may recover such kind of results if X is replaced by a real interpolation space between X and $D(A)$, say $Y_r^\theta := (X, D(A))_{\theta, r}$ $(\theta, r) \in (0, 1) \times [1, +\infty]$. Indeed, a theorem of Da Prato and Grisvard, see [DPG75, Theorem 4.15], gives us that, provided either

- $0 \in \rho(A)$ and $T \in (0, +\infty]$,
- $0 \in \sigma(A)$ and $T \in (0, +\infty)$,

for $q \in [1, +\infty)$, $\theta \in (0, 1)$, the solution u to (1.8) belongs to $C^0_b([0, T], Y_q^{1+\theta-1/q})$ and satisfies

$$(1.12) \quad \|u\|_{L^\infty([0, T], Y_q^{1+\theta-1/q})} \lesssim_{A, q, (T)} \|(\partial_t u, Au)\|_{L^q((0, T), Y_q^\theta)} \lesssim_{A, q, (T)} \|f\|_{L^q((0, T), Y_q^\theta)}.$$

If one wants to recover global in time estimates in (1.12) by means of [DPG75, Theorem 4.15], we have to assume that A is invertible, hence $0 \in \rho(A)$. However, for $q = 1$ similar estimates have been shown for several non-invertible operators and were of major importance to achieve existence in critical function spaces for some fluid dynamic problems like global well-posedness of Navier–Stokes equations, even for inhomogeneous flows, or free boundary problems, see for instance [Che99, DM09, DM15, OS16, OS21, OS22].

While the work of Ogawa and Shimizu [OS22] provides a powerful framework for many applications, we are mainly restricted to a specific class of second order elliptic operators with “smooth enough” coefficients. A different and more abstract approach was brought by the recent work of Danchin, Hieber, Mucha and Tolksdorf [DHMT21],

where the idea was to give a homogeneous version of the Da Prato–Grisvard theorem [DPG75, Theorem 4.15], in the sense that (1.12) holds with implicit constant uniform with respect to the time variable even when $0 \in \sigma(A)$. But further assumptions have to be made, mainly the injectivity of A on X . Their idea was to replace the real interpolation space $Y_q^\theta = (X, D(A))_{\theta,q}$ in (1.12) by

$$(X, D(\mathring{A}))_{\theta,q}$$

where $D(\mathring{A})$ is called the homogeneous domain of A and stands *morally* for the closure of $D(A)$ with respect to the (semi-)norm $\|A \cdot\|_X$. Such kind of investigation was already started by Haak, Haase, and Kuntzmann in [HHK06], then developed in Haase’s book [Haa06, Chapter 6], but the *completion* is considered instead of the closure.

If A is a non-degenerate elliptic operator of order $2m$ equal to its principal part, densely defined on $X = L^p(\mathbb{R}^n)$, such that $D(A) = H^{2m,p}(\mathbb{R}^n)$, we should have $D(\mathring{A}) = \dot{H}^{2m,p}(\mathbb{R}^n)$ which encounter no trouble of definition, if one consider the construction of homogeneous Sobolev and Besov spaces as equivalence classes of tempered distributions up to a polynomial, see [Haa06, Chapter 8, Section 3], [Tri83, Chapter 5]. In this case, we obtain

$$(X, D(A))_{\theta,q} = B_{p,q}^{2m\theta}(\mathbb{R}^n) \quad \text{and} \quad (X, D(\mathring{A}))_{\theta,q} = \dot{B}_{p,q}^{2m\theta}(\mathbb{R}^n).$$

However, if one wants to consider a similar problem on a domain, here the half-space \mathbb{R}_+^n , with some boundary conditions, with the definition of homogeneous function spaces as class of tempered distributions up to a polynomial, it is not clear that we can make a proper meaning of boundary conditions or traces. To overcome such difficulties, a construction of homogeneous Sobolev spaces $\dot{H}^{s,p}(\mathbb{R}_+^n)$ and a review of homogeneous Besov spaces on \mathbb{R}_+^n allowing to check interpolation between homogeneous spaces, and to recover boundary conditions in some cases, is done in [DHMT21, Chapters 3 & 4] continued in [Gau24b]. This construction is based on, and consistent with, the one of homogeneous Besov spaces on the whole space achieved in [BCD11, Chapter 2]. This leads in some cases to non-complete normed vector spaces as $D(\mathring{A})$ could be if one wants to consider it as the (moral) closure of $D(A)$ in an appropriate subspace of $\dot{H}^{2m,p}(\mathbb{R}_+^n)$, which may be not complete. That is why the construction of the homogeneous domain of A , its real interpolation spaces with X , and the homogeneous Da Prato–Grisvard theorem from [DHMT21, Chapter 2] are interesting: they allow $D(\mathring{A})$ to be a non-complete normed vector space⁽⁴⁾. This could be necessary if one wants to deal with boundary conditions. One can even recover the construction given in [Haa06, Chapter 6] when the completion is used instead of the closure to construct the homogeneous domain.

We also mention that the homogeneous operator and interpolation theory revisited by Danchin, Hieber, Mucha, and Tolksdorf in [DHMT21, Chapter 2] also gives a way to circumvent the lack of global-in-time estimates in the usual L^q -maximal regularity

⁽⁴⁾We notice that real interpolation of non-complete vector space makes sense, see [BL76, Chapter 3] where completeness is not needed to deal with the K -method.

framework, for the trace estimate issue (1.11). This has been done by the author in a previous paper, see [Gau24a, Sections 1, 2 & 4].

Finally, if one applies the homogeneous interpolation and then the homogeneous Da Prato–Grisvard theorem as done in [DHMT21, Chapters 2, 3 & 4], choosing $X \subset L^p(\mathbb{R}_+^n)$ to be a closed subspace, and $D(A)$ to be a closed subset of $H^{2m,p}(\mathbb{R}_+^n)$, it would lead to $L_t^1(\dot{B}_{p,1}^{2m\theta})$ -maximal regularity results with $\theta \in (0, 1)$. Proceeding this way disallow to obtain $L_t^1(\dot{B}_{p,1}^0)$ or $L_t^1(\dot{B}_{p,1}^\alpha)$ -maximal regularity results, for $\alpha < 0$. Our idea is to replace the use of $L^p(\mathbb{R}_+^n)$ as a ground space by $\dot{H}^{s,p}(\mathbb{R}_+^n)$, with $p \in (1, +\infty)$, $s \in (-1 + 1/p, 1/p)$, so that we may expect to realize A on $\dot{H}^{s,p}(\mathbb{R}_+^n)$ with domain

$$D(A) \subset \dot{H}^{s,p}(\mathbb{R}_+^n) \cap \dot{H}^{s+2m,p}(\mathbb{R}_+^n).$$

Therefore, for $s \in (-1 + 1/p, 1/p)$ and $\theta \in (0, 1)$, it seems reasonable to expect $L_t^1(\dot{B}_{p,1}^{s+2m\theta})$ -maximal regularity results, and then recover maximal regularity for some non-positive index of regularity.

To reach such realizations of A on homogeneous Sobolev spaces of fractional order on the whole space or on the half-space, we are going to use the construction started [DHMT21, Chapter 3], and continued in [Gau24b, Sections 2 & 3]. The Appendix A is dedicated to the meaning of partial traces in such function spaces to ensure that one can realize operators with boundary conditions on $\dot{H}^{s,p}(\mathbb{R}_+^n)$, provided $s \in (-1 + 1/p, 1/p)$, see also [Gau24b, Section 4] for usual trace results. We will also provide additional tools that will be useful to compute homogeneous interpolation spaces in presence of boundary conditions, as in Section 3.

In our case, considering first the Hodge Laplacian, then the Hodge–Stokes operator, we will be able to apply the homogeneous Da Prato–Grisvard theorem [DHMT21, Theorem 2.20], as well as the usual L^q -maximal regularity for UMD Banach spaces or [Gau24a, Theorem 4.7], to reach various maximal regularity results as the next one.

THEOREM 1.3 (see Theorems 3.17, 3.18, 3.19 & 3.22). — *Let $p \in (1, +\infty)$, $q \in [1, +\infty)$, $s \in (-1 + 1/p, 1/p + 2/q)$, $s, s + 2 - 2/q \notin \mathbb{N} + \frac{1}{p}$, such that $(\mathcal{C}_{s+2-2/q,p,q})^{(5)}$ is satisfied and let $T \in (0, +\infty]$.*

For any

$$f \in L^q\left((0, T), \dot{B}_{p,q,\mathcal{H}}^{s,\sigma}(\mathbb{R}_+^n, \Lambda)\right), u_0 \in \dot{B}_{p,q,\mathcal{H}}^{2+s-\frac{2}{q},\sigma}(\mathbb{R}_+^n, \Lambda),$$

there exists a unique mild solution

$$u \in C_b^0\left([0, T], \dot{B}_{p,q,\mathcal{H}}^{2+s-\frac{2}{q},\sigma}(\mathbb{R}_+^n, \Lambda)\right)$$

⁽⁵⁾The condition (\mathcal{C}_{\dots}) is here to ensure the completeness of considered function spaces. See $(\mathcal{C}_{s,p,q})$ below.

to

$$(HSS) \quad (6) \quad \left\{ \begin{array}{ll} \partial_t u - \Delta u = f, & \text{on } (0, T) \times \mathbb{R}_+^n, \\ d^* u = 0, & \text{on } (0, T) \times \mathbb{R}_+^n, \\ \nu \lrcorner du|_{\partial \mathbb{R}_+^n} = 0, & \text{on } (0, T) \times \partial \mathbb{R}_+^n, \\ \nu \lrcorner u|_{\partial \mathbb{R}_+^n} = 0, & \text{on } (0, T) \times \partial \mathbb{R}_+^n, \\ u(0) = u_0, & \text{in } \dot{B}_{p,q}^{2+s-\frac{2}{q}}(\mathbb{R}_+^n, \Lambda), \end{array} \right.$$

with the estimate

$$\begin{aligned} \|u\|_{L^\infty([0,T], \dot{B}_{p,q}^{2+s-\frac{2}{q}}(\mathbb{R}_+^n))} + \|(\partial_t u, \nabla^2 u)\|_{L^q((0,T), \dot{B}_{p,q}^s(\mathbb{R}_+^n))} \\ \lesssim_{p,q,s,n} \|f\|_{L^q((0,T), \dot{B}_{p,q}^s(\mathbb{R}_+^n))} + \|u_0\|_{\dot{B}_{p,q}^{2+s-\frac{2}{q}}(\mathbb{R}_+^n)}. \end{aligned}$$

In the case $q = +\infty$, if we assume in addition $u_0 \in \dot{D}_p^s(\mathbb{A}_{\mathcal{H}}^2)$, we have

$$\|(\partial_t u, \nabla^2 u)\|_{L^\infty([0,T], \dot{B}_{p,\infty}^s(\mathbb{R}_+^n))} \lesssim_{p,s,n} \|f\|_{L^\infty((0,T), \dot{B}_{p,\infty}^s(\mathbb{R}_+^n))} + \|\mathbb{A}_{\mathcal{H}} u_0\|_{\dot{B}_{p,\infty}^s(\mathbb{R}_+^n)}.$$

1.2. Notations, definitions, and usual concepts

Throughout this paper the dimension will be $n \geq 2$, and \mathbb{N} will be the set of non-negative integers. For $a, b \in \mathbb{R}$ with $a \leq b$, we write $[[a, b]] := [a, b] \cap \mathbb{Z}$.

For two real numbers $A, B \in \mathbb{R}$, $A \lesssim_{a,b,c} B$ means that there exists a constant $C > 0$ depending on a, b, c such that $A \leq CB$. When both $A \lesssim_{a,b,c} B$ and $B \lesssim_{a,b,c} A$ are true, we simply write $A \sim_{a,b,c} B$. When the number of indices is overloaded, we allow ourselves to write $A \lesssim_{a,b,c}^{d,e,f} B$ instead of $A \lesssim_{a,b,c,d,e,f} B$.

1.2.1. Smooth and measurable functions

Denote by $\mathcal{S}(\mathbb{R}^n, \mathbb{C})$ the space of complex valued Schwartz function, and $\mathcal{S}'(\mathbb{R}^n, \mathbb{C})$ its dual called the space of tempered distributions. The Fourier transform on $\mathcal{S}'(\mathbb{R}^n, \mathbb{C})$ is written \mathcal{F} , and is pointwise defined for any $f \in L^1(\mathbb{R}^n, \mathbb{C})$ by

$$\mathcal{F}f(\xi) := \int_{\mathbb{R}^n} f(x) e^{-ix \cdot \xi} dx, \xi \in \mathbb{R}^n.$$

Additionally, for $p \in [1, +\infty]$, we will write $p' = \frac{p}{p-1}$ its Hölder conjugate.

For any $m \in \mathbb{N}$, the map $\nabla^m : \mathcal{S}'(\mathbb{R}^n, \mathbb{C}) \rightarrow \mathcal{S}'(\mathbb{R}^n, \mathbb{C}^{n^m})$ is defined as $\nabla^m u := (\partial^\alpha u)_{|\alpha|=m}$. We denote by

$$\left(e^{t\Delta} \right)_{t \geq 0} \quad \text{and} \quad \left(e^{-t(-\Delta)^{\frac{1}{2}}} \right)_{t \geq 0}$$

respectively the heat and Poisson semigroup on \mathbb{R}^n . We also introduce operators ∇' and Δ' which are respectively the gradient and the Laplacian on \mathbb{R}^{n-1} identified with the $n - 1$ first variables of \mathbb{R}^n , i.e. $\nabla' = (\partial_{x_1}, \dots, \partial_{x_{n-1}})$ and $\Delta' = \partial_{x_1}^2 + \dots + \partial_{x_{n-1}}^2$.

⁽⁶⁾ For introductory purpose, the notations here are either not precise enough or quite redundant. For instance, the condition $d^* u = 0$ already implies the boundary condition $\nu \lrcorner u|_{\partial \mathbb{R}_+^n} = 0$.

When Ω is an open set of \mathbb{R}^n , $C_c^\infty(\Omega, \mathbb{C})$ is the set of smooth compactly supported function in Ω , and $\mathcal{D}'(\Omega, \mathbb{C})$ is its topological dual. For $p \in [1, +\infty)$, $L^p(\Omega, \mathbb{C})$ is the normed vector space of complex valued (Lebesgue-) measurable functions whose p^{th} power is integrable with respect to the Lebesgue measure, $\mathcal{S}(\overline{\Omega}, \mathbb{C})$ (*resp.* $\mathcal{S}_0(\overline{\Omega}, \mathbb{C})$, $C_c^\infty(\overline{\Omega}, \mathbb{C})$) stands for functions which are restrictions on Ω of elements of $\mathcal{S}(\mathbb{R}^n, \mathbb{C})$ (*resp.* $\mathcal{S}_0(\mathbb{R}^n, \mathbb{C})$, $C_c^\infty(\mathbb{R}^n, \mathbb{C})$). Unless the contrary is explicitly stated, we will always identify $L^p(\Omega, \mathbb{C})$ (*resp.* $C_c^\infty(\Omega, \mathbb{C})$) as the subspace of functions in $L^p(\mathbb{R}^n, \mathbb{C})$ (*resp.* $C_c^\infty(\mathbb{R}^n, \mathbb{C})$) supported in $\overline{\Omega}$ (*resp.* in Ω) through the extension by 0 outside Ω . $L^\infty(\Omega, \mathbb{C})$ stands for the space of essentially bounded (Lebesgue-) measurable functions.

For $s \in \mathbb{R}$, $p \in [1, +\infty)$, $\ell_s^p(\mathbb{Z}, \mathbb{C})$, stands for the normed vector space of p -summable sequences of complex numbers with respect to the counting measure $2^{ksp}dk$; $\ell_s^\infty(\mathbb{Z}, \mathbb{C})$ stands for sequences $(x_k)_{k \in \mathbb{Z}}$ such that $(2^{ks}x_k)_{k \in \mathbb{Z}}$ is bounded. More generally, when X is a Banach space, for $p \in [1, +\infty]$, one may also consider $L^p(\Omega, X)$ which stands for the space of (Bochner-)measurable functions $u : \Omega \rightarrow X$, such that $t \mapsto \|u(t)\|_X \in L^p(\Omega, \mathbb{R})$, similarly one may consider $\ell_s^p(\mathbb{Z}, X)$.

1.2.2. (Bi)Sectorial operators on Banach spaces

We introduce the following subsets of the complex plane

$$\Sigma_\mu := \{ z \in \mathbb{C}^* : |\arg(z)| < \mu \}, \text{ if } \mu \in (0, \pi),$$

$$S_\mu := (-\Sigma_\mu) \cup \Sigma_\mu, \text{ if } \mu \in \left(0, \frac{\pi}{2}\right),$$

we also define $\Sigma_0 := (0, +\infty)$, $S_0 := \mathbb{R} \setminus \{0\}$, and later we are going to consider $\overline{\Sigma}_\mu$, and \overline{S}_μ their closure.

An operator $(D(A), A)$ on a Banach space X , over the field of complex numbers, is said to be ω -sectorial if for a fixed $\omega \in (0, \pi)$ both conditions are satisfied

- (1) $\sigma(A) \subset \overline{\Sigma}_\omega$, where $\sigma(A)$ stands for the spectrum of A ;
- (2) For all $\mu \in (\omega, \pi)$, $\sup_{\lambda \in \mathbb{C} \setminus \overline{\Sigma}_\mu} \|\lambda(\lambda I - A)^{-1}\|_{X \rightarrow X} < +\infty$.

Similarly, $(D(A), A)$ is said to be ω -bisectorial, for a fixed $\omega \in (0, \frac{\pi}{2})$, if $\sigma(A) \subset \overline{S}_\omega$, and for all $\mu \in (\omega, \frac{\pi}{2})$, $\sup_{\lambda \in \mathbb{C} \setminus \overline{S}_\mu} \|\lambda(\lambda I - A)^{-1}\|_{X \rightarrow X} < +\infty$.

The following two propositions are classical and well-known. It will be of paramount importance throughout the present work.

PROPOSITION 1.4 ([Haa06, Proposition 2.1.1]). — *Let $(D(A), A)$ be a sectorial operator on a Banach space X . Then the following assertions hold.*

- (1) *If $k \in \mathbb{N}$, and $x \in \overline{D(A)}$, then*

$$\lim_{t \rightarrow +\infty} t^k(tI + A)^{-k}x = x \text{ and } \lim_{t \rightarrow +\infty} A^k(tI + A)^{-k}x = 0.$$

- (2) *If $k \in \mathbb{N}$, and $x \in \overline{R(A)}$, then*

$$\lim_{t \rightarrow 0} t^k(tI + A)^{-k}x = 0 \text{ and } \lim_{t \rightarrow 0} A^k(tI + A)^{-k}x = x.$$

In particular, $N(A) \cap \overline{R(A)} = \{0\}$, so that $X = \overline{R(A)}$ implies that A is injective.

- (3) For every $k \in \mathbb{N}$, $D(A^k) \cap R(A^k)$ is dense in $\overline{D(A)} \cap \overline{R(A)}$.
- (4) If X is reflexive, then A is densely defined and induces a topological decomposition

$$X = N(A) \oplus \overline{R(A)}.$$

The presentation of the preceding statement follows [Ege15, Proposition 3.2.2], where a similar assertion is made for bisectorial operators, with suitable adjustments. For the next proposition, we refer to [Haa06, Proposition 3.4.1], one may also see [ABHN11, Proposition 3.7.2, Theorem 3.7.11].

PROPOSITION 1.5. — *Let $(D(A), A)$ be an ω -sectorial operator on a Banach space X , with $\omega \in [0, \frac{\pi}{2})$. There exists a unique holomorphic family of operators $(T(z))_{z \in \Sigma_{\frac{\pi}{2}-\omega} \cup \{0\}}$ such that the following assertions hold:*

- (1) For all $\phi \in (\omega, \frac{\pi}{2})$, $(T(z))_{z \in \Sigma_{\frac{\pi}{2}-\phi} \cup \{0\}}$ is a family of uniformly bounded operators such that

$$T(0) = I \quad \text{and} \quad T(z + z') = T(z)T(z') \quad \text{for all } z, z' \in \Sigma_{\frac{\pi}{2}-\omega} \cup \{0\}.$$

- (2) If $x \in \overline{D(A)} \cap \overline{R(A)}$, then for each $\psi \in (0, \frac{\pi}{2} - \omega)$,

$$T(z)x \xrightarrow[\substack{|z| \rightarrow 0 \\ |\arg(z)| \leq \psi}]{\hspace{1.5cm}} x \quad \text{and} \quad T(z)x \xrightarrow[\substack{|z| \rightarrow +\infty \\ |\arg(z)| \leq \psi}]{\hspace{1.5cm}} 0.$$

- (3) If $x \in X$, then $T(z)x \in D(A^n)$, for all $z \in \Sigma_{\frac{\pi}{2}-\omega}$, $n \in \mathbb{N}$, and the map $z \mapsto T(z)x$ is holomorphic on $\Sigma_{\frac{\pi}{2}-\omega}$ with derivatives of order $k \in \mathbb{N}$,

$$T^{(k)}(z)x = (-A)^k T(z)x \quad \text{for all } z \in \Sigma_{\frac{\pi}{2}-\omega}.$$

- (4) For $\alpha \in \mathbb{C}_+$, $z \in \Sigma_{\frac{\pi}{2}-\phi}$, with $\phi \in (\omega, \frac{\pi}{2})$, one has

$$\|A^\alpha T(z)\|_{X \rightarrow X} \lesssim_{A, \phi, \alpha} \frac{1}{|z|^{\Re \alpha}}.$$

Therefore, one writes $e^{-zA} := T(z)$ for all $z \in \Sigma_{\frac{\pi}{2}-\omega} \cup \{0\}$, and $(e^{-zA})_{z \in \Sigma_{\frac{\pi}{2}-\omega} \cup \{0\}}$ is called the (holomorphic) C_0 -semigroup with generator $-A$.

Provided $\mu \in (0, \pi)$, we denote by $\mathbf{H}^\infty(\Sigma_\mu)$, the set of bounded holomorphic functions on Σ_μ , the same goes with S_μ instead of Σ_μ , for $\mu \in [0, \frac{\pi}{2})$.

If $(D(A), A)$ is ω -(bi)sectorial with $\omega \in [0, \pi)$ (resp. $[0, \frac{\pi}{2})$), for $\mu \in (\omega, \pi)$ (resp. $(\omega, \frac{\pi}{2})$) we say that A admits a *bounded* (or $\mathbf{H}^\infty(\Sigma_\mu)$ - (resp. $\mathbf{H}^\infty(S_\mu)$ -)) *holomorphic functional calculus* on X (of angle μ), if for $\theta \in (\omega, \mu)$, there exists a constant K_θ , such that for all $f \in \mathbf{H}^\infty(\Sigma_\theta)$ (resp. $\mathbf{H}^\infty(S_\theta)$), we have that

$$\|f(A)\|_{X \rightarrow X} \leq K_\theta \|f\|_{L^\infty}.$$

For all $x \in D(A) \cap R(A)$, $f(A)x$ is defined by the following convergent integral,

$$(1.13) \quad f(A)x = \frac{1}{2i\pi} \int_{\partial \Sigma_\theta} f(z)(zI - A)^{-1}x \, dz,$$

and the same goes with ∂S_θ instead of $\partial \Sigma_\theta$ for the bisectorial case, both boundaries being oriented counterclockwise.

We also say that A has *bounded imaginary powers* (BIP) of type θ_A if $f(z) = z^{is}$ plugged in (1.13) yields a bounded linear operator for all $s \in \mathbb{R}$, and

$$\theta_A := \inf \left\{ \omega \geq 0 \mid \sup_{s \in \mathbb{R}} e^{-\omega|s|} \|A^{is}\|_{X \rightarrow X} < +\infty \right\}.$$

Notice that if A has a bounded holomorphic functional calculus then it has bounded imaginary powers.

The functional calculus of sectorial operators is widely reviewed in several references but we mention here Haase’s book [Haa06]. However, there are only few references known to the author that deal with a systematic treatment of bisectorial operators, Duelli and Weis as well as Egert did such a treatment see [DW05] and [Ege15, Chapter 3].

1.2.3. Interpolation of normed vector spaces

Let $(X, \|\cdot\|_X)$ and $(Y, \|\cdot\|_Y)$ be two normed vector spaces. We write $X \hookrightarrow Y$ to say that X embeds continuously in Y . Now let us recall briefly basics of interpolation theory. If there exists a Hausdorff topological vector space Z , such that $X, Y \subset Z$, then $X \cap Y$ and $X + Y$ are normed vector spaces with their canonical norms, and one can define the K -functional of $z \in X + Y$, for any $t > 0$ by

$$K(t, z, X, Y) := \inf_{\substack{(x,y) \in X \times Y, \\ z=x+y}} (\|x\|_X + t \|y\|_Y).$$

This allows us to construct, for any $\theta \in (0, 1)$, $q \in [1, +\infty]$, the real interpolation spaces between X and Y with indexes θ, q as

$$(X, Y)_{\theta, q} := \left\{ x \in X + Y \mid t \mapsto t^{-\theta} K(t, x, X, Y) \in L_*^q(\mathbb{R}_+) \right\},$$

where $L_*^q(\mathbb{R}_+) := L^q((0, +\infty), dt/t)$.

The interested reader could check [BL76, Lun18, Tri78] for more information about interpolation theory and its applications.

1.2.4. Sobolev and Besov spaces on \mathbb{R}^n

To deal with Sobolev and Besov spaces on the whole space, we need to introduce Littlewood–Paley decomposition given by $\phi \in C_c^\infty(\mathbb{R}^n)$, radial, real-valued, non-negative, such that

- $\text{supp } \phi \subset B(0, 4/3)$;
- $\phi|_{B(0, 3/4)} = 1$;

so we define the following functions for any $j \in \mathbb{Z}$ for all $\xi \in \mathbb{R}^n$,

$$\phi_j(\xi) := \phi(2^{-j}\xi), \quad \psi_j(\xi) := \phi_j(\xi/2) - \phi_j(\xi),$$

and the family $(\psi_j)_{j \in \mathbb{Z}}$ has the following properties

- $\text{supp } (\psi_j) \subset \{ \xi \in \mathbb{R}^n \mid 3 \cdot 2^{j-2} \leq |\xi| \leq 2^{j+3}/3 \}$;
- $\forall \xi \in \mathbb{R}^n \setminus \{0\}, \sum_{j=-M}^N \psi_j(\xi) \xrightarrow{N, M \rightarrow +\infty} 1$.

Such a family $(\phi, (\psi_j)_{j \in \mathbb{Z}})$ is called a Littlewood–Paley family. Now, we consider the two following families of operators associated with their Fourier multipliers:

- The *homogeneous* family of Littlewood–Paley dyadic decomposition operators $(\dot{\Delta}_j)_{j \in \mathbb{Z}}$, where

$$\dot{\Delta}_j := \mathcal{F}^{-1} \psi_j \mathcal{F},$$

- The *inhomogeneous* family of Littlewood–Paley dyadic decomposition operators $(\Delta_k)_{k \in \mathbb{Z}}$, where

$$\Delta_{-1} := \mathcal{F}^{-1} \phi \mathcal{F},$$

$$\Delta_k := \dot{\Delta}_k \text{ for any } k \geq 0, \text{ and } \Delta_k := 0 \text{ for any } k \leq -2.$$

One may notice, as a direct application of Young’s inequality for the convolution, that they are all uniformly bounded families of operators on $L^p(\mathbb{R}^n)$, $p \in [1, +\infty]$.

Both families of operators lead for $s \in \mathbb{R}$, $p, q \in [1, +\infty]$, $u \in \mathcal{S}'(\mathbb{R}^n)$, to the following quantities,

$$\begin{aligned} \|u\|_{\dot{B}_{p,q}^s(\mathbb{R}^n)} &= \left\| \left(2^{ks} \|\Delta_k u\|_{L^p(\mathbb{R}^n)} \right)_{k \in \mathbb{Z}} \right\|_{\ell^q(\mathbb{Z})} \\ \text{and } \|u\|_{\dot{B}_{p,q}^s(\mathbb{R}^n)} &= \left\| \left(2^{js} \|\dot{\Delta}_j u\|_{L^p(\mathbb{R}^n)} \right)_{j \in \mathbb{Z}} \right\|_{\ell^q(\mathbb{Z})}, \end{aligned}$$

respectively named the inhomogeneous and homogeneous Besov norms, but the homogeneous norm is not really a norm since $\|u\|_{\dot{B}_{p,q}^s(\mathbb{R}^n)} = 0$ does not imply that $u = 0$. Thus, following [BCD11, Chapter 2] and [DHMT21, Chapter 3], we introduce a subspace of tempered distributions such that $\|\cdot\|_{\dot{B}_{p,q}^s(\mathbb{R}^n)}$ is point-separating, say

$$\mathcal{S}'_h(\mathbb{R}^n) := \left\{ u \in \mathcal{S}'(\mathbb{R}^n) \mid \forall \Theta \in C_c^\infty(\mathbb{R}^n), \|\Theta(\lambda \mathfrak{D})u\|_{L^\infty(\mathbb{R}^n)} \xrightarrow{\lambda \rightarrow +\infty} 0 \right\},$$

where for $\lambda > 0$, $\Theta(\lambda \mathfrak{D})u = \mathcal{F}^{-1} \Theta(\lambda \cdot) \mathcal{F} u$. Notice that $\mathcal{S}'_h(\mathbb{R}^n)$ does not contain any non-zero polynomials, and for any $p \in [1, +\infty)$, $L^p(\mathbb{R}^n) \subset \mathcal{S}'_h(\mathbb{R}^n)$.

One can also define the following quantities called the inhomogeneous and homogeneous Sobolev spaces’ potential norms

$$\|u\|_{H^{s,p}(\mathbb{R}^n)} := \left\| (I - \Delta)^{\frac{s}{2}} u \right\|_{L^p(\mathbb{R}^n)} \quad \text{and} \quad \|u\|_{\dot{H}^{s,p}(\mathbb{R}^n)} := \left\| \sum_{j \in \mathbb{Z}} (-\Delta)^{\frac{s}{2}} \dot{\Delta}_j u \right\|_{L^p(\mathbb{R}^n)},$$

where $(-\Delta)^{\frac{s}{2}}$ is understood on $u \in \mathcal{S}'_h(\mathbb{R}^n)$ by the action on its dyadic decomposition, i.e.

$$(-\Delta)^{\frac{s}{2}} \dot{\Delta}_j u := \mathcal{F}^{-1} |\xi|^s \mathcal{F} \dot{\Delta}_j u,$$

which gives a family of C^∞ functions with at most polynomial growth.

Hence for any $p, q \in [1, +\infty]$, $s \in \mathbb{R}$, we define

- the inhomogeneous and homogeneous Sobolev (Bessel and Riesz potential) spaces,

$$\begin{aligned} H^{s,p}(\mathbb{R}^n) &= \left\{ u \in \mathcal{S}'(\mathbb{R}^n) \mid \|u\|_{H^{s,p}(\mathbb{R}^n)} < +\infty \right\}, \\ \dot{H}^{s,p}(\mathbb{R}^n) &= \left\{ u \in \mathcal{S}'_h(\mathbb{R}^n) \mid \|u\|_{\dot{H}^{s,p}(\mathbb{R}^n)} < +\infty \right\}; \end{aligned}$$

- and the inhomogeneous and homogeneous Besov spaces,

$$B_{p,q}^s(\mathbb{R}^n) = \left\{ u \in \mathcal{S}'(\mathbb{R}^n) \mid \|u\|_{B_{p,q}^s(\mathbb{R}^n)} < +\infty \right\},$$

$$\dot{B}_{p,q}^s(\mathbb{R}^n) = \left\{ u \in \mathcal{S}'_h(\mathbb{R}^n) \mid \|u\|_{\dot{B}_{p,q}^s(\mathbb{R}^n)} < +\infty \right\};$$

which are all normed vector spaces.

The treatment of homogeneous Besov spaces $\dot{B}_{p,q}^s(\mathbb{R}^n)$, $s \in \mathbb{R}$, $p, q \in [1, +\infty]$, defined on $\mathcal{S}'_h(\mathbb{R}^n)$ has been done in an extensive manner in [BCD11, Chapter 2]. The corresponding construction for homogeneous Sobolev spaces $\dot{H}^{s,p}(\mathbb{R}^n)$, $s \in \mathbb{R}$, $p \in (1, +\infty)$ has been achieved after wise, see [BCD11, Chapter 1] only for the case $p = 2$, [DHMT21, Chapter 3] only for the case $s \in \mathbb{N}$, and [Gau24b, Section 2] for the case $s \in \mathbb{R}$.

The following subspace of Schwartz functions, say

$$\mathcal{S}_0(\mathbb{R}^n) := \{ u \in \mathcal{S}(\mathbb{R}^n) \mid 0 \notin \text{supp}(\mathcal{F}f) \},$$

is a nice dense subspace in $L^p(\mathbb{R}^n)$, $H^{s,p}(\mathbb{R}^n)$, $\dot{H}^{s,p}(\mathbb{R}^n)$, $B_{p,q}^s(\mathbb{R}^n)$ and $\dot{B}_{p,q}^s(\mathbb{R}^n)$, for all $p \in (1, +\infty)$, $q \in [1, +\infty)$, $s \in \mathbb{R}$

The inhomogeneous spaces $L^p(\mathbb{R}^n)$, $H^{s,p}(\mathbb{R}^n)$, and $B_{p,q}^s(\mathbb{R}^n)$ are all complete for all $p, q \in [1, +\infty]$, $s \in \mathbb{R}$, but in this setting homogeneous spaces are no longer always complete (see [BCD11, Proposition 1.34, Remark 2.26]). Indeed, it can be shown (see [BCD11, Theorem 2.25]) that homogeneous Besov spaces $\dot{B}_{p,q}^s(\mathbb{R}^n)$ are complete whenever $(s, p, q) \in \mathbb{R} \times (1, +\infty) \times [1, +\infty]$ satisfies

$$(C_{s,p,q}) \quad \left[s < \frac{n}{p} \right] \text{ or } \left[q = 1 \text{ and } s \leq \frac{n}{p} \right],$$

From now, and until the end of this paper, we write $(C_{s,p})$ for the statement $(C_{s,p,p})$. One may show that, similarly, $\dot{H}^{s,p}(\mathbb{R}^n)$ is complete whenever $(C_{s,p})$ is satisfied, see [Gau24b, Proposition 2.2].

We recall that all $s > 0$, $(p, q) \in (1, +\infty) \times [1, +\infty]$, we have $L^p(\mathbb{R}^n) \cap \dot{H}^{s,p}(\mathbb{R}^n) = H^{s,p}(\mathbb{R}^n)$, and $L^p(\mathbb{R}^n) \cap \dot{B}_{p,q}^s(\mathbb{R}^n) = B_{p,q}^s(\mathbb{R}^n)$ with equivalent norms, see [BL76, Theorem 6.3.2] for more details.

According to [BL76, Section 6.4], for all $s \in \mathbb{R}$, $p, q \in (1, +\infty) \times [1, +\infty]$, $H^{s,p}(\mathbb{R}^n)$ and $B_{p,q}^s(\mathbb{R}^n)$ are both complete, and moreover, they are reflexive when $q \neq 1, +\infty$, and we have

$$(1.14) \quad (H^{s,p}(\mathbb{R}^n))' = H^{-s,p'}(\mathbb{R}^n), \quad (B_{p,q}^s(\mathbb{R}^n))' = B_{p',q'}^{-s}(\mathbb{R}^n),$$

$$(1.15) \quad (B_{p,\infty}^s(\mathbb{R}^n))' = B_{p',1}^{-s}(\mathbb{R}^n), \quad (B_{p,1}^s(\mathbb{R}^n))' = B_{p',\infty}^{-s}(\mathbb{R}^n).$$

We recall also the usual real interpolation identities,

$$(H^{s_0,p}(\mathbb{R}^n), H^{s_1,p}(\mathbb{R}^n))_{\theta,q} = B_{p,q}^s(\mathbb{R}^n), \quad (B_{p,q_0}^{s_0}(\mathbb{R}^n), B_{p,q_1}^{s_1}(\mathbb{R}^n))_{\theta,q} = B_{p,q}^s(\mathbb{R}^n),$$

whenever $(p, q_0, q_1, q) \in [1, +\infty] \times [1, +\infty]^3$ ($p \neq 1, +\infty$, when dealing with Sobolev (Bessel potential) spaces), $\theta \in (0, 1)$, $s_0 \neq s_1$ two real numbers, such that

$$s := (1 - \theta)s_0 + \theta s_1,$$

see [BL76, Theorem 6.4.5]. A similar statement is available for homogeneous function spaces.

PROPOSITION 1.6 ([Gau24b, Theorem 2.6]). — *Let $(p, q, q_0, q_1) \in (1, +\infty) \times [1, +\infty]^3$, $s_0, s_1 \in \mathbb{R}$, such that $s_0 \neq s_1$, and set*

$$s := (1 - \theta)s_0 + \theta s_1.$$

Assuming $(\mathcal{C}_{s_0,p})$ (resp. $(\mathcal{C}_{s_0,p,q_0})$), we get the following

$$(1.16) \quad \left(\dot{H}^{s_0,p}(\mathbb{R}^n), \dot{H}^{s_1,p}(\mathbb{R}^n) \right)_{\theta,q} = \left(\dot{B}_{p,q_0}^{s_0}(\mathbb{R}^n), \dot{B}_{p,q_1}^{s_1}(\mathbb{R}^n) \right)_{\theta,q} = \dot{B}_{p,q}^s(\mathbb{R}^n).$$

If moreover, one consider $p_0, p_1 \in (1, +\infty)$, and assume that (\mathcal{C}_{s_0,p_0}) and (\mathcal{C}_{s_1,p_1}) are true then also is $(\mathcal{C}_{s,p_\theta})$ and

$$(1.17) \quad \left[\dot{H}^{s_0,p_0}(\mathbb{R}^n), \dot{H}^{s_1,p_1}(\mathbb{R}^n) \right]_{\theta} = \dot{H}^{s,p_\theta}(\mathbb{R}^n),$$

and if $(\mathcal{C}_{s_0,p_0,q_0})$ and $(\mathcal{C}_{s_1,p_1,q_1})$ are satisfied then $(\mathcal{C}_{s,p_\theta,q_\theta})$ is also satisfied with

$$(1.18) \quad \left[\dot{B}_{p_0,q_0}^{s_0}(\mathbb{R}^n), \dot{B}_{p_1,q_1}^{s_1}(\mathbb{R}^n) \right]_{\theta} = \dot{B}_{p_\theta,q_\theta}^s(\mathbb{R}^n),$$

where $\frac{1}{p_\theta} := \frac{1-\theta}{p_0} + \frac{\theta}{p_1}$ and q_θ defined similarly.

PROPOSITION 1.7 ([Gau24b, Proposition 2.9]). — *For all $p \in (1, +\infty)$, $q \in [1, +\infty]$, for all $s \in (-1 + \frac{1}{p}, \frac{1}{p})$, for all $u \in \dot{H}^{s,p}(\mathbb{R}^n)$ (resp. $\dot{B}_{p,q}^s(\mathbb{R}^n)$),*

$$\left\| \mathbb{1}_{\mathbb{R}_+^n} u \right\|_{\dot{H}^{s,p}(\mathbb{R}^n)} \lesssim_{s,p,n} \|u\|_{\dot{H}^{s,p}(\mathbb{R}^n)} \quad \left(\text{resp. } \left\| \mathbb{1}_{\mathbb{R}_+^n} u \right\|_{\dot{B}_{p,q}^s(\mathbb{R}^n)} \lesssim_{s,p,n} \|u\|_{\dot{B}_{p,q}^s(\mathbb{R}^n)} \right).$$

The same results still holds with (H, B) instead of (\dot{H}, \dot{B}) .

1.2.5. Homogeneous Sobolev and Besov spaces on \mathbb{R}_+^n

Let $s \in \mathbb{R}$, $p \in (1, +\infty)$, $q \in [1, +\infty]$. Then for any $X \in \{B_{p,q}^s, \dot{B}_{p,q}^s, H^{s,p}, \dot{H}^{s,p}\}$, we define

$$X(\mathbb{R}_+^n) := X(\mathbb{R}^n)|_{\mathbb{R}_+^n},$$

with the usual quotient norm

$$\|u\|_{X(\mathbb{R}_+^n)} := \inf_{\substack{\tilde{u} \in X(\mathbb{R}^n), \\ \tilde{u}|_{\mathbb{R}_+^n} = u}} \|\tilde{u}\|_{X(\mathbb{R}^n)}.$$

A direct consequence of the definition of those spaces is the density of $\mathcal{S}_0(\overline{\mathbb{R}_+^n}) \subset \mathcal{S}(\overline{\mathbb{R}_+^n})$ in each of them, and also the completeness and reflexivity when their counterpart on \mathbb{R}^n also have the corresponding property. We can also define,

$$X_0(\mathbb{R}_+^n) := \left\{ u \in X(\mathbb{R}^n) \mid \text{supp } u \subset \overline{\mathbb{R}_+^n} \right\},$$

with natural induced norm $\|u\|_{X_0(\mathbb{R}_+^n)} := \|u\|_{X(\mathbb{R}^n)}$.

We have

- **density results** [Gau24b, Proposition 3.9, Corollary 3.12]:
for $p \in (1, +\infty)$, $q \in [1, +\infty)$, $s > -1 + \frac{1}{p}$, when $(\mathcal{C}_{s,p})$ is satisfied for Sobolev spaces and $(\mathcal{C}_{s,p,q})$ in the case of Besov spaces,

$$(1.19) \quad \dot{H}_0^{s,p}(\mathbb{R}_+^n) = \overline{C_c^\infty(\mathbb{R}_+^n)}^{\|\cdot\|_{\dot{H}^{s,p}(\mathbb{R}^n)}}, \quad \text{and} \quad \dot{B}_{p,q,0}^s(\mathbb{R}_+^n) = \overline{C_c^\infty(\mathbb{R}_+^n)}^{\|\cdot\|_{\dot{B}_{p,q}^s(\mathbb{R}^n)}};$$

- **duality results**, [Gau24b, Propositions 3.11 & 3.23]:
for all $p \in (1, +\infty)$, $q \in (1, +\infty]$, $s > -1 + \frac{1}{p}$, when $(\mathcal{C}_{s,p})$ is satisfied for Sobolev spaces and $(\mathcal{C}_{s,p,q})$ in the case of Besov spaces,

$$(1.20) \quad (\dot{H}^{s,p}(\mathbb{R}_+^n))' = \dot{H}_0^{-s,p'}(\mathbb{R}_+^n), \quad (\dot{B}_{p',q'}^{-s}(\mathbb{R}_+^n))' = \dot{B}_{p,q,0}^s(\mathbb{R}_+^n),$$

$$(1.21) \quad (\dot{H}_0^{s,p}(\mathbb{R}_+^n))' = \dot{H}^{-s,p'}(\mathbb{R}_+^n), \quad (\dot{B}_{p',q',0}^{-s}(\mathbb{R}_+^n))' = \dot{B}_{p,q}^s(\mathbb{R}_+^n).$$

- **intersections are well-defined and complete** [Gau24b, Lemma 2.5, Proposition 3.3, Proposition 3.20]:
for $p_0, p_1 \in (1, +\infty)$, $q_0, q_1 \in [1, +\infty]$, $s_j > -1 + \frac{1}{p_j}$, $j \in \{0, 1\}$, when (\mathcal{C}_{s_0,p_0}) is satisfied for Sobolev spaces, or $(\mathcal{C}_{s_0,p_0,q_0})$ in the case of Besov spaces, we have that

$$(1.22) \quad \dot{H}^{s_0,p_0}(\mathbb{R}_+^n) \cap \dot{H}^{s_1,p_1}(\mathbb{R}_+^n) = [\dot{H}^{s_0,p_0} \cap \dot{H}^{s_1,p_1}](\mathbb{R}_+^n),$$

$$(1.23) \quad \dot{B}_{p_0,q_0}^{s_0}(\mathbb{R}_+^n) \cap \dot{B}_{p_1,q_1}^{s_1}(\mathbb{R}_+^n) = [\dot{B}_{p_0,q_0}^{s_0} \cap \dot{B}_{p_1,q_1}^{s_1}](\mathbb{R}_+^n);$$

so that each space is complete. Moreover, it admits $\mathcal{S}_0(\overline{\mathbb{R}_+^n})$ as dense subspace whenever $q_0, q_1 < +\infty$.

- **interpolation results**, [Gau24b, Propositions 3.17 & 3.22]:
if $(\dot{h}, \dot{b}) \in \{(\dot{H}, \dot{B}), (\dot{H}_0, \dot{B}_{\cdot, \cdot, 0})\}$, with $(p, q, q_0, q_1) \in (1, +\infty) \times [1, +\infty]^3$ ($p, p_j \neq 1, +\infty$ is assumed, when dealing with Sobolev (Riesz potential) spaces), $\theta \in (0, 1)$, $s_j, s > -1 + 1/p_j$, $j \in \{0, 1\}$, with $s > -1 + 1/p$, where s_0, s_1, s are three real numbers, with

$$s = (1 - \theta)s_0 + \theta s_1,$$

such that $(\mathcal{C}_{s,p,q})$ is satisfied. Then, one has

$$(1.24) \quad \begin{aligned} (\dot{h}^{s_0,p}(\mathbb{R}_+^n), \dot{h}^{s_1,p}(\mathbb{R}_+^n))_{\theta,q} &= \dot{b}_{p,q}^s(\mathbb{R}_+^n), \\ (\dot{b}_{p,q_0}^{s_0}(\mathbb{R}_+^n), \dot{b}_{p,q_1}^{s_1}(\mathbb{R}_+^n))_{\theta,q} &= \dot{b}_{p,q}^s(\mathbb{R}_+^n). \end{aligned}$$

Note that, due to Proposition 1.7, we have for free the following equalities of homogeneous Sobolev and Besov spaces, with equivalent norms, for all $p \in (1, +\infty)$, $q \in [1, +\infty]$, $s \in (-1 + \frac{1}{p}, \frac{1}{p})$,

$$(1.25) \quad \dot{H}^{s,p}(\mathbb{R}_+^n) = \dot{H}_0^{s,p}(\mathbb{R}_+^n), \quad \dot{B}_{p,q}^s(\mathbb{R}_+^n) = \dot{B}_{p,q,0}^s(\mathbb{R}_+^n).$$

We also provide an additional interpolation inequality.

LEMMA 1.8. — *Let $1 < p_0, p_1 < +\infty$, $s_0, s_1 \in \mathbb{R}$, such that $s_j > -1 + 1/p_j$, $j \in \{0, 1\}$, and (\mathcal{C}_{s_0,p_0}) is satisfied. For $\theta \in (0, 1)$, we set $s := (1 - \theta)s_0 + \theta s_1$, $1/p := (1 - \theta)/p_0 + \theta/p_1$.*

Then, for all $u \in \dot{H}^{s_0,p_0}(\mathbb{R}_+^n) \cap \dot{H}^{s_1,p_1}(\mathbb{R}_+^n)$, we have $u \in \dot{H}^{s,p}(\mathbb{R}_+^n)$, with the estimate

$$\|u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} \lesssim_{s,p,n,\theta} \|u\|_{\dot{H}^{s_0,p_0}(\mathbb{R}_+^n)}^{(1-\theta)} \|u\|_{\dot{H}^{s_1,p_1}(\mathbb{R}_+^n)}^\theta.$$

A similar result holds for Besov spaces, replacing $(\dot{H}^{s_0,p_0}, \dot{H}^{s_1,p_1})$ and the condition (\mathcal{C}_{s_0,p_0}) by $(\dot{B}_{p_0,q_0}^{s_0}, \dot{B}_{p_1,q_1}^{s_1})$ and the condition $(\mathcal{C}_{s_0,p_0,q_0})$.

Proof. — Let $u \in \dot{H}^{s_0,p_0}(\mathbb{R}_+^n) \cap \dot{H}^{s_1,p_1}(\mathbb{R}_+^n)$, then the extension operator E from [Gau24b, Corollary 3.4], is such that $Eu \in \dot{H}^{s_0,p_0}(\mathbb{R}^n) \cap \dot{H}^{s_1,p_1}(\mathbb{R}^n)$. Therefore, by [Gau24b, Proposition 2.2 (vii), Corollary 3.4] and the definition of function spaces by restriction, one obtains

$$\begin{aligned} \|u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} &\leq \|Eu\|_{\dot{H}^{s,p}(\mathbb{R}^n)} \lesssim_{s,p,n,\theta} \|Eu\|_{\dot{H}^{s_0,p_0}(\mathbb{R}^n)}^{(1-\theta)} \|Eu\|_{\dot{H}^{s_1,p_1}(\mathbb{R}^n)}^\theta \\ &\lesssim_{s,p,n,\theta} \|u\|_{\dot{H}^{s_0,p_0}(\mathbb{R}_+^n)}^{(1-\theta)} \|u\|_{\dot{H}^{s_1,p_1}(\mathbb{R}_+^n)}^\theta. \end{aligned}$$

This ends the proof. □

1.2.6. Operators on Sobolev and Besov spaces

We introduce domains for an operator A acting on Sobolev or Besov spaces, denoting

- $D_p^s(A)$ (resp. $\dot{D}_p^s(A)$) its domain on $H^{s,p}$ (resp. $\dot{H}^{s,p}$);
- $D_{p,q}^s(A)$ (resp. $\dot{D}_{p,q}^s(A)$) its domain on $B_{p,q}^s$ (resp. $\dot{B}_{p,q}^s$);
- $D_p(A) = D_p^0(A) = \dot{D}_p^0(A)$ its domain on L^p .

Similarly, $N_p^s(A)$, $N_{p,q}^s(A)$ will stand for its nullspace on $H^{s,p}$ and $B_{p,q}^s$, and range spaces will be given respectively by $R_p^s(A)$ and $R_{p,q}^s(A)$. We replace N and R by \dot{N} and \dot{R} for their corresponding sets on homogeneous function spaces.

If the operator A has different realizations depending on various function spaces and on the considered open set, we may write its domain $D(A, \Omega)$, and similarly for its nullspace N and range space R . We omit the open set Ω if there is no possible confusion.

Acknowledgments

The author would like to thank Sylvie Monniaux for her numerous feedbacks, which helped to improve the manuscript. The author would also like to thank Yoshikazu Giga for personal discussions that heavily inspired the last Section 3.3. The author is greatly indebted to the anonymous referees for their careful proofreading and useful suggestions that helped to substantially improve the manuscript.

2. Hodge Laplacians, Hodge decompositions and Hodge–Stokes operators

This section is dedicated to the study of Hodge Laplacians, the Hodge decomposition and Hodge–Stokes operators, on Sobolev and Besov spaces on \mathbb{R}^n and \mathbb{R}_+^n .

We will first introduce here the formalism of differential forms in the Euclidean setting. Resolvent estimates for the Hodge Laplacian and Hodge–Stokes like operators on the whole space will follow from standard Fourier and Harmonic analysis, from which we will deduce the related Hodge decomposition on \mathbb{R}^n as well as the boundedness of holomorphic functional calculus for each operator.

Secondly, we are going to give all corresponding similar results for the Hodge Laplacians, the Hodge decomposition and Hodge–Stokes operators on the half-space \mathbb{R}_+^n . Those results are going to be built from what happens on the whole space \mathbb{R}^n , mimicking the behavior of Dirichlet and Neumann Laplacians on the half-space, see [Gau24b, Section 5].

2.1. Differential forms on Euclidean space, and corresponding function spaces

Here Ω stands for a domain of \mathbb{R}^n with at least, if not empty, Lipschitz boundary. The open set Ω will be specified later on to be either, the whole space \mathbb{R}^n or the half-space \mathbb{R}_+^n . Recall briefly that $\partial\mathbb{R}^n = \emptyset$, and $\partial\mathbb{R}_+^n = \mathbb{R}^{n-1} \times \{0\}$. We also recall that the outer normal unit at $\partial\mathbb{R}_+^n$ is $\nu = -\mathbf{e}_n$, where $(\mathbf{e}_k)_{k \in \llbracket 1, n \rrbracket}$ is the canonical basis of \mathbb{R}^n , identified with its dual basis denoted by $(dx_k)_{k \in \llbracket 1, n \rrbracket}$, where $dx_k(\mathbf{e}_j) = \mathbb{1}_{\{k\}}(j)$, $(k, j) \in \llbracket 1, n \rrbracket^2$.

Following [MM18, Mon21], we introduce the *exterior derivative* $d := \nabla \wedge = \sum_{k=1}^n \partial_{x_k} \mathbf{e}_k \wedge$ and the *interior derivative* (or *coderivative*) $\delta := -\nabla \lrcorner = -\sum_{k=1}^n \partial_{x_k} \mathbf{e}_k \lrcorner$ acting on differential forms on a domain $\Omega \subset \mathbb{R}^n$, i.e. acting on functions defined on Ω which take values in the complexified exterior algebra $\Lambda = \Lambda^0 \oplus \Lambda^1 \oplus \dots \oplus \Lambda^n$ of \mathbb{R}^n . We allow us a slight abuse of notation: here we will not distinguish vectors of \mathbb{R}^n , vector fields, and 1-differential forms.

We also recall that for $k \in \llbracket 0, n \rrbracket$, $u \in \Lambda^k$ can be uniquely determined by $(u_I)_{I \in \mathcal{I}_n^k} \in \mathbb{C}^{\binom{n}{k}}$ such that

$$u = \sum_{I \in \mathcal{I}_n^k} u_I dx_I,$$

where $\mathcal{I}_n^k = \{(\ell_j)_{j \in \llbracket 1, k \rrbracket} \in \llbracket 1, n \rrbracket^k \mid \ell_j < \ell_{j+1}\}$, with $\|\mathcal{I}_n^k\| = \binom{n}{k}$, and u_I and dx_I stands respectively for $u_{\ell_1 \ell_2 \dots \ell_k}$ and $dx_{\ell_1} \wedge dx_{\ell_2} \wedge \dots \wedge dx_{\ell_k}$ whenever $I = (\ell_j)_{j \in \llbracket 1, k \rrbracket}$.

One may also notice that such representation of k -differential forms with increasing index is possible due to symmetry properties (i.e. $dx_\ell \wedge dx_k = -dx_k \wedge dx_\ell$ for all $k, \ell \in \llbracket 1, n \rrbracket$).

In particular, remark that $\Lambda^0 \simeq \mathbb{C}$, the space of complex scalars, and more generally $\Lambda^k \simeq \mathbb{C}^{\binom{n}{k}}$, so that $\Lambda \simeq \mathbb{C}^{2^n}$. We also set $\Lambda^\ell = \{0\}$ if $\ell < 0$ or $\ell > n$.

On the exterior algebra Λ , the basic operations are

- (i) the exterior product $\wedge : \Lambda^k \times \Lambda^\ell \rightarrow \Lambda^{k+\ell}$,
- (ii) the interior product $\lrcorner : \Lambda^k \times \Lambda^\ell \rightarrow \Lambda^{\ell-k}$,
- (iii) the Hodge star operator $\star : \Lambda^\ell \rightarrow \Lambda^{n-\ell}$,
- (iv) the inner product $\langle \cdot, \cdot \rangle : \Lambda^\ell \times \Lambda^\ell \rightarrow \mathbb{C}$.

If $a \in \Lambda^1, u \in \Lambda^\ell$ and $v \in \Lambda^{\ell+1}$, then

$$\langle a \wedge u, v \rangle = \langle u, a \lrcorner v \rangle.$$

For more details, we refer to, e.g., [AM04, Section 2] and [CM10, Section 3], noting that both papers contain some historical background (and being careful that δ has the opposite sign in [AM04]). One may also consult [DC94] for an introduction from the euclidean setting point of view, and [Jos11, Section 1-3] for basic and usual properties in the more general Riemannian setting⁽⁷⁾. We recall the relation between d and δ via the Hodge star operator:

$$\star \delta u = (-1)^\ell d(\star u) \quad \text{and} \quad \star du = (-1)^{\ell-1} \delta(\star u) \quad \text{for an } \ell\text{-form } u.$$

In dimension $n = 3$, this gives (see [CM10]) for a vector $a \in \mathbb{R}^3$ identified with a 1-form

- u scalar, interpreted as 0 -form: $a \wedge u = ua, a \lrcorner u = 0$;
- u scalar, interpreted as 3 -form: $a \wedge u = 0, a \lrcorner u = ua$;
- u vector, interpreted as 1 -form: $a \wedge u = a \times u, a \lrcorner u = a \cdot u$;
- u vector, interpreted as 2 -form: $a \wedge u = a \cdot u, a \lrcorner u = -a \times u$.

From now and until the end of the present paper, if $p \in (1, +\infty), q \in [1, +\infty], s \in \mathbb{R}, k \in \llbracket 0, n \rrbracket$ and $X^s \in \{H^{s,p}, \dot{H}^{s,p}, B_{p,q}^s, \dot{B}_{p,q}^s\}$, then $X^s(\Omega, \Lambda^k)$ stands for k -differential forms whose coefficients lie in $X^s(\Omega)$, i.e. $X^s(\Omega, \Lambda^k) \simeq X^s(\Omega, \mathbb{C}^{\binom{n}{k}})$. One may also consider similarly $X_0^s(\Omega, \Lambda^k)$.

Operators d and δ are differential operators such that $d^2 = d \circ d = 0$ and $\delta^2 = \delta \circ \delta = 0$, and each of them are bounded seen as linear operators $X^s(\Omega, \Lambda) \rightarrow X^{s-1}(\Omega, \Lambda)$.

We recall the following integration by parts formula, for all $u, v \in \mathcal{S}(\overline{\Omega}, \Lambda)$,

$$(2.1) \quad \int_{\Omega} \langle du(x), v(x) \rangle dx = \int_{\Omega} \langle u(x), \delta v(x) \rangle dx + \int_{\partial\Omega} \langle u(x), \nu \lrcorner v(x) \rangle d\sigma_x,$$

$$(2.2) \quad \int_{\Omega} \langle \delta u(x), v(x) \rangle dx = \int_{\Omega} \langle u(x), dv(x) \rangle dx + \int_{\partial\Omega} \langle u(x), \nu \wedge v(x) \rangle d\sigma_x,$$

which are true since we are in the Euclidean setting and where ν is the outer unit normal identified as a 1-form. More generally, for all $T \in \mathcal{D}'(\Omega, \Lambda^k) \simeq \mathcal{D}'(\Omega, \mathbb{C}^{\binom{n}{k}}), k \in \llbracket 0, n \rrbracket$, we define

$$\begin{aligned} \langle dT, \phi \rangle_{\Omega} &:= \langle T, \delta\phi \rangle_{\Omega} \quad \text{for all } \phi \in C_c^\infty(\Omega, \Lambda^{k+1}), \\ \langle \delta T, \psi \rangle_{\Omega} &:= \langle T, d\psi \rangle_{\Omega} \quad \text{for all } \psi \in C_c^\infty(\Omega, \Lambda^{k-1}). \end{aligned}$$

⁽⁷⁾Notice that the Riemannian setting presented by Jost deals with compact manifold but a lot of computations remain true in their full generality, due to local behavior of each operation (Hodge star operator, exterior and interior products etc.)

In particular, one may see those operators as unbounded ones and introduce their respective domains on $L^p(\Omega, \Lambda^k)$, $k \in \llbracket 0, n \rrbracket$, denoted by $D_p(d, \Lambda^k)$ and $D_p(\delta, \Lambda^k)$ defined as

$$D_p(d, \Lambda^k) := \left\{ u \in L^p(\Omega, \Lambda^k) \mid du \in L^p(\Omega, \Lambda^{k+1}) \right\}$$

$$\text{and } D_p(\delta, \Lambda^k) := \left\{ u \in L^p(\Omega, \Lambda^k) \mid \delta u \in L^p(\Omega, \Lambda^{k-1}) \right\},$$

as well as their ranges

$$R_p(d, \Lambda^k) := \left\{ v \in L^p(\Omega, \Lambda^k) \mid v = du, u \in D_p(d, \Lambda^{k-1}) \right\}$$

$$\text{and } R_p(\delta, \Lambda^k) := \left\{ v \in L^p(\Omega, \Lambda^k) \mid v = \delta u, u \in D_p(d, \Lambda^{k+1}) \right\}.$$

We can introduce their corresponding counterparts on homogeneous Sobolev spaces scales, $\dot{D}_p^s(d, \Lambda^k)$ on $\dot{H}^{s,p}$, the same goes for inhomogeneous Sobolev spaces $D_p^s(d, \Lambda^k)$ on $H^{s,p}$. The same goes with the interior derivative δ instead of d . One may proceed in a similar fashion considering their domains on inhomogeneous and homogeneous Besov spaces.

As a sequence of (densely defined but not necessarily closed) unbounded operators, we get:

$$\begin{array}{ccccccccccc} d & : & X^s(\Omega, \Lambda^0) & \longrightarrow & X^s(\Omega, \Lambda^1) & \longrightarrow & X^s(\Omega, \Lambda^2) & \longrightarrow & \dots & \longrightarrow & X^s(\Omega, \Lambda^{n-1}) & \longrightarrow & X^s(\Omega, \Lambda^n) & \longrightarrow & 0 \\ 0 & \longleftarrow & X^s(\Omega, \Lambda^0) & \longleftarrow & X^s(\Omega, \Lambda^1) & \longleftarrow & X^s(\Omega, \Lambda^2) & \longleftarrow & \dots & \longleftarrow & X^s(\Omega, \Lambda^{n-1}) & \longleftarrow & X^s(\Omega, \Lambda^n) & : & \delta. \end{array}$$

In dimension $n = 3$, one can specialize, by means of the identification $\Lambda^k \simeq \mathbb{C}^{\binom{3}{k}}$, as

$$\begin{array}{ccccccccccc} d & : & X^s(\Omega, \mathbb{C}) & \xrightarrow{\nabla} & X^s(\Omega, \mathbb{C}^3) & \xrightarrow{\text{curl}} & X^s(\Omega, \mathbb{C}^3) & \xrightarrow{\text{div}} & X^s(\Omega, \mathbb{C}) & \longrightarrow & 0 \\ 0 & \longleftarrow & X^s(\Omega, \mathbb{C}) & \xleftarrow{\text{div}} & X^s(\Omega, \mathbb{C}^3) & \xleftarrow{\text{curl}} & X^s(\Omega, \mathbb{C}^3) & \xleftarrow{\nabla} & X^s(\Omega, \mathbb{C}) & : & \delta. \end{array}$$

Thus for arbitrary dimension n , the operator d restricted to its action on $X^s(\Omega, \Lambda^1)$, with value in $X^{s-1}(\Omega, \Lambda^2)$, and δ restricted to its action on $X^s(\Omega, \Lambda^{n-1})$, with value in $X^{s-1}(\Omega, \Lambda^{n-2})$, are fair consistent generalizations of the curl operator on \mathbb{R}^3 . Since in dimension n higher than 4, $n - 1 \neq 2$, we also have to distinguish their dual operators: the operator d restricted to its action on $X^s(\Omega, \Lambda^{n-2})$ and the operator δ restricted to its action on $X^s(\Omega, \Lambda^2)$ which are fair consistent generalizations of the dual operator ${}^t\text{curl}$ (usually fully identified with the curl operator) on \mathbb{R}^3 .

We can use (2.1) and (2.2) to consider adjoints of d and δ in the sense of maximal adjoint operators in the Hilbert space $L^2(\Omega, \Lambda)$, so that we will see later, e.g. Lemma 2.11, that they have the following exact description of their domains

$$D_2(d^*, \Lambda^k) = \left\{ u \in d_2(\delta, \Lambda^k) \mid \nu \lrcorner u|_{\partial\Omega} = 0 \right\}$$

$$\text{and } D_2(\delta^*, \Lambda^k) = \left\{ u \in d_2(d, \Lambda^k) \mid \nu \wedge u|_{\partial\Omega} = 0 \right\}.$$

One can also see those adjoint operators through the following L^2 -closures of unbounded operators,

$$(D_2(d^*, \Lambda^k), d^*) = \overline{(C_c^\infty(\Omega, \Lambda^k), \delta)} \quad \text{and} \quad (D_2(\delta^*, \Lambda^k), \delta^*) = \overline{(C_c^\infty(\Omega, \Lambda^k), d)}.$$

DEFINITION 2.1. —

- (i) The **Hodge–Dirac operator** on Ω with **tangential boundary conditions** is defined as

$$D_t := \delta^* + \delta.$$

Its square denoted by $-\Delta_{\mathcal{H},t} := D_t^2 = \delta^*\delta + \delta\delta^*$, is called the (negative) **Hodge Laplacian with relative boundary conditions** (also called **generalised Dirichlet boundary conditions**)

$$\nu \wedge u|_{\partial\Omega} = 0, \text{ and } \nu \wedge \delta u|_{\partial\Omega} = 0.$$

The restriction to scalar functions $u : \Omega \rightarrow \Lambda^0$ gives $-\Delta_{\mathcal{H},t}u = \delta\delta^*u = -\Delta_{\mathcal{D}}u$, where $-\Delta_{\mathcal{D}}$ is the Dirichlet Laplacian.

- (ii) The **Hodge–Dirac operator** on Ω with **normal boundary conditions** is defined as

$$D_n := d + d^*.$$

Its square denoted by $-\Delta_{\mathcal{H},n} := D_n^2 = dd^* + d^*d$, is called the (negative) **Hodge Laplacian with absolute boundary conditions** (also called **generalised Neumann boundary conditions**)

$$\nu \lrcorner u|_{\partial\Omega} = 0, \text{ and } \nu \lrcorner du|_{\partial\Omega} = 0.$$

The restriction to scalar functions $u : \Omega \rightarrow \Lambda^0$ gives $-\Delta_{\mathcal{H},n}u = d^*du = -\Delta_{\mathcal{N}}u$, where $-\Delta_{\mathcal{N}}$ is the Neumann Laplacian.

Notation 2.2. — When it does not matter $(\mathfrak{d}, D, -\Delta_{\mathcal{H}})$ will stand either for $(\delta, D_t, -\Delta_{\mathcal{H},t})$ or $(d, D_n, -\Delta_{\mathcal{H},n})$, just writing

$$-\Delta_{\mathcal{H}} = D^2 = \mathfrak{d}\mathfrak{d}^* + \mathfrak{d}^*\mathfrak{d}.$$

Remark 2.3. — We make three independent remarks:

- The exterior and interior derivatives, as well as the Hodge–Dirac operators are a priori unbounded operators only defined on the biggest space $L^2(\Omega, \Lambda)$. However, throughout this paper we will use some semantic distortion referring to Hodge–Dirac operators as “unbounded operators on $L^2(\Omega, \Lambda^k)$ ”, $k \in \llbracket 0, n \rrbracket$, by their natural restriction to differential forms of degree k , such as

$$\begin{aligned} D_n : D_2(D_n, \Lambda^k) &\longrightarrow L^2(\Omega, \Lambda^{k-1}) \oplus L^2(\Omega, \Lambda^{k+1}) \\ &= L^2(\Omega, \Lambda^{k-1} \oplus \Lambda^{k+1}) \subset L^2(\Omega, \Lambda), \end{aligned}$$

and similarly for D_t , even if the range is not a subset of $L^2(\Omega, \Lambda^k)$. This misuse will remain also for other function spaces that could replace L^2 . More generally, for $m \in \llbracket 0, n \rrbracket$, for $0 \leq k_0 < k_1 < \dots < k_m \leq n$, we always use the canonical identification

$$C_c^\infty(\Omega, \Lambda^{k_0} \oplus \dots \oplus \Lambda^{k_m}) \hookrightarrow C_c^\infty(\Omega, \Lambda) \text{ and } \mathcal{D}'(\Omega, \Lambda^{k_0} \oplus \dots \oplus \Lambda^{k_m}) \hookrightarrow \mathcal{D}'(\Omega, \Lambda)$$

with identically zero coefficients on remaining indices.

- We recall here that, if $\Omega \subset \mathbb{R}^3$ is an open set with, say at least, Lipschitz boundary, one has formally for u with value in $\Lambda^1 \simeq \mathbb{C}^3$ or $\Lambda^2 \simeq \mathbb{C}^3$,

$$-\Delta_{\mathcal{H},t}u = -\Delta_{\mathcal{H},n}u = \operatorname{curl} \operatorname{curl} u - \nabla \operatorname{div} u$$

with either one of the following couple of boundary conditions

$$\left[u \cdot \nu_{|\partial\Omega} = 0, \nu \times \operatorname{curl} u_{|\partial\Omega} = 0 \right] \text{ or } \left[u \times \nu_{|\partial\Omega} = 0, (\operatorname{div} u)\nu_{|\partial\Omega} = 0 \right].$$

- In the case of $\Omega = \mathbb{R}^n$, notice that no boundary value comes in, hence $d^* = \delta$, $\delta^* = d$, so that

$$D. = D_t = D_n = (d + \delta),$$

$$\text{and } -\Delta_{\mathcal{H}} = -\Delta_{\mathcal{H},t} = -\Delta_{\mathcal{H},n} = (d + \delta)^2 = d\delta + \delta d.$$

DEFINITION 2.4. —

- (i) The orthogonal projector defined on $L^2(\Omega, \Lambda^k)$ onto $N_2(d^*, \Lambda^k)$ is denoted by \mathbb{P} and called the **generalized Helmholtz–Leray (or Leray) projector**.
- (ii) The (bounded) orthogonal projector defined on $L^2(\Omega, \Lambda^k)$ onto $N_2(\delta, \Lambda^k)$ is denoted by \mathbb{Q} .
- (iii) For $p \in (1, +\infty)$, $s \in \mathbb{R}$ and $k \in \llbracket 0, n \rrbracket$, we say that $H^{s,p}(\Omega, \Lambda^k)$ admits a **Hodge decomposition** if $(D_p^s(d, \Lambda^k), d)$, $(D_p^s(\delta, \Lambda^k), \delta)$ and their respective adjoints are closable and

$$\begin{aligned} (\mathfrak{H}_p^s) \quad H^{s,p}(\Omega, \Lambda^k) &= N_p^s(\mathfrak{d}, \Lambda^k) \oplus \overline{R_p^s(\mathfrak{d}^*, \Lambda^k)}, \\ &= \overline{R_p^s(\mathfrak{d}, \Lambda^k)} \oplus N_p^s(\mathfrak{d}^*, \Lambda^k), \end{aligned}$$

holds in the topological sense. We keep the same definition of the Hodge decomposition on other function spaces replacing $(H^{s,p}, D_p^s, R_p^s, N_p^s)$ by either $(\dot{H}^{s,p}, \dot{D}_p^s, \dot{R}_p^s, \dot{N}_p^s)$, $(B_{p,q}^s, d_{p,q}^s, R_{p,q}^s, N_{p,q}^s)$, or even by $(\dot{B}_{p,q}^s, \dot{D}_{p,q}^s, \dot{R}_{p,q}^s, \dot{N}_{p,q}^s)$, with $q \in [1, +\infty]$.

One can notice that in the case of vector fields (identified with 1-forms, i.e. $L^2(\Omega, \Lambda^1) \simeq L^2(\Omega, \mathbb{C}^n)$), we can identify \mathbb{P} as the usual Helmholtz–Leray projector on divergence free vector fields with null normal trace at the boundary

$$\mathbb{P} : L^2(\Omega, \mathbb{C}^n) \longrightarrow L^2_\sigma(\Omega) = \left\{ u \in L^2(\Omega, \mathbb{C}^n) \mid \operatorname{div} u = 0, u \cdot \nu_{|\partial\Omega} = 0 \right\}.$$

It gives the following classical orthogonal, topological, Hodge decomposition, see [Soh01, Chapter 2, Section 2.5],

$$L^2(\Omega, \mathbb{C}^n) = L^2_\sigma(\Omega) \overset{\perp}{\oplus} \nabla \dot{H}^1(\Omega, \mathbb{C}),$$

for any sufficiently nice domains Ω , say for instance with uniform Lipschitz boundary, see [Soh01, Lemma 2.5.3].

Before investigating the Hodge decomposition and the functional analytic properties of the Hodge Laplacian on differential forms on function spaces in \mathbb{R}_+^n , we want to know a bit more about the whole space case. In the next subsection, devoted to the whole space, we gather well known facts and results which lack explicit references in the literature to the best of author’s knowledge.

2.2. The case of the whole space

On the whole space \mathbb{R}^n the action of the Laplacian and the Hodge decomposition for vector fields is well known in the literature on usual spaces as Lebesgue spaces $L^p(\mathbb{R}^n, \mathbb{C}^n)$, $p \in (1, +\infty)$, and so is the case of inhomogeneous and homogeneous Sobolev and Besov spaces. Our main goal here is to extend and summarize those results with the formalism of differential forms.

To do so, we introduce an extension of the Fourier transform to differential forms whose coefficients lie in the space of complex valued Schwartz functions $\mathcal{S}(\mathbb{R}^n, \mathbb{C})$, or in the space of tempered distribution $\mathcal{S}'(\mathbb{R}^n, \mathbb{C})$.

- For all $u \in L^1(\mathbb{R}^n, \Lambda^k) \simeq L^1(\mathbb{R}^n, \mathbb{C}^{\binom{n}{k}})$, $k \in \llbracket 0, n \rrbracket$, we define

$$\mathcal{F}u := \sum_{I \in \mathcal{I}_n^k} \mathcal{F}u_I d\xi_I \in C_0^0(\mathbb{R}^n, \Lambda^k).$$

Hence, as in the scalar valued case, the Fourier transform \mathcal{F} induces a topological automorphism of $\mathcal{S}(\mathbb{R}^n, \Lambda^k) \simeq \mathcal{S}(\mathbb{R}^n, \mathbb{C}^{\binom{n}{k}})$.

- For $k \in \llbracket 0, n \rrbracket$, we write $\mathcal{S}'(\mathbb{R}^n, \Lambda^k) := (\mathcal{S}(\mathbb{R}^n, \Lambda^k))' \simeq \mathcal{S}'(\mathbb{R}^n, \mathbb{C}^{\binom{n}{k}})$. Similarly, the Fourier transform \mathcal{F} is an automorphism of $\mathcal{S}'(\mathbb{R}^n, \Lambda^k)$.
- For all $T \in \mathcal{S}'(\mathbb{R}^n, \Lambda^k)$, $k \in \llbracket 0, n \rrbracket$, we define

$$\begin{aligned} \langle dT, \phi \rangle_{\mathbb{R}^n} &:= \langle T, \delta\phi \rangle_{\mathbb{R}^n} \text{ for all } \phi \in \mathcal{S}(\mathbb{R}^n, \Lambda^{k+1}), \\ \langle \delta T, \psi \rangle_{\mathbb{R}^n} &:= \langle T, d\psi \rangle_{\mathbb{R}^n} \text{ for all } \psi \in \mathcal{S}(\mathbb{R}^n, \Lambda^{k-1}). \end{aligned}$$

The following lemma is straightforward and fundamental for our analysis.

LEMMA 2.5. — For all $u \in \mathcal{S}(\mathbb{R}^n, \Lambda^k)$, $k \in \llbracket 0, n \rrbracket$, for all $\xi \in \mathbb{R}^n$,

$$\mathcal{F}[du](\xi) = i\xi \wedge \mathcal{F}u(\xi) \quad \text{and} \quad \mathcal{F}[\delta u](\xi) = -i\xi \lrcorner \mathcal{F}u(\xi).$$

Remark 2.6. — This is somewhat consistent, when $n = 3$, with formulas like

$$\mathcal{F}[\text{curl } u](\xi) = i\xi \times \mathcal{F}u(\xi) \quad \text{and} \quad \mathcal{F}[\text{div } u](\xi) = i\xi \cdot \mathcal{F}u(\xi), \quad u \in \mathcal{S}(\mathbb{R}^3, \mathbb{C}^3), \xi \in \mathbb{R}^3.$$

From there, the identity for all differential forms u of degree k , and all vector $v \in \mathbb{R}^n$,

$$v \wedge (v \lrcorner u) + v \lrcorner (v \wedge u) = |v|^2 u,$$

with the use of Lemma 2.5 yields with Remark 2.3 that, for all $u \in \mathcal{S}'(\mathbb{R}^n, \Lambda^k)$, $k \in \llbracket 0, n \rrbracket$,

$$\begin{aligned} \mathcal{F}[-\Delta_{\mathcal{H}}u](\xi) &= \mathcal{F}[(\delta d + d\delta)u](\xi) = i\xi \wedge (-i\xi \lrcorner u) + -i\xi \lrcorner (i\xi \wedge u) \\ &= |\xi|^2 \cdot \mathcal{F}u(\xi) = \mathcal{F}[-\Delta u](\xi). \end{aligned}$$

Hence, the Hodge Laplacian on the whole space is nothing but the scalar Laplacian applied separately to each component of a differential form so that its properties are carried over by the scalar Laplacian. We state then a very well known result adapted to our setting.

Before the statement, we recall that, for $1 < p, \tilde{p} < +\infty$, $1 \leq q, \tilde{q} \leq +\infty$, $s, \alpha \in \mathbb{R}$, the intersection spaces

$$[\dot{H}^{s,p} \cap \dot{H}^{\alpha,\tilde{p}}](\mathbb{R}^n, \Lambda), [\dot{B}_{p,q}^s \cap \dot{B}_{\tilde{p},\tilde{q}}^\alpha](\mathbb{R}^n, \Lambda) \subset \mathcal{S}'_h(\mathbb{R}^n, \Lambda)$$

are complete whenever $(\mathcal{C}_{s,p})$ is satisfied for Sobolev spaces, and when $(\mathcal{C}_{s,p,q})$ is satisfied for Besov spaces. See [Gau24b, Lemma 2.5] and [BCD11, Proposition 2.17].

THEOREM 2.7. — *Let $p, \tilde{p} \in (1, +\infty)$, $q, \tilde{q} \in [1, +\infty]$, $s, \alpha \in \mathbb{R}$, and $k \in \llbracket 0, n \rrbracket$. The Hodge Laplacian is an injective operator on $\mathcal{S}'_h(\mathbb{R}^n, \Lambda^k)$, and satisfies the following properties*

(i) *For $f \in \mathcal{S}'_h(\mathbb{R}^n, \Lambda^k)$, consider the problem*

$$-\Delta u = f \quad \text{in } \mathbb{R}^n.$$

(a) *If $f \in [\dot{H}^{s,p} \cap \dot{H}^{\alpha,\tilde{p}}](\mathbb{R}^n, \Lambda^k)$, and $(\mathcal{C}_{s+2,p})$ is satisfied, then there exists a unique solution $u \in [\dot{H}^{s+2,p} \cap \dot{H}^{\alpha+2,\tilde{p}}](\mathbb{R}^n, \Lambda^k)$ with the estimates,*

$$\|d\delta u\|_{\dot{H}^{s,p}(\mathbb{R}^n)} + \|\delta du\|_{\dot{H}^{s,p}(\mathbb{R}^n)} \lesssim_{p,n,s} \|\nabla^2 u\|_{\dot{H}^{s,p}(\mathbb{R}^n)} \lesssim_{p,n,s} \|f\|_{\dot{H}^{s,p}(\mathbb{R}^n)},$$

$$\|d\delta u\|_{\dot{H}^{\alpha,\tilde{p}}(\mathbb{R}^n)} + \|\delta du\|_{\dot{H}^{\alpha,\tilde{p}}(\mathbb{R}^n)} \lesssim_{\tilde{p},n,\alpha} \|\nabla^2 u\|_{\dot{H}^{\alpha,\tilde{p}}(\mathbb{R}^n)} \lesssim_{\tilde{p},n,\alpha} \|f\|_{\dot{H}^{\alpha,\tilde{p}}(\mathbb{R}^n)}.$$

In particular, $-\Delta : [\dot{H}^{s+2,p} \cap \dot{H}^{\alpha+2,\tilde{p}}](\mathbb{R}^n, \Lambda^k) \rightarrow [\dot{H}^{s,p} \cap \dot{H}^{\alpha,\tilde{p}}](\mathbb{R}^n, \Lambda^k)$ is an isomorphism of Banach spaces.

(b) *The result still holds if we replace $(\dot{H}^{s,p}, \dot{H}^{\alpha,\tilde{p}}, \dot{H}^{s+2,p}, \dot{H}^{\alpha+2,\tilde{p}})$ by the Besov spaces $(\dot{B}_{p,q}^s, \dot{B}_{\tilde{p},\tilde{q}}^\alpha, \dot{B}_{p,q}^{s+2}, \dot{B}_{\tilde{p},\tilde{q}}^{\alpha+2})$.*

(ii) *For $\mu \in [0, \pi)$, $\lambda \in \Sigma_\mu$, $f \in \mathcal{S}'(\mathbb{R}^n, \Lambda^k)$, consider the problem*

$$\lambda u - \Delta u = f \quad \text{in } \mathbb{R}^n.$$

(a) *If $f \in \dot{H}^{s,p}(\mathbb{R}^n, \Lambda^k)$, then the resolvent problem above admits a unique solution $u \in [\dot{H}^{s,p} \cap \dot{H}^{s+2,p}](\mathbb{R}^n, \Lambda^k)$ with the estimates,*

$$|\lambda| \|u\|_{\dot{H}^{s,p}(\mathbb{R}^n)} + |\lambda|^{\frac{1}{2}} \|\nabla u\|_{\dot{H}^{s,p}(\mathbb{R}^n)} + \|\nabla^2 u\|_{\dot{H}^{s,p}(\mathbb{R}^n)} \lesssim_{p,n,s,\mu} \|f\|_{\dot{H}^{s,p}(\mathbb{R}^n)},$$

$$|\lambda|^{\frac{1}{2}} \|(d + \delta)u\|_{\dot{H}^{s,p}(\mathbb{R}^n)} + \|d\delta u\|_{\dot{H}^{s,p}(\mathbb{R}^n)} + \|\delta du\|_{\dot{H}^{s,p}(\mathbb{R}^n)} \lesssim_{p,n,s,\mu} \|f\|_{\dot{H}^{s,p}(\mathbb{R}^n)}.$$

In particular, $\lambda I - \Delta : [\dot{H}^{s,p} \cap \dot{H}^{s+2,p}](\mathbb{R}^n, \Lambda^k) \rightarrow \dot{H}^{s,p}(\mathbb{R}^n, \Lambda^k)$ is an isomorphism of Banach spaces whenever $(\mathcal{C}_{s,p})$ is satisfied.

Furthermore, the result still holds, replacing $(\dot{H}^{s,p}, \dot{H}^{s,p} \cap \dot{H}^{s+2,p})$ by $(H^{s,p}, H^{s+2,p})$ without any restriction on (s, p) .

(b) *The result above still holds replacing $(\dot{H}^{s,p}, \dot{H}^{s,p} \cap \dot{H}^{s+2,p})$ and $(\mathcal{C}_{s,p})$ by $(\dot{B}_{p,q}^s, \dot{B}_{p,q}^s \cap \dot{B}_{p,q}^{s+2})$ and $(\mathcal{C}_{s,p,q})$, or even by $(B_{p,q}^s, B_{p,q}^{s+2})$ without any restriction on (s, p, q) .*

(iii) *For any $\mu \in (0, \pi)$, the operator $-\Delta$ admits a $\mathbf{H}^\infty(\Sigma_\mu)$ -functional calculus on the function spaces: $[\dot{H}^{s,p} \cap \dot{H}^{\alpha,\tilde{p}}](\mathbb{R}^n, \Lambda^k)$, $(\mathcal{C}_{s,p})$ being satisfied, $\dot{B}_{p,q}^s(\mathbb{R}^n, \Lambda^k)$, $(\mathcal{C}_{s,p,q})$ being satisfied, and on both $H^{s,p}(\mathbb{R}^n, \Lambda^k)$ and $B_{p,q}^s(\mathbb{R}^n, \Lambda^k)$ without any restriction on (s, p, q) .*

Proof. —

Step 1: the scalar Laplacian is injective on $\mathcal{S}'_h(\mathbb{R}^n, \mathbb{C})$. For $f \in \mathcal{S}'(\mathbb{R}^n, \mathbb{C})$, let $u, v \in \mathcal{S}'_h(\mathbb{R}^n, \mathbb{C})$, such that

$$\forall \phi \in \mathcal{S}(\mathbb{R}^n, \mathbb{C}), \langle u, -\Delta\phi \rangle_{\mathbb{R}^n} = \langle f, \phi \rangle_{\mathbb{R}^n} = \langle v, -\Delta\phi \rangle_{\mathbb{R}^n}.$$

Therefore, it follows that $w := u - v \in \mathcal{S}'_h(\mathbb{R}^n, \mathbb{C})$ satisfies

$$\forall \phi \in \mathcal{S}(\mathbb{R}^n, \mathbb{C}), \langle w, -\Delta\phi \rangle_{\mathbb{R}^n} = 0.$$

Hence, one may apply the Fourier transform, to obtain

$$\forall \phi \in \mathcal{S}(\mathbb{R}^n, \mathbb{C}), \langle \mathcal{F}w, |\cdot|^2 \mathcal{F}^{-1}\phi \rangle_{\mathbb{R}^n} = 0,$$

so in particular, for test function in the form $\phi := \mathcal{F}[\frac{\psi}{\|\cdot\|^2}]$, with $\psi \in C_c^\infty(\mathbb{R}^n \setminus \{0\}, \mathbb{C})$ (notice that one can see that $\mathcal{F}[\frac{\psi}{|\cdot|^2}] \in \mathcal{S}(\mathbb{R}^n, \mathbb{C})$), we deduce

$$\forall \psi \in C_c^\infty(\mathbb{R}^n \setminus \{0\}, \mathbb{C}), \langle \mathcal{F}w, \psi \rangle_{\mathbb{R}^n} = 0.$$

Following the proof of [DHMT21, Lemma 3.6], we conclude that $w = 0$ (since $w \in \mathcal{S}'_h(\mathbb{R}^n, \mathbb{C})$ and $\mathcal{S}'_h(\mathbb{R}^n, \mathbb{C})$ does not contain any polynomial), so $u = v$ in $\mathcal{S}'(\mathbb{R}^n, \mathbb{C})$.

Step 2: • For the point (i), it suffices to follow [DM15, Lemma 3.1.1]: In **Step 1**, we have proved uniqueness of solution in $\mathcal{S}'_h(\mathbb{R}^n, \Lambda)$, therefore it suffices to construct a solution. For $f \in \mathcal{S}_0(\mathbb{R}^n, \Lambda)$, the solution is given by

$$u = \mathcal{F}^{-1} \left[\xi \mapsto \frac{\mathcal{F}f(\xi)}{|\xi|^2} \right].$$

We have the estimates

$$\|\nabla^2 u\|_{\dot{H}^{s,p}(\mathbb{R}^n)} \lesssim_{p,n,s} \|f\|_{\dot{H}^{s,p}(\mathbb{R}^n)} \quad \text{and} \quad \|\nabla^2 u\|_{\dot{H}^{\alpha,\tilde{p}}(\mathbb{R}^n)} \lesssim_{\tilde{p},n,\alpha} \|f\|_{\dot{H}^{\alpha,\tilde{p}}(\mathbb{R}^n)}.$$

Since $[\dot{H}^{s,p} \cap \dot{H}^{\alpha,\tilde{p}}](\mathbb{R}^n, \Lambda)$, $[\dot{H}^{s+2,p} \cap \dot{H}^{\alpha+2,\tilde{p}}](\mathbb{R}^n, \Lambda) \subset \mathcal{S}'_h(\mathbb{R}^n, \Lambda)$ are both complete with $\mathcal{S}_0(\mathbb{R}^n, \Lambda)$ as a dense subspace, see [Gau24b, Lemma 2.5], the result still holds for all $f \in [\dot{H}^{s,p} \cap \dot{H}^{\alpha,\tilde{p}}](\mathbb{R}^n, \Lambda)$.

For the case of Besov spaces, we have to proceed a bit differently, since q and \tilde{q} may take the value $+\infty$. First, when $q < +\infty$, as before, we can proceed by density of $\mathcal{S}_0(\mathbb{R}^n, \Lambda)$. When $q = +\infty$ the result follows from real interpolation. Thus, when $(\mathcal{C}_{s+2,p,q})$ is satisfied, we have an isomorphism

$$-\Delta : \dot{B}_{p,q}^{s+2}(\mathbb{R}^n, \Lambda) \longrightarrow \dot{B}_{p,q}^s(\mathbb{R}^n, \Lambda).$$

Now, let $f \in \dot{B}_{p,q}^s(\mathbb{R}^n, \Lambda) \cap \dot{B}_{\tilde{p},\tilde{q}}^\alpha(\mathbb{R}^n, \Lambda) \subset \mathcal{S}'_h(\mathbb{R}^n)$, then there exists a unique $u \in \dot{B}_{p,q}^{s+2}(\mathbb{R}^n, \Lambda) \subset \mathcal{S}'_h(\mathbb{R}^n)$ such that

$$-\Delta u = f.$$

We want to show that $u \in \dot{B}_{\tilde{p},\tilde{q}}^{\alpha+2}(\mathbb{R}^n, \Lambda)$. Since $f \in \dot{B}_{\tilde{p},\tilde{q}}^\alpha(\mathbb{R}^n, \Lambda)$, applying the Littlewood–Paley decomposition and Bernstein’s inequality [BCD11, Lemma 2.1], one obtains

$$2^{2j} \|\dot{\Delta}_j u\|_{L^{\tilde{p}}(\mathbb{R}^n)} \sim_{\tilde{p},n} \|\dot{\Delta}_j [\Delta u]\|_{L^{\tilde{p}}(\mathbb{R}^n)} = \|\dot{\Delta}_j f\|_{L^{\tilde{p}}(\mathbb{R}^n)}.$$

Finally, taking the $\ell_\alpha^{\tilde{q}}(\mathbb{Z})$ -norm on both sides yields

$$\|u\|_{\dot{B}_{p,\tilde{q}}^{\alpha+2}(\mathbb{R}^n)} \sim_{\tilde{p},n,\alpha} \|f\|_{\dot{B}_{p,\tilde{q}}^\alpha(\mathbb{R}^n)}.$$

• For the point (ii), see [ABHN11, Example 3.7.6, Theorem 3.7.11] which imply that for $\mu \in [0, \pi)$, all $p \in [1, +\infty]$, all $\lambda \in \Sigma_\mu$, all $f \in L^p(\mathbb{R}^n, \Lambda)$,

$$\|\lambda(\lambda I - \Delta)^{-1} f\|_{L^p(\mathbb{R}^n)} \lesssim_{p,n,\mu} \|f\|_{L^p(\mathbb{R}^n)}.$$

We also have $(\lambda I - \Delta)^{-1} C_0^0(\mathbb{R}^n, \Lambda) \subset C_0^0(\mathbb{R}^n, \Lambda)$. Moreover, it is not difficult to show that

$$f \mapsto \left[\xi \mapsto \frac{f(\xi)}{\lambda + |\xi|^2} \right]$$

maps continuously $\mathcal{S}(\mathbb{R}^n, \Lambda)$ into itself boundedly. Therefore, the conjugation by the Fourier transform implies that

$$(\lambda I - \Delta)^{-1} : \mathcal{S}(\mathbb{R}^n, \Lambda) \longrightarrow \mathcal{S}(\mathbb{R}^n, \Lambda)$$

is well-defined and bounded, and by duality so is

$$(\lambda I - \Delta)^{-1} : \mathcal{S}'(\mathbb{R}^n, \Lambda) \longrightarrow \mathcal{S}'(\mathbb{R}^n, \Lambda).$$

Finally, we also have $(\lambda I - \Delta)^{-1} \mathcal{S}'_h(\mathbb{R}^n, \Lambda) \subset \mathcal{S}'_h(\mathbb{R}^n, \Lambda)$. Indeed, for $u \in \mathcal{S}'_h(\mathbb{R}^n, \Lambda)$, $\Theta \in C_c^\infty(\mathbb{R}^n)$, we have

$$\begin{aligned} & \|\Theta(\zeta \mathcal{D})(\lambda I - \Delta)^{-1} u\|_{L^\infty(\mathbb{R}^n)} \\ &= \|(\lambda I - \Delta)^{-1} \Theta(\zeta \mathcal{D}) u\|_{L^\infty(\mathbb{R}^n)} \lesssim_{n,\mu} \frac{1}{|\lambda|} \|\Theta(\zeta \mathcal{D}) u\|_{L^\infty(\mathbb{R}^n)} \xrightarrow{\zeta \rightarrow +\infty} 0. \end{aligned}$$

The estimates are then direct consequences of the L^p -ones. Let $f \in \dot{H}^{s,p}(\mathbb{R}^n, \Lambda)$, we have $u := (\lambda I - \Delta)^{-1} f \in \mathcal{S}'_h(\mathbb{R}^n)$, and by the Fatou Lemma

$$\begin{aligned} \|u\|_{\dot{H}^{s,p}(\mathbb{R}^n)} &\leq \liminf_{N \rightarrow +\infty} \left\| \sum_{j=-N}^N (-\Delta)^{\frac{s}{2}} \dot{\Delta}_j u \right\|_{L^p(\mathbb{R}^n)} \\ &\leq \liminf_{N \rightarrow +\infty} \left\| \sum_{j=-N}^N (\lambda I - \Delta)^{-1} (-\Delta)^{\frac{s}{2}} \dot{\Delta}_j f \right\|_{L^p(\mathbb{R}^n)} \\ &\lesssim_{p,n,\mu} \frac{1}{|\lambda|} \liminf_{N \rightarrow +\infty} \left\| \sum_{j=-N}^N (-\Delta)^{\frac{s}{2}} \dot{\Delta}_j f \right\|_{L^p(\mathbb{R}^n)} \\ &\lesssim_{p,n,\mu} \frac{1}{|\lambda|} \|f\|_{\dot{H}^{s,p}(\mathbb{R}^n)}. \end{aligned}$$

For the remaining estimates in the case of homogeneous Sobolev spaces, one may proceed similarly. Note that we only took advantage of the completeness of Lebesgue spaces up to now.

For Besov spaces, we have by Bernstein’s inequality [BCD11, Lemma 2.1],

$$\|\dot{\Delta}_j u\|_{L^p(\mathbb{R}^n)} = \|(\lambda I - \Delta)^{-1} \dot{\Delta}_j f\|_{L^p(\mathbb{R}^n)} \lesssim_{p,n,\mu} \frac{1}{|\lambda|} \|\dot{\Delta}_j f\|_{L^p(\mathbb{R}^n)},$$

but, we also have

$$2^{2j} \left\| \dot{\Delta}_j u \right\|_{L^p(\mathbb{R}^n)} \sim_{p,n} \left\| \dot{\Delta}_j [\Delta u] \right\|_{L^p(\mathbb{R}^n)} = \left\| \Delta (\lambda I - \Delta)^{-1} \dot{\Delta}_j f \right\|_{L^p(\mathbb{R}^n)} \\ \lesssim_{p,n,\mu} \left\| \dot{\Delta}_j f \right\|_{L^p(\mathbb{R}^n)}.$$

Therefore, taking the $\ell_s^q(\mathbb{Z})$ -norms yields the desired estimates

$$|\lambda| \|u\|_{\dot{B}_{p,q}^s(\mathbb{R}^n)} \lesssim_{p,n,\mu} \|f\|_{\dot{B}_{p,q}^s(\mathbb{R}^n)} \text{ and } \|u\|_{\dot{B}_{p,q}^{s+2}(\mathbb{R}^n)} \lesssim_{p,n,\mu} \|f\|_{\dot{B}_{p,q}^s(\mathbb{R}^n)}.$$

The remaining estimates follow the same lines.

• For the point (iii), the result on $L^p(\mathbb{R}^n, \mathbb{C})$ is a consequence of a more general one which is [Haa06, Proposition 8.3.4]. \square

Similarly, thanks again to standard Fourier analysis, we can introduce appropriate differential form-valued versions of Riesz transforms for the Hodge Laplacian. Their boundedness on appropriate function spaces are again carried over by their scalar analogue, and a direct consequence will be an explicit formula for our generalized Leray projector \mathbb{P} on \mathbb{R}^n .

To do so, we notice that one can write associated Fourier symbols, thanks to Lemma 2.5, to obtain

$$d(-\Delta)^{-\frac{1}{2}} = \sum_{k=1}^n R_k \mathbf{e}_k \wedge \quad \text{and} \quad \delta(-\Delta)^{-\frac{1}{2}} = - \sum_{k=1}^n R_k \mathbf{e}_k \lrcorner$$

where for $k \in \llbracket 1, n \rrbracket$, R_k is the k^{th} Riesz transform on \mathbb{R}^n given by the Fourier symbol $\xi \mapsto \frac{i\xi_k}{|\xi|}$, which is well known to be bounded on $L^p(\mathbb{R}^n, \mathbb{C})$, $1 < p < +\infty$, see [Ste70, Chapter 2, Theorem 1 & Chapter 3, Section 1]. Therefore, the next proposition follows naturally.

PROPOSITION 2.8. — *Let $p, \tilde{p} \in (1, +\infty)$, $q, \tilde{q} \in [1, +\infty]$, $s, \alpha \in \mathbb{R}$, and let $k \in \llbracket 0, n \rrbracket$. The operators*

$$d(-\Delta)^{-\frac{1}{2}}, \delta(-\Delta)^{-\frac{1}{2}}, d\delta(-\Delta)^{-1} \text{ and } \delta d(-\Delta)^{-1}$$

are all well-defined bounded linear operators,

- On $[\dot{H}^{s,p} \cap \dot{H}^{\alpha,\tilde{p}}](\mathbb{R}^n, \Lambda)$, and on $[\dot{B}_{p,q}^s \cap \dot{B}_{\tilde{p},\tilde{q}}^\alpha](\mathbb{R}^n, \Lambda)$, $(\mathcal{C}_{s,p})$ being satisfied in case of Sobolev spaces, $(\mathcal{C}_{s,p,q})$ in the case of Besov spaces. Moreover, we have decoupled estimates,

$$\|Tu\|_{X^s(\mathbb{R}^n)} \lesssim_{p,n,s} \|u\|_{X^s(\mathbb{R}^n)}$$

and

$$\|Tu\|_{X^\alpha(\mathbb{R}^n)} \lesssim_{p,n,\alpha} \|u\|_{X^\alpha(\mathbb{R}^n)}, \quad u \in [X^s \cap X^\alpha](\mathbb{R}^n, \Lambda^k),$$

where $(X^s, X^\alpha) \in \{(\dot{H}^{s,p}, \dot{H}^{\alpha,\tilde{p}}), (\dot{B}_{p,q}^s, \dot{B}_{\tilde{p},\tilde{q}}^\alpha)\}$ and T denotes any of the operators above.

- On $H^{s,p}(\mathbb{R}^n, \Lambda)$, and on $B_{p,q}^s(\mathbb{R}^n, \Lambda)$, without any restriction on (s, p, q) .

Moreover, the following identity holds

$$\left(d(-\Delta)^{-\frac{1}{2}} + \delta(-\Delta)^{-\frac{1}{2}} \right)^2 = I.$$

THEOREM 2.9. — *Let $p \in (1, +\infty)$, $q \in [1, +\infty]$, $s \in \mathbb{R}$, and let $k \in \llbracket 0, n \rrbracket$. The following hold*

(i) *The following equality is true, whenever $(\mathcal{C}_{s,p})$ is satisfied,*

$$\dot{N}_p^s(\mathfrak{d}, \mathbb{R}^n, \Lambda^k) = \overline{\dot{R}_p^s(\mathfrak{d}, \mathbb{R}^n, \Lambda^k)}^{\|\cdot\|_{\dot{H}^{s,p}(\mathbb{R}^n)}},$$

and still holds replacing \mathfrak{d} by δ .

(ii) *The (generalized) Helmholtz–Leray projector is well-defined and bounded as a linear map*

$$\mathbb{P} : \dot{H}^{s,p}(\mathbb{R}^n, \Lambda^k) \longrightarrow \dot{N}_p^s(\delta, \mathbb{R}^n, \Lambda^k),$$

whenever $(\mathcal{C}_{s,p})$ is satisfied. Moreover, the following identity is true

$$\mathbb{P} = \mathbb{I} - \mathfrak{d}(-\Delta)^{-1}\delta.$$

(iii) *The following Hodge decomposition holds whenever $(\mathcal{C}_{s,p})$ is satisfied,*

$$\dot{H}^{s,p}(\mathbb{R}^n, \Lambda^k) = \dot{N}_p^s(\delta, \mathbb{R}^n, \Lambda^k) \oplus \dot{N}_p^s(\mathfrak{d}, \mathbb{R}^n, \Lambda^k).$$

Everything above still holds replacing $(\dot{H}^{s,p}, \dot{N}_p^s, \dot{R}_p^s)$ by either $(\dot{B}_{p,q}^s, \dot{N}_{p,q}^s, \dot{R}_{p,q}^s)$, $(\mathcal{C}_{s,p,q})$ being satisfied, $(H^{s,p}, N_p^s, R_p^s)$ or $(B_{p,q}^s, N_{p,q}^s, R_{p,q}^s)$ without any restriction on (s, p, q) . In case of Besov spaces with $q = +\infty$, the density result of point (i) only holds in the weak sense.*

Remark 2.10. — *On Λ^1 -valued functions identified with vector fields one recovers the usual well known formula, i.e.*

$$\mathbb{P} = \mathbb{I} + \nabla(-\Delta)^{-1}\operatorname{div}.$$

Proof. —

Step 1: The orthogonal projector \mathbb{P} is originally defined only as an operator

$$\mathbb{P} : L^2(\mathbb{R}^n, \Lambda^k) \longrightarrow N_2(\delta, \mathbb{R}^n, \Lambda^k).$$

We claim that \mathbb{P} is equal to the operator formally given by

$$\tilde{\mathbb{P}} := \mathbb{I} - \mathfrak{d}(-\Delta)^{-1}\delta.$$

The proof is standard, and works as in the case of vector fields, and then is left to the reader.

Step 2: Previous step and Proposition 2.8 give that \mathbb{P} is bounded $\dot{H}^{s,p}(\mathbb{R}^n, \Lambda^k)$.

For $u \in \dot{H}^{s,p}(\mathbb{R}^n, \Lambda^k)$, $v \in \mathcal{S}(\mathbb{R}^n)$, we regularize with the resolvent, to compute

$$\begin{aligned} \langle \mathbb{P}u, dv \rangle_{\mathbb{R}^n} &= \lim_{\lambda \rightarrow 0_+} \langle u, dv \rangle_{\mathbb{R}^n} - \langle \mathfrak{d}(\lambda\mathbb{I} - \Delta)^{-1}\delta u, dv \rangle_{\mathbb{R}^n} \\ &= \lim_{\lambda \rightarrow 0_+} \langle u, dv \rangle_{\mathbb{R}^n} - \langle u, (\lambda\mathbb{I} - \Delta)^{-1}\mathfrak{d}\delta dv \rangle_{\mathbb{R}^n} \\ &= \lim_{\lambda \rightarrow 0_+} \langle u, dv \rangle_{\mathbb{R}^n} - \langle u, (\lambda\mathbb{I} - \Delta)^{-1}(-\Delta)dv \rangle_{\mathbb{R}^n} \\ &= \langle u, dv \rangle_{\mathbb{R}^n} - \langle u, dv \rangle_{\mathbb{R}^n} = 0. \end{aligned}$$

Hence $\mathbb{P}\dot{H}^{s,p}(\mathbb{R}^n, \Lambda^k) \subset \dot{N}_p^s(\delta, \mathbb{R}^n, \Lambda^k)$, and we even have $\mathbb{P}|_{\dot{N}_p^s(\delta, \mathbb{R}^n, \Lambda^k)} = \text{I}$, so that

$$\mathbb{P}\dot{H}^{s,p}(\mathbb{R}^n, \Lambda^k) = \dot{N}_p^s(\delta, \mathbb{R}^n, \Lambda^k).$$

Similarly, $[\text{I} - \mathbb{P}]\dot{H}^{s,p}(\mathbb{R}^n, \Lambda^k) \subset \dot{N}_p^s(\underline{d}, \mathbb{R}^n, \Lambda^k)$, and $[\text{I} - \mathbb{P}]|_{\dot{N}_p^s(\underline{d}, \mathbb{R}^n, \Lambda^k)} = \text{I}$, which comes from

$$[\text{I} - \mathbb{P}]u = \lim_{\lambda \rightarrow 0^+} d(\lambda \text{I} - \Delta)^{-1} \delta u, \quad u \in \dot{H}^{s,p}(\mathbb{R}^n, \Lambda^k).$$

This also gives $\dot{N}_p^s(\underline{d}, \mathbb{R}^n, \Lambda^k) = \overline{\dot{R}_p^s(\underline{d}, \mathbb{R}^n, \Lambda^k)}^{\|\cdot\|_{\dot{H}^{s,p}(\mathbb{R}^n)}}$.

The proof is straightforward the same for other function spaces. □

2.3. The case of the half-space \mathbb{R}_+^n

2.3.1. L^2 -theory for Hodge Laplacians and the Hodge decomposition

The following lemma is fundamental for the analysis of the L^2 theory of the Hodge Laplacian when one has explicit access to the boundary, and, moreover, several proofs presented here do not depend on the open set Ω , here $\Omega = \mathbb{R}_+^n$, and remain valid as long as integration by parts formulas (2.1) and (2.2) and partial trace results for vector fields are available.

LEMMA 2.11. — *Let $k \in \llbracket 0, n \rrbracket$. We set*

$$\begin{aligned} D_2(\delta, \mathbb{R}_+^n, \Lambda^k) &:= \left\{ u \in D_2(\delta, \mathbb{R}_+^n, \Lambda^k) \mid \nu \lrcorner u|_{\partial\mathbb{R}_+^n} = 0 \right\}, \\ D_2(\underline{d}, \mathbb{R}_+^n, \Lambda^k) &:= \left\{ u \in D_2(\underline{d}, \mathbb{R}_+^n, \Lambda^k) \mid \nu \wedge u|_{\partial\mathbb{R}_+^n} = 0 \right\}. \end{aligned}$$

The operator $(d_2(\underline{d}, \mathbb{R}_+^n, \Lambda^k), \underline{d})$ is an unbounded densely defined closed operator, with adjoint

$$(2.3) \quad (D_2(\delta^*, \mathbb{R}_+^n, \Lambda^{k+1}), \delta^*) = (D_2(\delta, \mathbb{R}_+^n, \Lambda^{k+1}), \delta).$$

Similarly, $(d_2(\delta, \mathbb{R}_+^n, \Lambda^k), \delta)$ is an unbounded densely defined closed operator, with adjoint

$$(2.4) \quad (D_2(\delta^*, \mathbb{R}_+^n, \Lambda^{k-1}), \delta^*) = (D_2(\underline{d}, \mathbb{R}_+^n, \Lambda^{k-1}), \underline{d}).$$

Proof. — Closedness is straightforward by Theorem A.1, and the fact that both are densely defined is straightforward since the space $C_c^\infty(\mathbb{R}_+^n, \Lambda^k)$ is contained in both domains. We just prove the duality identity (2.3), the proof of (2.4) is similar. Let $u \in D_2(\delta, \mathbb{R}_+^n, \Lambda^k)$, then for all $v \in \mathcal{S}_0(\overline{\mathbb{R}_+^n}, \Lambda^k)$, we can use Theorem A.1, to obtain that

$$\langle v, \delta u \rangle_{\mathbb{R}_+^n} = \langle dv, u \rangle_{\mathbb{R}_+^n}.$$

Thus, by Cauchy–Schwarz inequality

$$\left| \langle dv, u \rangle_{\mathbb{R}_+^n} \right| \leq \| \delta u \|_{L^2(\mathbb{R}_+^n)} \| v \|_{L^2(\mathbb{R}_+^n)}.$$

Hence, $v \mapsto \langle dv, u \rangle_{\mathbb{R}_+^n}$ extends uniquely as a bounded linear functional on $L^2(\mathbb{R}_+^n, \Lambda^k)$, so that necessarily $(D_2(\underline{\delta}, \mathbb{R}_+^n, \Lambda^k), \delta) \subset (D_2(d^*, \mathbb{R}_+^n, \Lambda^{k+1}), d^*)$.

For the reverse inclusion, let $u \in D_2(d^*, \mathbb{R}_+^n, \Lambda^{k+1})$, for all $v \in D_2(d, \mathbb{R}_+^n, \Lambda^k)$, we have

$$\langle v, d^*u \rangle_{\mathbb{R}_+^n} = \langle dv, u \rangle_{\mathbb{R}_+^n}.$$

In particular, for $v \in C_c^\infty(\mathbb{R}_+^n, \Lambda^k)$, it yields that

$$\langle v, d^*u \rangle_{\mathbb{R}_+^n} = \langle dv, u \rangle_{\mathbb{R}_+^n} = \langle v, \delta u \rangle_{\mathbb{R}_+^n}.$$

Hence, $d^*u = \delta u$ in $\mathcal{D}'(\mathbb{R}_+^n, \Lambda^k)$, then in $L^2(\mathbb{R}_+^n, \Lambda^k)$, so that for all $v \in D_2(d, \mathbb{R}_+^n, \Lambda^k)$,

$$\langle v, \delta u \rangle_{\mathbb{R}_+^n} = \langle dv, u \rangle_{\mathbb{R}_+^n}.$$

From above equality, we apply Theorem A.1 to check $\nu \lrcorner u|_{\partial\mathbb{R}_+^n} = 0$ and deduce

$$(D_2(d^*, \mathbb{R}_+^n, \Lambda^{k+1}), d^*) \subset (D_2(\underline{\delta}, \mathbb{R}_+^n, \Lambda^{k+1}), \delta),$$

the proof being therefore complete. □

In particular, since $(D_2(d^*, \mathbb{R}_+^n, \Lambda^k), d^*)$ and $(D_2(\delta, \mathbb{R}_+^n, \Lambda^k), \delta)$ are closed operators, both

$$N_2(\delta, \mathbb{R}_+^n, \Lambda^k) \quad \text{and} \quad N_2(d^*, \mathbb{R}_+^n, \Lambda^k)$$

are closed subspaces of $L^2(\mathbb{R}_+^n, \Lambda^k)$. Thus, the following orthogonal projections are well-defined and bounded

$$\begin{aligned} \mathbb{P} : L^2(\mathbb{R}_+^n, \Lambda^k) &\longrightarrow N_2(d^*, \mathbb{R}_+^n, \Lambda^k), & [\mathbb{I} - \mathbb{P}] : L^2(\mathbb{R}_+^n, \Lambda^k) &\longrightarrow \overline{R_2(d, \mathbb{R}_+^n, \Lambda^k)}, \\ \mathbb{Q} : L^2(\mathbb{R}_+^n, \Lambda^k) &\longrightarrow N_2(\delta, \mathbb{R}_+^n, \Lambda^k), & [\mathbb{I} - \mathbb{Q}] : L^2(\mathbb{R}_+^n, \Lambda^k) &\longrightarrow \overline{R_2(\delta^*, \mathbb{R}_+^n, \Lambda^k)}, \end{aligned}$$

which induce topological Hodge decompositions

$$\begin{aligned} (\mathfrak{H}_2) \quad L^2(\mathbb{R}_+^n, \Lambda^k) &= \overline{R_2(d, \mathbb{R}_+^n, \Lambda^k)} \oplus^\perp N_2(d^*, \mathbb{R}_+^n, \Lambda^k), \\ &= \overline{R_2(\delta^*, \mathbb{R}_+^n, \Lambda^k)} \oplus^\perp N_2(\delta, \mathbb{R}_+^n, \Lambda^k). \end{aligned}$$

LEMMA 2.12. — For $k \in \llbracket 0, n \rrbracket$, the following Hodge–Dirac operators

$$\begin{aligned} (D_2(D_n, \mathbb{R}_+^n, \Lambda^k), D_n) &= (D_2(d, \mathbb{R}_+^n, \Lambda^k) \cap D_2(d^*, \mathbb{R}_+^n, \Lambda^k), d + d^*), \\ (D_2(D_t, \mathbb{R}_+^n, \Lambda^k), D_t) &= (D_2(\delta^*, \mathbb{R}_+^n, \Lambda^k) \cap D_2(\delta, \mathbb{R}_+^n, \Lambda^k), \delta^* + \delta), \end{aligned}$$

are both densely defined closed operators on $L^2(\mathbb{R}_+^n, \Lambda^k)$.

Proof. — Let $(u_j)_{j \in \mathbb{N}} \subset D_2(d, \mathbb{R}_+^n, \Lambda^k) \cap D_2(d^*, \mathbb{R}_+^n, \Lambda^k)$, and $(u, v) \in L^2(\mathbb{R}_+^n, \Lambda^k) \times L^2(\mathbb{R}_+^n, \Lambda)$ such that it satisfies

$$u_j \xrightarrow{j \rightarrow +\infty} u \quad \text{and} \quad D_n u_j \xrightarrow{j \rightarrow +\infty} v \quad \text{in } L^2(\mathbb{R}_+^n).$$

By the Hodge decomposition (5.2), there exists a unique couple of elements $(v_0, v_1) \in \mathbb{R}_2(d, \mathbb{R}_+^n, \Lambda) \times \mathbb{N}_2(d^*, \mathbb{R}_+^n, \Lambda)$ such that $v = v_0 + v_1$. Since u_j goes to u in $L^2(\mathbb{R}_+^n, \Lambda^k)$, by continuity of involved projectors, and uniqueness of decomposition, it follows that

$$du_j \xrightarrow{j \rightarrow +\infty} v_0 \quad \text{and} \quad d^*u_j \xrightarrow{j \rightarrow +\infty} v_1 \quad \text{in } L^2(\mathbb{R}_+^n, \Lambda).$$

But $(u_j)_{j \in \mathbb{N}}$ converges to u in $L^2(\mathbb{R}_+^n, \Lambda^k)$, so in particular in distributional sense, thus necessarily $(v_0, v_1) = (du, d^*u)$ and $v = D_n u$, i.e. $(D_2(D_n, \mathbb{R}_+^n, \Lambda^k), D_n)$ is closed on $L^2(\mathbb{R}_+^n, \Lambda^k)$. The proof ends here since one can reproduce all above arguments for $(D_2(D_t, \mathbb{R}_+^n, \Lambda^k), D_t)$. □

PROPOSITION 2.13. — *The Hodge–Dirac operator $(D_2(D., \mathbb{R}_+^n, \Lambda), D.)$ is an injective self-adjoint 0-bisectorial operator on $L^2(\mathbb{R}_+^n, \Lambda)$ so that it satisfies the following bound, for all $\theta \in (0, \frac{\pi}{2})$,*

$$(2.5) \quad \forall \mu \in \mathbb{C} \setminus \overline{S}_\theta, \left\| \mu(\mu I + D.)^{-1} \right\|_{L^2(\mathbb{R}_+^n) \rightarrow L^2(\mathbb{R}_+^n)} \leq \frac{1}{\sin(\theta)}.$$

Moreover, it admits a bounded $(\mathbf{H}^\infty(S_\theta)$ -)functional calculus on $L^2(\mathbb{R}_+^n, \Lambda)$ with bound 1, i.e., for all $f \in \mathbf{H}^\infty(S_\theta)$, $u \in L^2(\mathbb{R}_+^n, \Lambda)$,

$$(2.6) \quad \|f(D.)u\|_{L^2(\mathbb{R}_+^n)} \leq \|f\|_{L^\infty(S_\theta)} \|u\|_{L^2(\mathbb{R}_+^n)}.$$

Remark 2.14. — Proposition 2.13 does not depend on the fact $\Omega = \mathbb{R}_+^n$. See [MM18, Section 2] where the same result is stated for bounded (even weak-)Lipschitz domains.

Proof. — The resolvent bound (2.5) is usual since $(D_2(D.), D.)$ is self-adjoint by construction, see [Haa06, Proposition C.4.2]. The fact that it admits a bounded holomorphic functional calculus follows from [McI86, Section 10]. □

For $k \in \llbracket 0, n \rrbracket$, the Hodge Laplacian $(D(\Delta_{\mathcal{H}}, \mathbb{R}_+^n, \Lambda^k), -\Delta_{\mathcal{H}})$ can be realized on $L^2(\mathbb{R}_+^n, \Lambda^k)$ by means of densely defined, symmetric, accretive, continuous, closed, sesquilinear forms on $L^2(\mathbb{R}_+^n, \Lambda^k)$, for

$$(2.7) \quad \mathfrak{a}_{\mathcal{H}} : D_2(\mathfrak{a}_{\mathcal{H}}, \Lambda^k)^2 \ni (u, v) \longmapsto \int_{\mathbb{R}_+^n} \langle du(x), dv(x) \rangle dx + \int_{\mathbb{R}_+^n} \langle \delta u(x), \delta v(x) \rangle dx$$

with $D_2(\mathfrak{a}_{\mathcal{H},n}, \Lambda^k) = D_2(d, \Lambda^k) \cap D_2(d^*, \Lambda^k)$, $D_2(\mathfrak{a}_{\mathcal{H},t}) = D_2(\delta^*, \Lambda^k) \cap D_2(\delta, \Lambda^k)$, so that it is easy to see that both are closed, densely defined, non-negative self-adjoint operators on $L^2(\mathbb{R}_+^n, \Lambda^k)$. See [Ouh05, Chapter 1] for more details about realization of operators via sesquilinear forms on a Hilbert space. The next theorem is a standard consequence.

THEOREM 2.15. — *Let $k \in \llbracket 0, n \rrbracket$. The operator $(D_2(\Delta_{\mathcal{H}}, \mathbb{R}_+^n, \Lambda^k), -\Delta_{\mathcal{H}})$ is an injective non-negative self-adjoint and 0-sectorial operator on $L^2(\mathbb{R}_+^n, \Lambda^k)$, which admits a $\mathbf{H}^\infty(\Sigma_\theta)$ -functional calculus for all $\theta \in (0, \pi)$.*

Moreover, the following hold

- (i) $D_2(\Delta_{\mathcal{H}}, \mathbb{R}_+^n, \Lambda^k)$ is a closed subspace of $H^{2,2}(\mathbb{R}_+^n, \Lambda)$;

(ii) *Provided $\mu \in [0, \pi)$, for $\lambda \in \Sigma_\mu$, $f \in L^2(\mathbb{R}_+^n, \Lambda^k)$, then $u := (\lambda I - \Delta_{\mathcal{H}})^{-1} f$ satisfies*

$$(2.8) \quad |\lambda| \|u\|_{L^2(\mathbb{R}_+^n)} + |\lambda|^{\frac{1}{2}} \|D.u\|_{L^2(\mathbb{R}_+^n)} + \|\Delta u\|_{L^2(\mathbb{R}_+^n)} \lesssim_\mu \|f\|_{L^2(\mathbb{R}_+^n)};$$

$$(2.9) \quad |\lambda| \|u\|_{L^2(\mathbb{R}_+^n)} + |\lambda|^{\frac{1}{2}} \|\nabla u\|_{L^2(\mathbb{R}_+^n)} + \|\nabla^2 u\|_{L^2(\mathbb{R}_+^n)} \lesssim_{n,\mu} \|f\|_{L^2(\mathbb{R}_+^n)};$$

(iii) *The following resolvent identity holds for all $\mu \in [0, \pi)$, $\lambda \in \Sigma_\mu$, $f \in L^2(\mathbb{R}_+^n, \Lambda^k)$,*

$$E_{\mathcal{H}}(\lambda I - \Delta_{\mathcal{H}})^{-1} f = (\lambda I - \Delta)^{-1} E_{\mathcal{H}} f.$$

(For the definition of $E_{\mathcal{H}}$, see (2.10) below.)

Remark 2.16. — In above Theorem 2.15, points (i) and (iii), as well as the estimate (2.9) of point (ii) are the only points that rely on the fact that the considered open set is $\Omega = \mathbb{R}_+^n$, but mainly the point (iii) is used to deduce the previous ones. The beginning of the statement, as well as (2.8), does not rely on any particular structure, and remains true on any open set Ω .

Every other result below, in the present subsection about L^2 -theory of Hodge Laplacians and the Hodge decomposition, remain true on general domains Ω as long as one can show that the Hodge Laplacian is injective.

Before proving Theorem 2.15, following [Gau24b, Section 5] for $\mathcal{J} \in \{\mathcal{D}, \mathcal{N}\}$, we introduce the following extension operator defined for any measurable function u on \mathbb{R}_+^n , for almost every $x = (x', x_n) \in \mathbb{R}^n$:

$$E_{\mathcal{D}} u(x', x_n) := \begin{cases} u(x', x_n), & \text{if } (x', x_n) \in \mathbb{R}^{n-1} \times \mathbb{R}_+, \\ -u(x', -x_n), & \text{if } (x', x_n) \in \mathbb{R}^{n-1} \times \mathbb{R}_*^-; \end{cases}$$

$$E_{\mathcal{N}} u(x', x_n) := \begin{cases} u(x', x_n), & \text{if } (x', x_n) \in \mathbb{R}^{n-1} \times \mathbb{R}_+, \\ u(x', -x_n), & \text{if } (x', x_n) \in \mathbb{R}^{n-1} \times \mathbb{R}_*^-. \end{cases}$$

Now, we specify the definition of the extension operators $E_{\mathcal{H},j}$, $j \in \{\mathbf{n}, \mathbf{t}\}$, on measurable functions $u : \mathbb{R}_+^n \rightarrow \Lambda^k$, provided $k \in \llbracket 0, n \rrbracket$, $I \in \mathcal{I}_n^k$,

$$(2.10) \quad (E_{\mathcal{H},\mathbf{n}} u)_I := \begin{cases} E_{\mathcal{D}} u_I, & \text{if } n \in I, \\ E_{\mathcal{N}} u_I, & \text{if } n \notin I; \end{cases} \quad \text{and} \quad (E_{\mathcal{H},\mathbf{t}} u)_I := \begin{cases} E_{\mathcal{N}} u_I, & \text{if } n \in I, \\ E_{\mathcal{D}} u_I, & \text{if } n \notin I; \end{cases}$$

For $u : \mathbb{R}_+^n \rightarrow \Lambda^k$, we also set

$$\tilde{u}_j := [E_{\mathcal{H},j} u]_{|\mathbb{R}^n}.$$

By construction, for $j \in \{\mathbf{n}, \mathbf{t}\}$, $s \in (-1+1/p, 1/p)$, $p \in (1, +\infty)$, the Proposition 1.7 leads to the boundedness of

$$(2.11) \quad E_{\mathcal{H},j} : \dot{H}^{s,p}(\mathbb{R}_+^n, \Lambda) \rightarrow \dot{H}^{s,p}(\mathbb{R}^n, \Lambda).$$

The same result holds replacing $\dot{H}^{s,p}$ by either $H^{s,p}$, $B_{p,q}^s$, or even by $\dot{B}_{p,q}^s$, $q \in [1, +\infty]$.

LEMMA 2.17. — For all $u \in D_2(d, \mathbb{R}_+^n, \Lambda^k)$ (resp. $D_2(d^*, \mathbb{R}_+^n, \Lambda^k)$), we have

$$E_{\mathcal{H},n}u \in D_2(d, \mathbb{R}^n, \Lambda^k) \quad (\text{resp. } D_2(\delta, \mathbb{R}^n, \Lambda^k))$$

with the formula

$$dE_{\mathcal{H},n}u = E_{\mathcal{H},n}du \quad (\text{resp. } \delta E_{\mathcal{H},n}u = E_{\mathcal{H},n}d^*u).$$

Proof. — Let $u \in D_2(d, \mathbb{R}_+^n, \Lambda^k)$, for $v \in \mathcal{S}(\mathbb{R}^n, \Lambda^{k+1})$,

$$\begin{aligned} \langle E_{\mathcal{H},n}u, \delta v \rangle_{\mathbb{R}^n} &= \langle u, \delta v \rangle_{\mathbb{R}_+^n} + \langle \tilde{u}_n, \delta v \rangle_{\mathbb{R}_-^n} \\ &= \langle du, v \rangle_{\mathbb{R}_+^n} + \langle (-\mathbf{e}_n) \wedge u, v \rangle_{\partial\mathbb{R}_+^n} + \langle \tilde{d}u_n, v \rangle_{\mathbb{R}_-^n} + \langle (\mathbf{e}_n) \wedge \tilde{u}_n, v \rangle_{\partial\mathbb{R}_-^n} \\ &= \langle du, v \rangle_{\mathbb{R}_+^n} + \langle \tilde{d}u_n, v \rangle_{\mathbb{R}_-^n} \\ &= \langle E_{\mathcal{H},n}du, v \rangle_{\mathbb{R}^n}. \end{aligned}$$

Which holds, since $(\mathbf{e}_n) \wedge \tilde{u}_n(\cdot, 0) = (\mathbf{e}_n) \wedge u(\cdot, 0)$.

Now, if $u \in D_2(d^*, \mathbb{R}_+^n, \Lambda^k)$, for $v \in \mathcal{S}(\mathbb{R}^n, \Lambda^{k-1})$,

$$\begin{aligned} \langle E_{\mathcal{H},n}u, dv \rangle_{\mathbb{R}^n} &= \langle u, dv \rangle_{\mathbb{R}_+^n} + \langle \tilde{u}_n, dv \rangle_{\mathbb{R}_-^n} \\ &= \langle \delta u, v \rangle_{\mathbb{R}_+^n} + \langle (-\mathbf{e}_n) \lrcorner u, v \rangle_{\partial\mathbb{R}_+^n} + \langle \tilde{\delta}u_n, v \rangle_{\mathbb{R}_-^n} + \langle (\mathbf{e}_n) \lrcorner \tilde{u}_n, v \rangle_{\partial\mathbb{R}_-^n} \\ &= \langle \delta u, v \rangle_{\mathbb{R}_+^n} + \langle \tilde{\delta}u_n, v \rangle_{\mathbb{R}_-^n} \\ &= \langle E_{\mathcal{H},n}d^*u, v \rangle_{\mathbb{R}^n}. \end{aligned}$$

Which holds, since $(\mathbf{e}_n) \lrcorner \tilde{u}_n(\cdot, 0) = -(\mathbf{e}_n) \lrcorner u(\cdot, 0) = 0$. □

Proof of Theorem 2.15. — By the realization of the Hodge Laplacian by means of the sesquilinear form (2.7), we have $(D_2(\Delta_{\mathcal{H}}, \mathbb{R}_+^n, \Lambda^k), -\Delta_{\mathcal{H}}) = (D_2(D^2, \mathbb{R}_+^n, \Lambda^k), D^2)$. Thus, as the square of a self-adjoint 0-bisectorial operator, the Hodge Laplacian is a non-negative self-adjoint 0-sectorial operator on $L^2(\mathbb{R}_+^n, \Lambda^k)$, and it also admits a bounded holomorphic functional calculus, see for instance [Ege15, Theorem 3.2.20]. In particular, (2.8) in point (ii) holds.

For now, we only consider the case $(D_2(\Delta_{\mathcal{H},n}, \mathbb{R}_+^n, \Lambda^k), -\Delta_{\mathcal{H},n})$, the proof could be achieved in similar fashion for $(D_2(\Delta_{\mathcal{H},t}, \mathbb{R}_+^n, \Lambda^k), -\Delta_{\mathcal{H},t})$.

For $\lambda \in \Sigma_\mu$, $\mu \in (0, \pi)$, $f \in L^2(\mathbb{R}_+^n, \Lambda^k)$, we set $U := (\lambda I - \Delta)^{-1}E_{\mathcal{H}}f \in H^{2,2}(\mathbb{R}^n, \Lambda^k)$. By construction, as in the proof of [Gau24b, Proposition 5.1], for $I \in \mathcal{I}_n^k$ we have

$$\begin{aligned} U|_{\partial\mathbb{R}_+^n} &= 0, \text{ provided } n \in I, \\ \partial_{x_n} U|_{\partial\mathbb{R}_+^n} &= 0, \text{ provided } n \notin I. \end{aligned}$$

Therefore, we obtain first,

$$\begin{aligned} \nu \lrcorner u|_{\partial\mathbb{R}_+^n} &= -\mathbf{e}_n \lrcorner u|_{\partial\mathbb{R}_+^n} = (-1)^k \sum_{1 \leq \ell_1 < \dots < \ell_{k-1} < n} u_{\ell_1 \ell_2 \dots \ell_{k-1} n}(\cdot, 0) dx_{\ell_1} \wedge \dots \wedge dx_{\ell_{k-1}} \\ &= (-1)^k \sum_{I' \in \mathcal{I}_{n-1}^{k-1}} u_{I',n}(\cdot, 0) dx_{I'} = 0. \end{aligned}$$

Similarly, we get that

$$\begin{aligned} \nu \lrcorner du|_{\partial\mathbb{R}_+^n} &= -\mathbf{e}_n \lrcorner du|_{\partial\mathbb{R}_+^n} \\ &= -\sum_{j=1}^{n-1} \sum_{1 \leq \ell_1 < \dots < \ell_{k-1} < n} \partial_{x_j} u_{\ell_1 \dots \ell_{k-1} n}(\cdot, 0) dx_j \wedge dx_{\ell_1} \wedge \dots \wedge dx_{\ell_{k-1}} \\ &\quad - \sum_{1 \leq \ell_1 < \dots < \ell_{k-1} < \ell_k < n} \partial_{x_n} u_{\ell_1 \dots \ell_k}(\cdot, 0) dx_{\ell_1} \wedge \dots \wedge dx_{\ell_k} \\ &= 0. \end{aligned}$$

From above calculations, we deduce that $u := U|_{\mathbb{R}_+^n}$ is such that $u \in H^{2,2}(\mathbb{R}_+^n, \Lambda^k) \cap D_2(\Delta_{\mathcal{H},n}, \mathbb{R}_+^n, \Lambda^k)$, and

$$\lambda u - \Delta u = f \text{ in } \mathbb{R}_+^n.$$

Hence, by uniqueness $(\lambda I - \Delta_{\mathcal{H}})^{-1} f = [(\lambda I - \Delta)^{-1} E_{\mathcal{H}} f]_{\mathbb{R}_+^n}$. One may conclude following the lines of the proof of [Gau24b, Proposition 5.1], using Lemma 2.17. \square

LEMMA 2.18. — *Provided $k \in \llbracket 0, n \rrbracket$, $\mu \in (0, \pi)$, $\lambda \in \Sigma_\mu$, the following commutation identities hold,*

- (1) $\mathbb{P}(\lambda I - \Delta_{\mathcal{H},n})^{-1} f = (\lambda I - \Delta_{\mathcal{H},n})^{-1} \mathbb{P} f$, for all $f \in L^2(\mathbb{R}_+^n, \Lambda^k)$;
- (2) $d(\lambda I - \Delta_{\mathcal{H},n})^{-1} f = (\lambda I - \Delta_{\mathcal{H},n})^{-1} df$, for all $f \in D_2(d, \mathbb{R}_+^n, \Lambda^k)$;
- (3) $d^*(\lambda I - \Delta_{\mathcal{H},n})^{-1} f = (\lambda I - \Delta_{\mathcal{H},n})^{-1} d^* f$, for all $f \in D_2(d^*, \mathbb{R}_+^n, \Lambda^k)$.

Every above identities still hold replacing $(\mathbf{n}, \mathbb{P}, d, d^*)$ by $(\mathbf{t}, \mathbb{Q}, \delta^*, \delta)$.

Remark 2.19. — Lemma 2.18 does not depend on the domain $\Omega = \mathbb{R}_+^n$ since its proof only relies on the use of the sesquilinear form associated with the Hodge Laplacian.

Proof. —

Step 0: For $u \in D_2(D_n, \mathbb{R}_+^n, \Lambda^k)$, we have $\mathbb{P}u \in D_2(D_n, \mathbb{R}_+^n, \Lambda^k)$. Indeed, by definition $\mathbb{P}u \in N_2(d^*, \mathbb{R}_+^n, \Lambda^k)$. Hence, it remains to prove $\mathbb{P}u \in D_2(d, \mathbb{R}_+^n, \Lambda^k)$. For $\varphi \in C_c^\infty(\mathbb{R}_+^n, \Lambda)$, we have

$$\langle \mathbb{P}u, \delta\varphi \rangle_{\mathbb{R}_+^n} = \langle \mathbb{P}u, d^*\varphi \rangle_{\mathbb{R}_+^n} = \langle u, \mathbb{P}d^*\varphi \rangle_{\mathbb{R}_+^n} = \langle u, d^*\varphi \rangle_{\mathbb{R}_+^n} = \langle du, \varphi \rangle_{\mathbb{R}_+^n}.$$

Therefore, $d\mathbb{P}u = du$ in $\mathcal{D}'(\mathbb{R}_+^n, \Lambda)$, hence $d\mathbb{P}u \in L^2(\mathbb{R}_+^n, \Lambda)$.

Now, for all $u, v \in D_2(D_n, \mathbb{R}_+^n, \Lambda)$, since $d\mathbb{P}w = dw$ and $d^*\mathbb{P}w = 0$ for $w \in \{u, v\}$, we have

$$\begin{aligned} \mathbf{a}_{\mathcal{H},n}(\mathbb{P}u, v) &= \langle d\mathbb{P}u, dv \rangle_{\mathbb{R}_+^n} + \langle d^*\mathbb{P}u, d^*v \rangle_{\mathbb{R}_+^n} \\ &= \langle du, dv \rangle_{\mathbb{R}_+^n} \\ &= \langle du, d\mathbb{P}v \rangle_{\mathbb{R}_+^n} \\ &= \langle du, d\mathbb{P}v \rangle_{\mathbb{R}_+^n} + \langle d^*u, d^*\mathbb{P}v \rangle_{\mathbb{R}_+^n} \\ &= \mathbf{a}_{\mathcal{H},n}(u, \mathbb{P}v). \end{aligned}$$

Step 1: Let $\mu \in (0, \pi)$, $\lambda \in \Sigma_\mu$, and $f \in L^2(\mathbb{R}_+^n, \Lambda^k)$, we set $u := (\lambda I - \Delta_{\mathcal{H},n})^{-1} f$, then for all $v \in D_2(D_n, \mathbb{R}_+^n, \Lambda)$,

$$\begin{aligned} \lambda \langle \mathbb{P}u, v \rangle_{\mathbb{R}_+^n} + \mathfrak{a}_{\mathcal{H},n}(\mathbb{P}u, v) &= \lambda \langle u, \mathbb{P}v \rangle_{\mathbb{R}_+^n} + \mathfrak{a}_{\mathcal{H},n}(u, \mathbb{P}v) \\ &= \langle f, \mathbb{P}v \rangle_{\mathbb{R}_+^n} \\ &= \langle \mathbb{P}f, v \rangle_{\mathbb{R}_+^n}. \end{aligned}$$

Hence, by uniqueness of the solution to the resolvent problem in $L^2(\mathbb{R}_+^n, \Lambda)$, we deduce $\mathbb{P}u = (\lambda I - \Delta_{\mathcal{H},n})^{-1} \mathbb{P}f$.

Step 2: We use the same notations as the ones introduced in Step 1, but we assume that $f \in D_2(d, \mathbb{R}_+^n, \Lambda^k)$. For all $v \in D_2(\Delta_{\mathcal{H},n}, \mathbb{R}_+^n, \Lambda)$, since $d^2 = 0$, as well as $(d^*)^2 = 0$,

$$\begin{aligned} \langle df, v \rangle_{\mathbb{R}_+^n} &= \langle f, d^*v \rangle_{\mathbb{R}_+^n} \\ &= \lambda \langle u, d^*v \rangle_{\mathbb{R}_+^n} + \mathfrak{a}_{\mathcal{H},n}(u, d^*v) \\ &= \lambda \langle u, d^*v \rangle_{\mathbb{R}_+^n} + \langle du, dd^*v \rangle_{\mathbb{R}_+^n} + \langle d^*u, (d^*)^2v \rangle_{\mathbb{R}_+^n} \\ &= \lambda \langle du, v \rangle_{\mathbb{R}_+^n} + \langle d^*du, d^*v \rangle_{\mathbb{R}_+^n} \\ &= \lambda \langle du, v \rangle_{\mathbb{R}_+^n} + \langle ddu, dv \rangle_{\mathbb{R}_+^n} + \langle d^*du, d^*v \rangle_{\mathbb{R}_+^n} \\ &= \lambda \langle du, v \rangle_{\mathbb{R}_+^n} + \mathfrak{a}_{\mathcal{H},n}(du, v). \end{aligned}$$

Since the continuous embedding $D_2(\Delta_{\mathcal{H},n}, \mathbb{R}_+^n, \Lambda) \hookrightarrow D_2(D_n, \mathbb{R}_+^n, \Lambda)$ is dense, the equality above still holds for all $v \in D_2(D_n, \mathbb{R}_+^n, \Lambda)$. Hence, by uniqueness of the solution to the resolvent problem in $L^2(\mathbb{R}_+^n, \Lambda)$, we obtain $du = (\lambda I - \Delta_{\mathcal{H},n})^{-1} df$. The proof ends here since all remaining results can be proven similarly. \square

LEMMA 2.20. — Let $k \in \llbracket 0, n \rrbracket$, following operators

$$\begin{aligned} d(-\Delta_{\mathcal{H},n})^{-\frac{1}{2}} : L^2(\mathbb{R}_+^n, \Lambda^k) &\longrightarrow N_2(d, \mathbb{R}_+^n, \Lambda^{k+1}); \\ d^*(-\Delta_{\mathcal{H},n})^{-\frac{1}{2}} : L^2(\mathbb{R}_+^n, \Lambda^k) &\longrightarrow N_2(d^*, \mathbb{R}_+^n, \Lambda^{k-1}); \end{aligned}$$

are well-defined bounded linear operators on $L^2(\mathbb{R}_+^n, \Lambda)$, and are each-other's adjoint. Moreover, for all $u \in L^2(\mathbb{R}_+^n, \Lambda^k)$, we have

$$\left. \begin{aligned} &\|d(-\Delta_{\mathcal{H},n})^{-\frac{1}{2}}u - d(\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}}u\|_{L^2(\mathbb{R}_+^n)} \\ &\|d^*(-\Delta_{\mathcal{H},n})^{-\frac{1}{2}}u - d^*(\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}}u\|_{L^2(\mathbb{R}_+^n)} \end{aligned} \right\} \xrightarrow{\lambda \rightarrow 0_+} 0.$$

Everything still holds replacing (\mathfrak{n}, d, d^*) by $(\mathfrak{t}, \delta^*, \delta)$.

Proof. — We prove the $L^2(\mathbb{R}_+^n, \Lambda)$ -boundedness of $d(-\Delta_{\mathcal{H},n})^{-\frac{1}{2}}$ and compute its adjoint.

• We recall that the Hodge Laplacian is 0-sectorial and injective over $L^2(\mathbb{R}_+^n, \Lambda^k)$, so that by Proposition 1.4,

$$\overline{D_2\left((-\Delta_{\mathcal{H},n})^{1/2}, \mathbb{R}_+^n, \Lambda^k\right) \cap R_2\left((-\Delta_{\mathcal{H},n})^{1/2}, \mathbb{R}_+^n, \Lambda^k\right)}^{\|\cdot\|_{L^2(\mathbb{R}_+^n)}} = L^2\left(\mathbb{R}_+^n, \Lambda^k\right).$$

We use the bounded holomorphic functional calculus of D_n on $L^2(\mathbb{R}_+^n, \Lambda)$ provided by Proposition 2.13. By means of $f_\lambda : z \mapsto \frac{z}{\sqrt{\lambda+z^2}}$, and the boundedness of \mathbb{P} , we have, for all $\lambda \geq 0$, and all $u \in L^2(\mathbb{R}_+^n, \Lambda^k)$,

$$\begin{aligned} \|f_\lambda(D_n)u\|_{L^2(\mathbb{R}_+^n)}^2 &= \|D_n(\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}}u\|_{L^2(\mathbb{R}_+^n)}^2 \\ &= \|d(\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}}u\|_{L^2(\mathbb{R}_+^n)}^2 + \|d^*(\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}}u\|_{L^2(\mathbb{R}_+^n)}^2 \\ &\leq \|u\|_{L^2(\mathbb{R}_+^n)}^2. \end{aligned}$$

• We did the abuse of notation $f_0(D_n) = D_n(-\Delta_{\mathcal{H},n})^{-\frac{1}{2}}$. We clarify this point and show that for all $u \in L^2(\mathbb{R}_+^n, \Lambda^k)$,

$$\left. \begin{aligned} &\|d(-\Delta_{\mathcal{H},n})^{-\frac{1}{2}}u - d(\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}}u\|_{L^2(\mathbb{R}_+^n)} \\ &\|d^*(-\Delta_{\mathcal{H},n})^{-\frac{1}{2}}u - d^*(\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}}u\|_{L^2(\mathbb{R}_+^n)} \end{aligned} \right\} \xrightarrow{\lambda \rightarrow 0_+} 0.$$

We start with an element $v \in D_2((-\Delta_{\mathcal{H},n})^{1/2}, \Lambda^k) \cap R_2((-\Delta_{\mathcal{H},n})^{1/2}, \Lambda^k)$, so one can write $v = (-\Delta_{\mathcal{H},n})^{1/2}w$ for $w \in D_2(\Delta_{\mathcal{H},n}, \Lambda^k)$, and we obtain by Lemma 2.18:

$$\begin{aligned} &\|d(\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}}v - dw\|_{L^2(\mathbb{R}_+^n)}^2 + \|d^*(\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}}v - d^*w\|_{L^2(\mathbb{R}_+^n)}^2 \\ &= \|(-\Delta_{\mathcal{H},n})^{\frac{1}{2}}(\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}}dw - dw\|_{L^2(\mathbb{R}_+^n)}^2 \\ &\quad + \|(-\Delta_{\mathcal{H},n})^{\frac{1}{2}}(\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}}d^*w - d^*w\|_{L^2(\mathbb{R}_+^n)}^2 \\ &= \|(-\Delta_{\mathcal{H},n})^{\frac{1}{2}}(\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}}D_nw - D_nw\|_{L^2(\mathbb{R}_+^n)}^2 \\ &= \|(-\Delta_{\mathcal{H},n})^{\frac{1}{2}}(\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}}(-\Delta_{\mathcal{H},n})^{\frac{1}{2}}w - (-\Delta_{\mathcal{H},n})^{\frac{1}{2}}w\|_{L^2(\mathbb{R}_+^n)}^2 \\ &= \|(-\Delta_{\mathcal{H},n})^{\frac{1}{2}}(\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}}v - v\|_{L^2(\mathbb{R}_+^n)}^2 \end{aligned}$$

But thanks to [Haa06, Theorem 5.2.6], originally due to an idea of McIntosh [McI86], one has the representation formulas

$$v = \frac{1}{\sqrt{\pi}} \int_0^{+\infty} (-\Delta_{\mathcal{H},n})^{\frac{1}{2}} e^{t\Delta_{\mathcal{H},n}} v \frac{dt}{\sqrt{t}},$$

and

$$(-\Delta_{\mathcal{H},n})^{\frac{1}{2}}(\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}}v = \frac{1}{\sqrt{\pi}} \int_0^{+\infty} (-\Delta_{\mathcal{H},n})^{\frac{1}{2}} e^{-t\lambda} e^{t\Delta_{\mathcal{H},n}} v \frac{dt}{\sqrt{t}}.$$

Since $v \in D_2((-\Delta_{\mathcal{H},n})^{1/2}, \Lambda^k) \cap R_2((-\Delta_{\mathcal{H},n})^{1/2}, \Lambda^k)$ the integrals are absolutely convergent, and we can also apply the Dominated Convergence Theorem, so that

$$(2.12) \quad \begin{aligned} & \left\| d(\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}} v - dw \right\|_{L^2(\mathbb{R}_+^n)}^2 + \left\| d^*(\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}} v - d^*w \right\|_{L^2(\mathbb{R}_+^n)}^2 \\ &= \left\| (-\Delta_{\mathcal{H},n})^{\frac{1}{2}} (\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}} v - v \right\|_{L^2(\mathbb{R}_+^n)}^2 \xrightarrow{\lambda \rightarrow 0_+} 0. \end{aligned}$$

Now, recalling that the Hodge Laplacian is injective, we can write

$$D_n w = D_n (-\Delta_{\mathcal{H},n})^{-\frac{1}{2}} v = f_0(D_n)v,$$

so we obtain

$$\|f_0(D_n)v\|_{L^2(\mathbb{R}_+^n)}^2 = \left\| d(-\Delta_{\mathcal{H},n})^{-\frac{1}{2}} v \right\|_{L^2(\mathbb{R}_+^n)}^2 + \left\| d^*(-\Delta_{\mathcal{H},n})^{-\frac{1}{2}} v \right\|_{L^2(\mathbb{R}_+^n)}^2 \leq \|v\|_{L^2(\mathbb{R}_+^n)}^2.$$

Since $D_2((-\Delta_{\mathcal{H},n})^{1/2}, \Lambda^k) \cap R_2((-\Delta_{\mathcal{H},n})^{1/2}, \Lambda^k)$ is dense in $L^2(\mathbb{R}_+^n, \Lambda^k)$, everything coincides by uniqueness of continuous extensions.

It remains to relax the convergence to all $u \in L^2(\mathbb{R}_+^n, \Lambda^k)$. For $u \in L^2(\mathbb{R}_+^n, \Lambda^k)$, $\varepsilon > 0$, there exists $\tilde{u} \in D_2((-\Delta_{\mathcal{H},n})^{1/2}, \Lambda^k) \cap R_2((-\Delta_{\mathcal{H},n})^{1/2}, \Lambda^k)$ such that

$$\|u - \tilde{u}\|_{L^2(\mathbb{R}_+^n)} < \varepsilon.$$

We deduce for $\lambda > 0$,

$$\begin{aligned} & \left\| d(-\Delta_{\mathcal{H},n})^{-\frac{1}{2}} u - d(\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}} u \right\|_{L^2(\mathbb{R}_+^n)} \\ & \leq \left\| d(-\Delta_{\mathcal{H},n})^{-\frac{1}{2}} u - d(-\Delta_{\mathcal{H},n})^{-\frac{1}{2}} \tilde{u} \right\|_{L^2(\mathbb{R}_+^n)} \\ & \quad + \left\| d(-\Delta_{\mathcal{H},n})^{-\frac{1}{2}} \tilde{u} - d(\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}} \tilde{u} \right\|_{L^2(\mathbb{R}_+^n)} \\ & \quad + \left\| d(\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}} \tilde{u} - d(\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}} u \right\|_{L^2(\mathbb{R}_+^n)} \\ & < 2\varepsilon + \left\| d(-\Delta_{\mathcal{H},n})^{-\frac{1}{2}} \tilde{u} - d(\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}} \tilde{u} \right\|_{L^2(\mathbb{R}_+^n)}. \end{aligned}$$

So that, for λ large enough, since $\tilde{u} \in D_2((-\Delta_{\mathcal{H},n})^{1/2}, \Lambda^k) \cap R_2((-\Delta_{\mathcal{H},n})^{1/2}, \Lambda^k)$, we use (2.12) to reach the inequality

$$\left\| d(-\Delta_{\mathcal{H},n})^{-\frac{1}{2}} u - d(\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}} u \right\|_{L^2(\mathbb{R}_+^n)} < 3\varepsilon.$$

We proceed similarly for $d^*(-\Delta_{\mathcal{H},n})^{-\frac{1}{2}}$.

• Now, we compute the adjoint. The adjoint of $d(\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}}$, provided $\lambda > 0$, is $(\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}} d^* = d^*(\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}}$ up to a dense subset of $L^2(\mathbb{R}_+^n, \Lambda)$ (here $D_2(d^*, \mathbb{R}_+^n, \Lambda)$), thanks to Lemma 2.18. By previous steps, we can pass to the limit as λ goes to 0 in the L^2 inner product yielding the identity

$$\left(d(-\Delta_{\mathcal{H},n})^{-\frac{1}{2}} \right)^* = d^*(-\Delta_{\mathcal{H},n})^{-\frac{1}{2}}. \quad \square$$

We can summarize with the next theorem.

THEOREM 2.21. — *Let $k \in \llbracket 0, n \rrbracket$, the following assertions are true*

(1) *The following equality holds*

$$N_2(d, \mathbb{R}_+^n, \Lambda^k) = \overline{R_2(d, \mathbb{R}_+^n, \Lambda^k)}^{\|\cdot\|_{L^2(\mathbb{R}_+^n)}},$$

and still holds replacing d by d^ .*

(2) *The (generalized) Helmholtz–Leray projector*

$$\mathbb{P} : L^2(\mathbb{R}_+^n, \Lambda^k) \longrightarrow N^2(d^*, \mathbb{R}_+^n, \Lambda^k)$$

satisfies the identity

$$\mathbb{P} = I - d(-\Delta_{\mathcal{H},n})^{-\frac{1}{2}} d^*(-\Delta_{\mathcal{H},n})^{-\frac{1}{2}}.$$

(3) *The following Hodge decomposition holds*

$$L^2(\mathbb{R}_+^n, \Lambda^k) = N^2(d^*, \mathbb{R}_+^n, \Lambda^k) \oplus N^2(d, \mathbb{R}_+^n, \Lambda^k).$$

Moreover, the result remains true if we replace $(\mathfrak{n}, d, d^, \mathbb{P})$ by $(\mathfrak{t}, \delta^*, \delta, \mathbb{Q})$.*

Remark 2.22. — The Theorem 2.21 and the whole construction of this section mainly depends on the injectivity of the Laplacian: the construction is done via resolvent approximation, abstract functional calculus provided by the Hilbertian structure of $L^2(\mathbb{R}_+^n, \Lambda)$ and the self-adjointness of the Laplacian. Therefore, a such construction does not depend on the open set $\Omega = \mathbb{R}_+^n$.

To be more precise, the above Theorem 2.21 should remain true for all open sets Ω , say at least Lipschitz, that admit no non-zero harmonic forms. In the case of a bounded domain: the theorem remains true whenever all its Betti numbers vanish.

Proof. —

Step 1: Identity for \mathbb{P} . — From boundedness of the operators from Lemma 2.20, we deduce that the new operator $\overline{\mathbb{P}}$ defined for all $f \in L^2(\mathbb{R}_+^n, \Lambda)$ by

$$\overline{\mathbb{P}}f := f - d(-\Delta_{\mathcal{H},n})^{-\frac{1}{2}} d^*(-\Delta_{\mathcal{H},n})^{-\frac{1}{2}} f,$$

is well-defined and bounded on $L^2(\mathbb{R}_+^n, \Lambda)$. We are going to check that $\overline{\mathbb{P}}$ is an orthogonal projector, hence, firstly, a projector. By Proposition 1.4 and Lemma 2.20, we have

$$\begin{aligned} \overline{\mathbb{P}}^2 f &= \overline{\mathbb{P}}f - d(-\Delta_{\mathcal{H},n})^{-\frac{1}{2}} d^*(-\Delta_{\mathcal{H},n})^{-\frac{1}{2}} \overline{\mathbb{P}}f \\ &= \overline{\mathbb{P}}f - d(-\Delta_{\mathcal{H},n})^{-\frac{1}{2}} d^*(-\Delta_{\mathcal{H},n})^{-\frac{1}{2}} f + \left[d(-\Delta_{\mathcal{H},n})^{-\frac{1}{2}} d^*(-\Delta_{\mathcal{H},n})^{-\frac{1}{2}} \right]^2 f \\ &= \lim_{\lambda \rightarrow 0} \left(\overline{\mathbb{P}}f - d^* d (\lambda I - \Delta_{\mathcal{H},n})^{-1} f + \left[d^* d (\lambda I - \Delta_{\mathcal{H},n})^{-1} \right]^2 f \right) \\ &= \lim_{\lambda \rightarrow 0} \left(\overline{\mathbb{P}}f - \lambda d^* d (\lambda I - \Delta_{\mathcal{H},n})^{-2} f \right) \\ &= \overline{\mathbb{P}}f. \end{aligned}$$

By construction and by Lemma 2.20, $\overline{\mathbb{P}}$ is self-adjoint, hence orthogonal.

For all $f \in D_2(d^*, \mathbb{R}_+^n, \Lambda)$, by Proposition 1.4, Lemma 2.18 and Lemma 2.20, since $(d^*)^2 = 0$,

$$\begin{aligned} d^* \bar{\mathbb{P}} f &= \lim_{\lambda \rightarrow 0} \left(d^* f - d^* d (\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}} d^* (\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}} f \right) \\ &= \lim_{\lambda \rightarrow 0} \left(d^* f + \Delta_{\mathcal{H},n} (\lambda I - \Delta_{\mathcal{H},n})^{-1} d^* f \right) \\ &= (d^* f - d^* f) = 0. \end{aligned}$$

Thus, since the embedding $D_2(d^*, \mathbb{R}_+^n, \Lambda) \hookrightarrow L^2(\mathbb{R}_+^n, \Lambda)$ is dense, it follows that

$$R_2(\bar{\mathbb{P}}, \mathbb{R}_+^n, \Lambda) \subset N_2(d^*, \mathbb{R}_+^n, \Lambda).$$

For all $f \in N_2(d^*, \mathbb{R}_+^n, \Lambda)$,

$$\begin{aligned} \bar{\mathbb{P}} f &= \lim_{\lambda \rightarrow 0} \left(f - d (\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}} d^* (\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}} f \right) \\ &= \lim_{\lambda \rightarrow 0} \left(f - d (\lambda I - \Delta_{\mathcal{H},n})^{-1} d^* f \right) \\ &= (f + 0) = f. \end{aligned}$$

Hence, $\mathbb{P}|_{N_2(d^*, \mathbb{R}_+^n, \Lambda)} = I$.

By construction, we also have

$$R_2(I - \bar{\mathbb{P}}, \mathbb{R}_+^n, \Lambda) = R_2(d(-\Delta_{\mathcal{H},n})^{-\frac{1}{2}} d^* (-\Delta_{\mathcal{H},n})^{-\frac{1}{2}}, \mathbb{R}_+^n, \Lambda) \subset \overline{R_2(d, \mathbb{R}_+^n, \Lambda)};$$

so that by uniqueness of the orthogonal projection on $N_2(d^*, \mathbb{R}_+^n, \Lambda)$, $\bar{\mathbb{P}} = \mathbb{P}$.

Step 2: We notice first that the inclusion $\overline{R_2(d, \mathbb{R}_+^n, \Lambda^k)}^{\|\cdot\|_{L^2(\mathbb{R}_+^n)}} \subset N_2(d, \mathbb{R}_+^n, \Lambda^k)$ is true.

Now, for the reverse inclusion let $f \in N_2(d, \mathbb{R}_+^n, \Lambda^k)$, we have

$$f = \lim_{\lambda \rightarrow 0} -\Delta_{\mathcal{H},n} (\lambda I - \Delta_{\mathcal{H},n})^{-1} f = \lim_{\lambda \rightarrow 0} d d^* (\lambda I - \Delta_{\mathcal{H},n})^{-1} f (= [I - \mathbb{P}]f).$$

By construction, for all $\lambda > 0$, we have $d d^* (\lambda I - \Delta_{\mathcal{H},n})^{-1} f \in R_2(d, \mathbb{R}_+^n, \Lambda^k)$, so that the reverse inclusion $N_2(d, \mathbb{R}_+^n, \Lambda^k) \subset \overline{R_2(d, \mathbb{R}_+^n, \Lambda^k)}^{\|\cdot\|_{L^2(\mathbb{R}_+^n)}}$ holds. \square

2.3.2. $\dot{H}^{s,p}$ and $\dot{B}_{p,q}^s$ -theory for Hodge Laplacians and the Hodge decomposition

We start this new subsection claiming about closedness of the exterior and interior derivatives with and without 0 boundary conditions. The two following lemmas are straightforward.

LEMMA 2.23. — *Let $p \in (1, +\infty)$, $s \in (-1 + 1/p, 1/p)$, $k \in \llbracket 0, n \rrbracket$. With the same notations as in Lemma 2.11, the operators*

$$\left(\dot{D}_p^s(d, \mathbb{R}_+^n, \Lambda^k), d \right) \quad \text{and} \quad \left(\dot{D}_p^s(\delta, \mathbb{R}_+^n, \Lambda^k), \delta \right)$$

are densely defined closed operators on $\dot{H}^{s,p}(\mathbb{R}_+^n, \Lambda^k)$.

Moreover,

- the result still holds replacing $(\dot{H}^{s,p}, \dot{D}_p^s)$ by either $(H^{s,p}, D_p^s)$, $(B_{p,q}^s, D_{p,q}^s)$ or $(\dot{B}_{p,q}^s, \dot{D}_{p,q}^s)$, with $q \in [1, +\infty)$;

- in case of $(B_{p,\infty}^s, D_{p,\infty}^s)$ and $(\dot{B}_{p,\infty}^s, \dot{D}_{p,\infty}^s)$ above operators are only weak* densely defined, strongly closed operators;
- all the above results remain true exchanging the roles of d and δ .

LEMMA 2.24. — Let $p \in (1, +\infty)$, $s \in (-1 + 1/p, 1/p)$. With the same notations as in Lemma 2.11, the dual operator of $(\dot{D}_p^s(d, \mathbb{R}_+^n, \Lambda), d)$ on $\dot{H}^{s,p}(\mathbb{R}_+^n, \Lambda)$, is

$$(\dot{D}_{p'}^{-s}(d^*, \mathbb{R}_+^n, \Lambda), d^*) = (\dot{D}_{p'}^{-s}(\underline{\delta}, \mathbb{R}_+^n, \Lambda), \delta)$$

as an operator on $\dot{H}^{-s,p'}(\mathbb{R}_+^n, \Lambda)$.

Moreover,

- the result still holds replacing $(\dot{H}^{s,p}, \dot{D}_p^s, \dot{H}^{-s,p'}, \dot{d}_{p'}^{-s})$ by $(\dot{B}_{p,q}^s, \dot{D}_{p,q}^s, \dot{B}_{p',q'}^{-s}, \dot{d}_{p',q'}^{-s})$ with $q \in [1, +\infty)$;
- we may replace $(\dot{D}, \dot{H}, \dot{B})$ by (D, H, B) ;
- all the above results remain true exchanging the roles of d and δ .

Remark 2.25. — Notice that talking about $D_p^s(\underline{d}, \mathbb{R}_+^n, \Lambda)$ in above lemmas, with respect to notation introduced in Lemma 2.11, in particular the involved 0-boundary condition, actually makes sense, thanks to Theorem A.2.

Before we start our investigation of Hodge Laplacians and the Hodge decomposition, we need to show the closedness of Hodge–Dirac operators. In order to verify such a property, the next result will be of paramount importance, to reproduce the behavior obtained in the L^2 setting on other scales of function spaces. We mention that many results presented here will strongly depend on the fact that the considered open set is \mathbb{R}_+^n (mainly Lemma 2.26, and point (ii) of Theorem 2.29 which are widely used to construct other results of the present section).

The proof of the next lemma is identical to the one of Lemma 2.17.

LEMMA 2.26. — Let $p \in (1, +\infty)$, $s \in (-1 + 1/p, 1/p)$, $k \in \llbracket 0, n \rrbracket$. For all $u \in \dot{D}_p^s(d, \mathbb{R}_+^n, \Lambda^k)$ (resp. $\dot{D}_p^s(d^*, \mathbb{R}_+^n, \Lambda^k)$) we have

$$E_{\mathcal{H},n}u \in \dot{D}_p^s(d, \mathbb{R}^n, \Lambda^k) \quad (\text{resp. } \dot{D}_p^s(\delta, \mathbb{R}^n, \Lambda^k))$$

with formulas

$$dE_{\mathcal{H},n}u = E_{\mathcal{H},n}du \quad (\text{resp. } \delta E_{\mathcal{H},n}u = E_{\mathcal{H},n}d^*u).$$

Moreover,

- the result still holds replacing \dot{D}_p^s by $\dot{D}_{p,q}^s$ with $q \in [1, +\infty)$;
- we may replace \dot{D} by D ;
- all the above results remain true exchanging the roles of d and δ , and replacing n by t .

PROPOSITION 2.27. — Let $p \in (1, +\infty)$, $s \in (-1 + 1/p, 1/p)$, $k \in \llbracket 0, n \rrbracket$. The Hodge–Dirac operator

$$(\dot{D}_p^s(D_n, \mathbb{R}_+^n, \Lambda^k), D_n) = (\dot{D}_p^s(d, \mathbb{R}_+^n, \Lambda^k) \cap \dot{D}_p^s(d^*, \mathbb{R}_+^n, \Lambda^k), d + d^*)$$

is a densely defined closed operator on $\dot{H}^{s,p}(\mathbb{R}_+^n, \Lambda^k)$.

Moreover,

- the result still holds replacing $(\dot{H}^{s,p}, \dot{D}_p^s)$ by either $(H^{s,p}, D_p^s)$, $(B_{p,q}^s, D_{p,q}^s)$ or $(\dot{B}_{p,q}^s, \dot{D}_{p,q}^s)$, with $q \in [1, +\infty)$;
- in case of $(B_{p,\infty}^s, D_{p,\infty}^s)$ and $(\dot{B}_{p,\infty}^s, \dot{D}_{p,\infty}^s)$ above Hodge–Dirac operator is only weak* densely defined, and strongly closed;
- all above results remain true replacing (\mathbf{n}, d) by (\mathbf{t}, δ) .

Proof. — Let $(u_j)_{j \in \mathbb{N}} \subset \dot{D}_p^s(d, \mathbb{R}_+^n, \Lambda^k) \cap \dot{D}_p^s(d^*, \mathbb{R}_+^n, \Lambda^k)$, and $(u, v) \in \dot{H}^{s,p}(\mathbb{R}_+^n, \Lambda^k) \times \dot{H}^{s,p}(\mathbb{R}_+^n, \Lambda)$ satisfying

$$u_j \xrightarrow{j \rightarrow +\infty} u \quad \text{and} \quad D_{\mathbf{n}} u_j \xrightarrow{j \rightarrow +\infty} v \quad \text{in} \quad \dot{H}^{s,p}(\mathbb{R}_+^n, \Lambda).$$

We set for all $j \in \mathbb{N}$, $U_j := E_{\mathcal{H}, \mathbf{n}} u_j$, $U := E_{\mathcal{H}, \mathbf{n}} u$. By Lemma 2.26, we have for all $j \in \mathbb{Z}$ $U_j \in \dot{D}_p^s(d, \mathbb{R}^n, \Lambda^k) \cap \dot{D}_p^s(\delta, \mathbb{R}^n, \Lambda^k)$

$$DU_j = E_{\mathcal{H}, \mathbf{n}} D_{\mathbf{n}} u_j.$$

We also have,

$$DU_j \xrightarrow{j \rightarrow +\infty} V := E_{\mathcal{H}, \mathbf{n}} v \quad \text{in} \quad \dot{H}^{s,p}(\mathbb{R}^n, \Lambda).$$

By the Hodge decomposition on \mathbb{R}^n , check Theorem 2.9, there exists a unique couple $(V_0, V_1) \in \dot{R}_p^s(d, \mathbb{R}^n, \Lambda) \times \dot{N}_p^s(\delta, \mathbb{R}^n, \Lambda)$ such that $V = V_0 + V_1$. Since U_j goes to U in $\dot{H}^{s,p}(\mathbb{R}^n, \Lambda^k)$, by continuity of involved projectors, and uniqueness of decomposition, it follows that

$$dU_j \xrightarrow{j \rightarrow +\infty} V_0 \quad \text{and} \quad \delta U_j \xrightarrow{j \rightarrow +\infty} V_1 \quad \text{in} \quad \dot{H}^{s,p}(\mathbb{R}^n, \Lambda).$$

In particular, if we set $v_\ell := V_\ell|_{\mathbb{R}_+^n}$ for $\ell \in \{0, 1\}$, we necessarily have by restriction

$$du_j \xrightarrow{j \rightarrow +\infty} v_0 \quad \text{and} \quad \delta u_j \xrightarrow{j \rightarrow +\infty} v_1 \quad \text{in} \quad \dot{H}^{s,p}(\mathbb{R}_+^n, \Lambda).$$

But $(u_j)_{j \in \mathbb{N}}$ converge to u in $\dot{H}^{s,p}(\mathbb{R}_+^n, \Lambda^k)$, so in particular in distributional sense. Thus, necessarily $(v_0, v_1) = (du, \delta u)$ and $v = Du$. By continuity of trace provided by Theorem A.1, we also have $\nu \lrcorner u|_{\partial \mathbb{R}_+^n} = 0$, i.e. $(\dot{D}_p^s(D_{\mathbf{n}}, \mathbb{R}_+^n, \Lambda^k), D_{\mathbf{n}})$ is a closed operator on $\dot{H}^{s,p}(\mathbb{R}_+^n, \Lambda^k)$. The proof ends here, since one can reproduce all above arguments for $(\dot{D}_p^s(D_{\mathbf{t}}, \mathbb{R}_+^n, \Lambda^k), D_{\mathbf{t}})$, and also for all other kind of function spaces. \square

The next result about closedness of Hodge Laplacian admits a similar proof.

PROPOSITION 2.28. — *Let $p \in (1, +\infty)$, $s \in (-1 + 1/p, 1/p)$, $k \in \llbracket 0, n \rrbracket$. The Hodge Laplacian*

$$\left(\dot{D}_p^s(\Delta_{\mathcal{H}, \mathbf{n}}, \mathbb{R}_+^n, \Lambda^k), -\Delta_{\mathcal{H}, \mathbf{n}} \right) = \left(\dot{D}_p^s(D_{\mathbf{n}}^2, \mathbb{R}_+^n, \Lambda^k), D_{\mathbf{n}}^2 \right)$$

is a densely defined closed injective operator on $\dot{H}^{s,p}(\mathbb{R}_+^n, \Lambda^k)$. The following formula holds for all $u \in \dot{D}_p^s(\Delta_{\mathcal{H}, \mathbf{n}}, \mathbb{R}_+^n, \Lambda^k)$,

$$-\Delta E_{\mathcal{H}, \mathbf{n}} u = E_{\mathcal{H}, \mathbf{n}} [-\Delta_{\mathcal{H}, \mathbf{n}} u].$$

Moreover,

- the result still holds replacing $(\dot{H}^{s,p}, \dot{D}_p^s)$ by either $(H^{s,p}, D_p^s)$, $(B_{p,q}^s, D_{p,q}^s)$ or $(\dot{B}_{p,q}^s, \dot{D}_{p,q}^s)$, with $q \in [1, +\infty)$;

- in case of $(B_{p,\infty}^s, D_{p,\infty}^s)$ and $(\dot{B}_{p,\infty}^s, \dot{D}_{p,\infty}^s)$ the Hodge Laplacian is only weak* densely defined, and strongly closed;
- all above results remain true replacing \mathfrak{n} by \mathfrak{t} .

From there, the whole context has been established in order to be able to claim the next theorem.

THEOREM 2.29. — *Let $p \in (1, +\infty)$, $q \in [1, +\infty]$, $s \in (-1 + 1/p, 1/p)$, and $k \in \llbracket 0, n \rrbracket$.*

- (i) *For $\mu \in [0, \pi)$, $\lambda \in \Sigma_\mu$, if $f \in \dot{H}^{s,p}(\mathbb{R}_+^n, \Lambda^k)$ then the following resolvent problem*

$$\lambda u - \Delta_{\mathcal{H}} u = f \text{ in } \mathbb{R}_+^n,$$

admits a unique solution $u \in \dot{D}_p^s(\Delta_{\mathcal{H}}, \mathbb{R}_+^n, \Lambda^k) \subset [\dot{H}^{s,p} \cap \dot{H}^{s+2,p}](\mathbb{R}_+^n, \Lambda^k)$ with estimates,

$$\begin{aligned} |\lambda| \|u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} + |\lambda|^{\frac{1}{2}} \|\nabla u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} + \|\nabla^2 u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} &\lesssim_{p,n,s,\mu} \|f\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)}, \\ |\lambda|^{\frac{1}{2}} \|(d + \delta)u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} + \|d\delta u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} + \|\delta d u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} &\lesssim_{p,n,s,\mu} \|f\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)}. \end{aligned}$$

In particular, $\lambda I - \Delta_{\mathcal{H}} : \dot{D}_p^s(\Delta_{\mathcal{H}}, \mathbb{R}_+^n, \Lambda^k) \longrightarrow \dot{H}^{s,p}(\mathbb{R}_+^n, \Lambda^k)$ is an isomorphism of Banach spaces.

Furthermore, the result still holds replacing $(\dot{H}^{s,p}, \dot{H}^{s,p} \cap \dot{H}^{s+2,p}, \dot{D}_p^s)$ by $(H^{s,p}, H^{s+2,p}, D_p^s)$, $(\dot{B}_{p,q}^s, \dot{B}_{p,q}^s \cap \dot{B}_{p,q}^{s+2}, \dot{D}_{p,q}^s)$, or even by $(B_{p,q}^s, B_{p,q}^{s+2}, D_{p,q}^s)$.

- (ii) *For any $\mu \in (0, \pi)$, the operator $-\Delta_{\mathcal{H}}$ admits a $\mathbf{H}^\infty(\Sigma_\mu)$ -functional calculus on function spaces: $\dot{H}^{s,p}(\mathbb{R}_+^n, \Lambda^k)$, $\dot{B}_{p,q}^s(\mathbb{R}_+^n, \Lambda^k)$, $H^{s,p}(\mathbb{R}_+^n, \Lambda^k)$ and $B_{p,q}^s(\mathbb{R}_+^n, \Lambda^k)$.*

Moreover, the following resolvent identity holds on any previously mentioned function spaces,

$$E_{\mathcal{H}}(\lambda I - \Delta_{\mathcal{H}})^{-1} = (\lambda I - \Delta)^{-1} E_{\mathcal{H}}.$$

Proof. — For $\mu \in [0, \pi)$, $\lambda \in \Sigma_\mu$, if $f \in \dot{H}^{s,p}(\mathbb{R}_+^n, \Lambda^k)$, we have by Theorem 2.7 and (2.11):

$$(\lambda I - \Delta)^{-1} E_{\mathcal{H}} f \in [\dot{H}^{s,p} \cap \dot{H}^{s+2,p}](\mathbb{R}^n, \Lambda^k).$$

Thus, the definition of function spaces by restriction yields $u := [(\lambda I - \Delta)^{-1} E_{\mathcal{H}} f]_{|\mathbb{R}_+^n} \in [\dot{H}^{s,p} \cap \dot{H}^{s+2,p}](\mathbb{R}_+^n, \Lambda^k)$ and it satisfies

$$\lambda u - \Delta u = f \text{ in } \mathbb{R}_+^n,$$

so that by the definition of function spaces by restriction, Theorem 2.7, and the boundedness properties of $E_{\mathcal{H}}$ (2.11), we also have the estimates

$$\begin{aligned} |\lambda| \|u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} + |\lambda|^{\frac{1}{2}} \|\nabla u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} + \|\nabla^2 u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} &\lesssim_{p,n,s,\mu} \|f\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)}, \\ |\lambda|^{\frac{1}{2}} \|(d + \delta)u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} + \|d\delta u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} + \|\delta d u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} &\lesssim_{p,n,s,\mu} \|f\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)}. \end{aligned}$$

By density of $[L^2 \cap \dot{H}^{s,p}](\mathbb{R}_+^n, \Lambda^k)$ in $\dot{H}^{s,p}(\mathbb{R}_+^n, \Lambda^k)$, one may use Theorem 2.15 and continuity of traces provided by Theorem A.2 to show that necessarily $\nu \lrcorner u|_{\partial\mathbb{R}_+^n} = 0$ and $\nu \lrcorner du|_{\partial\mathbb{R}_+^n} = 0$ (or resp. $\nu \wedge u|_{\partial\mathbb{R}_+^n} = 0$ and $\nu \wedge \delta u|_{\partial\mathbb{R}_+^n} = 0$). Hence

$$u \in \dot{D}_p^s(\Delta_{\mathcal{H}}, \mathbb{R}_+^n, \Lambda^k).$$

Now assume $v \in \dot{D}_p^s(\Delta_{\mathcal{H}}, \mathbb{R}_+^n, \Lambda^k)$ satisfies

$$\lambda v - \Delta_{\mathcal{H}}v = f \text{ in } \mathbb{R}_+^n.$$

We apply Proposition 2.28 to claim that $V := E_{\mathcal{H}}v$ must satisfy

$$\lambda V - \Delta V = E_{\mathcal{H}}f \text{ in } \mathbb{R}^n.$$

Thus, necessarily $E_{\mathcal{H}}(\lambda I - \Delta_{\mathcal{H}})^{-1}f = (\lambda I - \Delta)^{-1}E_{\mathcal{H}}f$.

This resolvent identity leads to the construction of bounded $(\mathbf{H}^\infty(\Sigma_\mu)$ -)holomorphic functional calculus, given by the following identity for all $\Psi \in \mathbf{H}^\infty(\Sigma_\mu)$, $\mu \in (0, \pi)$:

$$E_{\mathcal{H}}\Psi(-\Delta_{\mathcal{H}}) = \Psi(-\Delta)E_{\mathcal{H}}.$$

The result for homogeneous Besov spaces $\dot{B}_{p,q}^s$, $q < +\infty$, and other similar inhomogeneous function spaces may be achieved in a similar manner. The case of inhomogeneous and homogeneous Besov spaces with $q = +\infty$ follows from real interpolation. □

The goal for now is to prove the Hodge decomposition. The idea is to prove that the representation formula of \mathbb{P} (resp. \mathbb{Q}) proved in Lemma 2.20 still makes sense on $\dot{H}^{s,p}(\mathbb{R}_+^n, \Lambda)$, $\dot{B}_{p,q}^s(\mathbb{R}_+^n, \Lambda)$, and their inhomogeneous counterparts. To do so, we adapt Lemma 2.18 in the present setting.

LEMMA 2.30. — *Let $p \in (1, +\infty)$, $s \in (-1 + 1/p, 1/p)$, $\mu \in [0, \pi)$, $\lambda \in \Sigma_\mu$, $t \geq 0$, $k \in \llbracket 0, n \rrbracket$. The following commutation identities hold,*

- (1) $d(\lambda I - \Delta_{\mathcal{H},n})^{-1}f = (\lambda I - \Delta_{\mathcal{H},n})^{-1}df$, for all $f \in \dot{D}_p^s(d, \mathbb{R}_+^n, \Lambda^k)$;
- (2) $d^*(\lambda I - \Delta_{\mathcal{H},n})^{-1}f = (\lambda I - \Delta_{\mathcal{H},n})^{-1}d^*f$, for all $f \in \dot{D}_p^s(d^*, \mathbb{R}_+^n, \Lambda^k)$;
- (3) $de^{t\Delta_{\mathcal{H},n}}f = e^{t\Delta_{\mathcal{H},n}}df$, for all $f \in \dot{D}_p^s(d, \mathbb{R}_+^n, \Lambda^k)$;
- (4) $d^*e^{t\Delta_{\mathcal{H},n}}f = e^{t\Delta_{\mathcal{H},n}}d^*f$, for all $f \in \dot{D}_p^s(d^*, \mathbb{R}_+^n, \Lambda^k)$.

Every above identities still hold replacing (\mathbf{n}, d, d^*) by $(\mathbf{t}, \delta^*, \delta)$, and \dot{d}_p^s by either d_p^s , $\dot{d}_{p,q}^s$ or even by $\dot{d}_{p,q}^s$, with $q \in [1, +\infty]$.

Proof. — Let $f \in \dot{D}_p^s(d, \mathbb{R}_+^n, \Lambda^k) \subset \dot{H}^{s,p}(\mathbb{R}_+^n, \Lambda^k)$, then by Theorem 2.29 there exists a unique $u \in \dot{D}_p^s(\Delta_{\mathcal{H},n}, \mathbb{R}_+^n, \Lambda^k) \subset \dot{D}_p^s(D_{\mathbf{n}}, \mathbb{R}_+^n, \Lambda^k) \subset \dot{D}_p^s(d, \mathbb{R}_+^n, \Lambda^k)$ such that

$$\lambda u - \Delta_{\mathcal{H},n}u = f.$$

Since $u, f \in \dot{D}_p^s(d, \mathbb{R}_+^n, \Lambda^k)$, we deduce that $\Delta_{\mathcal{H},n}u \in \dot{D}_p^s(d, \mathbb{R}_+^n, \Lambda^k)$ and we use $d^2 = 0$ to deduce

$$\lambda du - \Delta_{\mathcal{H},n}du = df.$$

Thus, we obtain that $du \in \dot{H}^{s,p}(\mathbb{R}_+^n, \Lambda^{k+1})$ is a solution of the equation

$$\lambda v - \Delta_{\mathcal{H},n}v = df.$$

Thus, uniqueness of the solution yields $du = (\lambda I - \Delta_{\mathcal{H},n})^{-1}df$. If it holds for resolvents, then it holds for semigroups, since we have the Cauchy integral formula, provided $\theta \in (\frac{\pi}{2}, \pi)$ and $t > 0$ are fixed,

$$e^{t\Delta_{\mathcal{H},n}} = \frac{1}{2\pi i} \int_{\gamma_t} e^{t\lambda} (\lambda I - \Delta_{\mathcal{H},n})^{-1} d\lambda,$$

where $\gamma_t = -\gamma_- + \gamma_0 + \gamma_+$ is the path given by

$$\gamma_{\pm} : [t^{-1}, +\infty) \rightarrow \mathbb{C}, \quad \gamma_{\pm}(s) := se^{\pm i\theta} \quad \text{and} \quad \gamma_0 : [-\theta, \theta] \rightarrow \mathbb{C}, \quad \gamma_0(s) := t^{-1}e^{is}.$$

□

PROPOSITION 2.31. — *Let $p \in (1, +\infty)$, $s \in (-1 + 1/p, 1/p)$, $k \in \llbracket 0, n \rrbracket$. For any $\lambda \geq 0$, following operators are well-defined and uniformly bounded with respect to λ*

$$\begin{aligned} d(\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}} : \dot{H}^{s,p}(\mathbb{R}_+^n, \Lambda^k) &\longrightarrow \dot{N}_p^s(d, \mathbb{R}_+^n, \Lambda^{k+1}); \\ d^*(\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}} : \dot{H}^{s,p}(\mathbb{R}_+^n, \Lambda^k) &\longrightarrow \dot{N}_p^s(d^*, \mathbb{R}_+^n, \Lambda^{k-1}). \end{aligned}$$

Moreover, the following identities also hold for all $\lambda > 0$:

- $d(\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}}f = (\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}}df$, for all $f \in \dot{D}_p^s(d, \mathbb{R}_+^n, \Lambda^k)$;
- $d^*(\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}}f = (\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}}d^*f$, for all $f \in \dot{D}_p^s(d^*, \mathbb{R}_+^n, \Lambda^k)$.

Everything still holds replacing (n, d, d^*) by (t, δ^*, δ) , and replacing $(\dot{H}^{s,p}, \dot{N}_p^s)$ by either $(\dot{B}_{p,q}^s, \dot{N}_{p,q}^s)$, $(H^{s,p}, N_p^s)$ or even by $(B_{p,q}^s, N_{p,q}^s)$ with $q \in [1, +\infty]$.

Proof. — For $\lambda \geq 0$, we introduce the representation formula,

$$(2.13) \quad (\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}}f = \frac{1}{\sqrt{\pi}} \int_0^{+\infty} e^{-\tau\lambda} e^{\tau\Delta_{\mathcal{H},n}} f \frac{d\tau}{\sqrt{\tau}}.$$

This representation formula makes sense thanks to holomorphic functional calculus, and the integral is absolutely convergent for every $f \in \dot{R}_p^s(\Delta_{\mathcal{H},n}, \mathbb{R}_+^n, \Lambda^k)$.

We use the definition of function spaces by restriction and the bounded holomorphic functional calculus, with the identity provided by point (ii) of Theorem 2.29, i.e. $E_{\mathcal{H},n}e^{\tau\Delta_{\mathcal{H},n}} = e^{\tau\Delta}E_{\mathcal{H},n}$, to obtain

$$\begin{aligned} \left\| d(\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}}f \right\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} + \left\| d^*(\lambda I - \Delta_{\mathcal{H},n})^{-\frac{1}{2}}f \right\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} \\ \lesssim_{k,n} \left\| \nabla(\lambda I - \Delta)^{-\frac{1}{2}}E_{\mathcal{H},n}f \right\|_{\dot{H}^{s,p}(\mathbb{R}^n)} \\ \lesssim_{n,k,s,p} \|f\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)}. \end{aligned}$$

Therefore, the boundedness follows by density of $\dot{R}_p^s(\Delta_{\mathcal{H},n}, \mathbb{R}_+^n, \Lambda^k)$ in $\dot{H}^{s,p}(\mathbb{R}_+^n, \Lambda^k)$. Commutations relations when $\lambda > 0$ follow from Lemma 2.30 and the representation formula (2.13). The boundedness on the Besov scale follows from real interpolation. □

According to more convenient and usual notations with respect to the field of partial differential equations we set new symbols.

Notation 2.32. — We introduce the following notations

$$\begin{aligned} \mathbf{H}_{\mathbf{n},\sigma}^{s,p} &:= \mathbf{N}_p^s(\mathbf{d}^*), & \mathbf{H}_{\gamma}^{s,p} &:= \mathbf{N}_p^s(\mathbf{d}) \quad \text{and} \quad \mathbf{H}_{\sigma}^{s,p} &:= \mathbf{N}_p^s(\delta), & \mathbf{H}_{\mathbf{t},\gamma}^{s,p} &:= \mathbf{N}_p^s(\delta^*); \\ \mathbf{B}_{p,q,\mathbf{n}}^{s,\sigma} &:= \mathbf{N}_{p,q}^s(\mathbf{d}^*), & \mathbf{B}_{p,q}^{s,\gamma} &:= \mathbf{N}_{p,q}^s(\mathbf{d}) \quad \text{and} \quad \mathbf{B}_{p,q}^{s,\sigma} &:= \mathbf{N}_{p,q}^s(\delta), & \mathbf{B}_{p,q,\mathbf{t}}^{s,\gamma} &:= \mathbf{N}_{p,q}^s(\delta^*). \end{aligned}$$

Then we are able to obtain the following result.

THEOREM 2.33. — *Let $p \in (1, +\infty)$, $q \in [1, +\infty]$, $s \in (-1 + 1/p, 1/p)$, and let $k \in \llbracket 0, n \rrbracket$. It holds that*

(i) *The following equality holds,*

$$\dot{\mathbf{N}}_p^s(\mathbf{d}, \mathbb{R}_+^n, \Lambda^k) = \overline{\dot{\mathbf{R}}_p^s(\mathbf{d}, \mathbb{R}_+^n, \Lambda^k)}^{\|\cdot\|_{\dot{\mathbf{H}}^{s,p}(\mathbb{R}_+^n)}},$$

and still holds replacing \mathbf{d} by \mathbf{d}^ .*

(ii) *The (generalized) Helmholtz–Leray projector is well-defined and bounded as a linear operator*

$$\mathbb{P} : \dot{\mathbf{H}}^{s,p}(\mathbb{R}_+^n, \Lambda^k) \longrightarrow \dot{\mathbf{H}}_{\mathbf{n},\sigma}^{s,p}(\mathbb{R}_+^n, \Lambda^k).$$

Moreover, the following identity is true

$$\mathbb{P} = \mathbf{I} - \mathbf{d}(-\Delta_{\mathcal{H},\mathbf{n}})^{-\frac{1}{2}} \mathbf{d}^*(-\Delta_{\mathcal{H},\mathbf{n}})^{-\frac{1}{2}}.$$

(iii) *The following Hodge decomposition holds*

$$\dot{\mathbf{H}}^{s,p}(\mathbb{R}_+^n, \Lambda^k) = \dot{\mathbf{H}}_{\mathbf{n},\sigma}^{s,p}(\mathbb{R}_+^n, \Lambda^k) \oplus \dot{\mathbf{H}}_{\gamma}^{s,p}(\mathbb{R}_+^n, \Lambda^k).$$

Moreover, the result remains true if we replace

- $\dot{\mathbf{H}}^{s,p}$ by $\dot{\mathbf{B}}_{p,q}^s$;
- $(\dot{\mathbf{H}}, \dot{\mathbf{B}})$ by (\mathbf{H}, \mathbf{B}) ;
- $(\mathbf{n}, \mathbf{d}, \mathbf{d}^*, \mathbb{P}, \sigma, \gamma)$ by $(\mathbf{t}, \delta^*, \delta, \mathbb{Q}, \gamma, \sigma)$.

In case of Besov spaces with $q = +\infty$, the density result of point (i) only holds in the weak sense.*

Proof. — One may reproduce the proofs of Theorem 2.21, thanks to Proposition 2.31 above. □

The following corollary is a direct consequence of the given expression for the Helmholtz–Leray projection in Theorem 2.33.

COROLLARY 2.34. — *Let $p \in (1, +\infty)$, $s \in (-1 + 1/p, 1/p)$, $\mu \in [0, \pi)$, $\lambda \in \Sigma_{\mu}$, $k \in \llbracket 0, n \rrbracket$. The following commutation identities hold for all $f \in \dot{\mathbf{H}}^{s,p}(\mathbb{R}_+^n, \Lambda^k)$, for all $t \geq 0$,*

$$\begin{aligned} (\lambda \mathbf{I} - \Delta_{\mathcal{H},\mathbf{n}})^{-1} \mathbb{P} f &= \mathbb{P} (\lambda \mathbf{I} - \Delta_{\mathcal{H},\mathbf{n}})^{-1} f, \\ e^{t\Delta_{\mathcal{H},\mathbf{n}}} \mathbb{P} f &= \mathbb{P} e^{t\Delta_{\mathcal{H},\mathbf{n}}} f. \end{aligned}$$

Above identity still holds replacing (\mathbf{n}, \mathbb{P}) by (\mathbf{t}, \mathbb{Q}) , and $\dot{\mathbf{H}}^{s,p}$ by either $\mathbf{H}^{s,p}$, $\mathbf{B}_{p,q}^s$ or even by $\dot{\mathbf{B}}_{p,q}^s$, with $q \in [1, +\infty]$.

2.3.3. Hodge–Stokes and Hodge–Maxwell operators

The present subsection is about discussing properties of Hodge–Stokes and Hodge–Maxwell operators. First, one can define **Hodge–Stokes operator**'s domain, for all $p \in (1, +\infty)$, $s \in (-1 + 1/p, 1/p)$, $k \in \llbracket 0, n \rrbracket$, by

$$(2.14) \quad \dot{D}_p^s(\mathbb{A}_{\mathcal{H},n}, \mathbb{R}_+^n, \Lambda^k) := \dot{H}_{n,\sigma}^{s,p}(\mathbb{R}_+^n, \Lambda^k) \cap \dot{D}_p^s(\Delta_{\mathcal{H},n}, \mathbb{R}_+^n, \Lambda^k),$$

and for all $u \in \dot{D}_p^s(\mathbb{A}_{\mathcal{H},n}, \mathbb{R}_+^n, \Lambda^k)$

$$(2.15) \quad \mathbb{A}_{\mathcal{H},n}u := d^*du = -\mathbb{P}\Delta u = -\Delta\mathbb{P}u = -\Delta u.$$

Above operator $\mathbb{A}_{\mathcal{H},n}$ is called the **Hodge–Stokes operator** with **absolute boundary conditions** which is a closed densely defined operator on $\dot{H}_{n,\sigma}^{s,p}(\mathbb{R}_+^n, \Lambda^k)$.

Similarly one can treat the case of **Hodge–Maxwell operators**,

$$(2.16) \quad \dot{D}_p^s(\mathbb{M}_{\mathcal{H},n}, \mathbb{R}_+^n, \Lambda^k) := \dot{H}_\gamma^{s,p}(\mathbb{R}_+^n, \Lambda^k) \cap \dot{D}_p^s(\Delta_{\mathcal{H},n}, \mathbb{R}_+^n, \Lambda^k),$$

and for all $u \in \dot{D}_p^s(\mathbb{M}_{\mathcal{H},n}, \mathbb{R}_+^n, \Lambda^k)$

$$(2.17) \quad \mathbb{M}_{\mathcal{H},n}u := dd^*u = -[I - \mathbb{P}]\Delta u = -\Delta[I - \mathbb{P}]u = -\Delta u.$$

The operator $\mathbb{M}_{\mathcal{H},n}$ defined as above is called the **Hodge–Maxwell operator** with **perfectly conductive wall boundary conditions** which is a closed densely defined operator on $\dot{H}_\gamma^{s,p}(\mathbb{R}_+^n, \Lambda^k)$.

Similarly, one may replace $(\mathbf{n}, d, \mathbb{P})$ by $(\mathbf{t}, \delta, \mathbb{Q})$, respectively in, (2.14) and (2.15), and in (2.16) and (2.17). This leads to the construction of

$$(2.18) \quad \left(\dot{D}_p^s(\mathbb{A}_{\mathcal{H},\mathbf{t}}, \mathbb{R}_+^n, \Lambda^k), \mathbb{A}_{\mathcal{H},\mathbf{t}}\right) \quad \text{and} \quad \left(\dot{D}_p^s(\mathbb{M}_{\mathcal{H},\mathbf{t}}, \mathbb{R}_+^n, \Lambda^k), \mathbb{M}_{\mathcal{H},\mathbf{t}}\right)$$

called respectively the **Hodge–Stokes operator** with **relative boundary conditions** and the **Hodge–Maxwell operator** with **relative boundary conditions** which are both closed densely defined operator on $\dot{H}_\sigma^{s,p}(\mathbb{R}_+^n, \Lambda^k)$ and $\dot{H}_{\mathbf{t},\gamma}^{s,p}(\mathbb{R}_+^n, \Lambda^k)$, respectively.

Those Hodge–Stokes and Hodge–Maxwell operator are still meaningful on other function spaces replacing $(\dot{H}^{s,p}, \dot{D}_p^s)$ by $(\dot{B}_{p,q}^s, \dot{D}_{p,q}^s)$, then $(\dot{H}, \dot{B}, \dot{D})$ by (H, B, D) .

Notice again the exception of Besov spaces, homogeneous and inhomogeneous, where the domains of any Hodge–Stokes and Hodge–Maxwell operators are only weak* dense in the case $q = +\infty$.

With the above definitions, Theorem 2.29, Corollary 2.34 and Theorem 2.33, we obtain for free the next theorem.

THEOREM 2.35. — *Let $p \in (1, +\infty)$, $s \in (-1 + 1/p, 1/p)$. For all $\mu \in (0, \pi)$, the operator $\mathbb{A}_{\mathcal{H},n}$ (resp. $\mathbb{M}_{\mathcal{H},n}$) admits a $\mathbf{H}^\infty(\Sigma_\mu)$ -functional calculus on $\dot{H}_{n,\sigma}^{s,p}(\mathbb{R}_+^n, \Lambda)$ (resp. $\dot{H}_\gamma^{s,p}(\mathbb{R}_+^n, \Lambda)$).*

Moreover, the result remains true if we replace

- $\dot{H}^{s,p}$ by $\dot{B}_{p,q}^s$, $q \in [1, +\infty]$;
- (\dot{H}, \dot{B}) by (H, B) ;
- $(\mathbf{n}, \sigma, \gamma, \mathbb{A}, \mathbb{M})$ by $(\mathbf{t}, \gamma, \sigma, \mathbb{M}, \mathbb{A})$.

3. L^q -maximal regularity with global-in-time estimates

In order to motivate the results in next Sections 3.1 and 3.2, we provide short reminders about recent advances for maximal regularity in the Sobolev framework provided by [DHMT21, Chapter 2] and [Gau24a]. We are going to follow the presentation from [Gau24a, Section 2].

First, let us consider $(D(A), A)$ a densely defined closed operator on a Banach space X . We recall, see [ABHN11, Theorem 3.7.11], that the two following assertions are equivalent:

- (1) A is ω -sectorial on X , with $\omega \in [0, \frac{\pi}{2})$;
- (2) $-A$ generates a bounded holomorphic semigroup on X , denoted by $(e^{-tA})_{t \geq 0}$.

Thus, provided that A is ω -sectorial on X for some $\omega \in [0, \frac{\pi}{2})$, for $T \in (0, +\infty]$, we look at the following abstract Cauchy problem,

$$(ACP) \quad \begin{cases} \partial_t u(t) + Au(t) = f(t), & 0 < t < T \\ u(0) = u_0, \end{cases}$$

where $f \in L^1_{loc}((0, T), X)$, $u_0 \in Y$, Y being some normed vector space depending on X and $D(A)$.

We want to look at global-in-time maximal regularity results for (ACP). To obtain estimates that are uniform in time, we require the involved function spaces to be **homogeneous**. This keypoint was captured in the work of Danchin, Hieber, Mucha and Tolksdorf [DHMT21, Chapter 2] to build a **homogeneous** version of the Da Prato–Grisvard theorem for injective sectorial operators under some additional assumptions on A . We are going to present briefly their construction to motivate the next section.

Before that, we introduce two quantities for $v \in X + D(A)$,

$$\|v\|_{\dot{\mathcal{D}}_A(\theta, q)} := \left(\int_0^{+\infty} \|t^{1-\theta} A e^{-tA} v\|_X^q \frac{dt}{t} \right)^{\frac{1}{q}}, \text{ and } \|v\|_{\mathcal{D}_A(\theta, q)} := \|v\|_X + \|v\|_{\dot{\mathcal{D}}_A(\theta, q)},$$

where $\theta \in (0, 1)$, $q \in [1, +\infty]$, with the special case

$$\|v\|_{\dot{\mathcal{D}}_A(\theta, \infty)} := \sup_{t > 0} \|t^{1-\theta} A e^{-tA} v\|_X.$$

This leads to the construction of the vector space

$$\mathcal{D}_A(\theta, q) := \{v \in X \mid \|v\|_{\dot{\mathcal{D}}_A(\theta, q)} < +\infty\}.$$

The vector space $\mathcal{D}_A(\theta, q)$ is known to be a Banach space under the norm $\|\cdot\|_{\mathcal{D}_A(\theta, q)}$ and, moreover, it satisfies the following equality with equivalence of norms

$$(3.1) \quad \mathcal{D}_A(\theta, q) = (X, D(A))_{\theta, q},$$

see [Haa06, Theorem 6.2.9]. If moreover, $0 \in \rho(A)$ it has been proved, [Haa06, Corollary 6.5.5], that $\|\cdot\|_{\dot{\mathcal{D}}_A(\theta, q)}$ and $\|\cdot\|_{\mathcal{D}_A(\theta, q)}$ are two equivalent norms on $\mathcal{D}_A(\theta, q)$. So we restrict ourselves to the case of injective operators.

ASSUMPTION 3.1. — *The operator $(D(A), A)$ is injective on X , and there exists a normed vector space $(Y, \|\cdot\|_Y)$, such that $D(A) \subset Y$, and for all $x \in D(A)$,*

$$(3.2) \quad \|Ax\|_X \sim_{X,Y,A} \|x\|_Y.$$

The idea is to construct a homogeneous version of A denoted \mathring{A} , defining first its domain

$$D(\mathring{A}) := \left\{ y \in Y \mid \exists (x_n)_{n \in \mathbb{N}} \subset D(A), \|y - x_n\|_Y \xrightarrow{n \rightarrow +\infty} 0 \right\}.$$

So that, for all $y \in d(\mathring{A})$, X being complete, it is meaning full to set

$$\mathring{A}y := \lim_{n \rightarrow +\infty} Ax_n.$$

Constructed this way, the operator \mathring{A} is then injective on $D(\mathring{A})$. We notice that $D(\mathring{A})$, endowed with the norm $\|\mathring{A} \cdot\|_X$, is a normed vector space, but not necessarily complete. We also need the existence of a Hausdorff topological vector space Z , such that $X, Y \subset Z$, and to consider the following assumption.

ASSUMPTION 3.2. — *The operator $(D(A), A)$ and the normed vector space Y are such that*

$$(3.3) \quad X \cap D(\mathring{A}) = D(A).$$

As a consequence of all above assumptions, we can extend naturally, see [DHMT21, Remark 2.7], $(e^{-tA})_{t \geq 0}$ to a C_0 -semigroup,

$$e^{-tA} : X + D(\mathring{A}) \longrightarrow X + D(\mathring{A}), t \geq 0,$$

so that, one can fully make sense of the following vector space,

$$\mathring{\mathcal{D}}_A(\theta, q) := \left\{ v \in X + D(\mathring{A}) \mid \|v\|_{\mathring{\mathcal{D}}_A(\theta, q)} < +\infty \right\}.$$

Similarly to what happens for $\mathcal{D}_A(\theta, q)$ in (3.1), it has been proved in [DHMT21, Proposition 2.12], that the following equality holds with equivalence of norms,

$$(3.4) \quad \mathring{\mathcal{D}}_A(\theta, q) = \left(X, D(\mathring{A}) \right)_{\theta, q},$$

but the possible lack of completeness of $D(\mathring{A})$ implies that $\mathring{\mathcal{D}}_A(\theta, q)$ is not necessarily complete. This has consequences on how to consider the forcing term f in (ACP), choosing $f \in L^q((0, T), \mathcal{D}_A(\theta, q))$ instead of $f \in L^q((0, T), \mathring{\mathcal{D}}_A(\theta, q))$ to avoid definition issues, the latter choice being possible when $\mathring{\mathcal{D}}_A(\theta, q)$ is a Banach space.

THEOREM 3.3 ([DHMT21, Theorem 2.20]). — *Let $\omega \in [0, \frac{\pi}{2})$, $(d(A), A)$ an ω -sectorial operator on a Banach space X such that Assumptions 3.1 and 3.2 are satisfied. Let $q \in [1, +\infty)$, $\theta \in (0, \frac{1}{q})$, $\theta_q := \theta + 1 - 1/q$, and let $T \in (0, +\infty]$.*

For $f \in L^q((0, T), \mathcal{D}_A(\theta, q))$ and $u_0 \in \mathring{\mathcal{D}}_A(\theta_q, q)$, the problem (ACP) admits an unique mild solution

$$u \in C_b^0([0, T], \mathring{\mathcal{D}}_A(\theta_q, q)),$$

such that $\partial_t u, Au \in L^q((0, T), \mathring{D}_A(\theta, q))$ with estimates,

$$(3.5) \quad \|u\|_{L^\infty([0, T], \mathring{D}_A(\theta, q))} + \|(\partial_t u, Au)\|_{L^q((0, T), \mathring{D}_A(\theta, q))} \\ \lesssim_A \|f\|_{L^q((0, T), \mathring{D}_A(\theta, q))} + \|u_0\|_{\mathring{D}_A(\theta, q)}.$$

In case $q = +\infty$, we assume in addition that $u_0 \in D(A^2)$ and then for each $\theta \in (0, 1)$,

$$(3.6) \quad \|(\partial_t u, Au)\|_{L^\infty([0, T], \mathring{D}_A(\theta, \infty))} \lesssim_A \|f\|_{L^\infty((0, T), \mathcal{D}_A(\theta, \infty))} + \|Au_0\|_{\mathring{D}_A(\theta, \infty)}.$$

One also have the following result.

THEOREM 3.4 ([Gau24a, Theorem 4.7]). — *Let $\omega \in [0, \frac{\pi}{2})$, $(d(A), A)$ an ω -sectorial operator on a UMD Banach space X , such that it has BIP on X of type $\theta_A < \frac{\pi}{2}$, and satisfies assumptions (3.1) and (3.2). Let $q \in (1, +\infty)$, $\alpha \in (-1 + 1/q, 1/q)$ and $T \in (0, +\infty]$. We set $\alpha_q := 1 + \alpha - 1/q$.*

For $f \in \dot{H}^{\alpha, q}((0, T), X)$, $u_0 \in \mathring{D}_A(\alpha_q, q)$, the problem (ACP) admits a unique mild solution $u \in C_b^0([0, T], \mathring{D}_A(\alpha_q, q))$ such that $\partial_t u, Au \in \dot{H}^{\alpha, q}((0, T), X)$ with estimate

$$(3.7) \quad \|u\|_{L^\infty([0, T], \mathring{D}_A(\alpha_q, q))} \lesssim_{A, q, \alpha} \|(\partial_t u, Au)\|_{\dot{H}^{\alpha, q}((0, T), X)} \\ \lesssim_{A, q, \alpha} \|f\|_{\dot{H}^{\alpha, q}((0, T), X)} + \|u_0\|_{\mathring{D}_A(\alpha_q, q)}.$$

Moreover, if $u_0 \in \mathcal{D}_A(\alpha_q, q)$ for all $\beta \in [0, 1]$,

$$(3.8) \quad \|(-\partial_t)^\beta A^{1-\beta} u\|_{\dot{H}^{\alpha, q}((0, T), X)} \lesssim_{A, q, \alpha} \|f\|_{\dot{H}^{\alpha, q}((0, T), X)} + \|u_0\|_{\mathring{D}_A(\alpha_q, q)}.$$

Remark 3.5. —

- In Theorem 3.4, Assumptions 3.1 and 3.2 are assumed here in order to ensure that $\mathring{D}_A(\theta, q)$ is a well-defined, even if not complete, normed vector space. The estimate (3.8) still holds for $u_0 \in \mathring{D}_A(1 + \alpha - 1/q, q)$ whenever this space is complete.
- If $u_0 = 0$, the estimate (3.8) remains valid if we replace the operator $(-\partial_t)^{1-\beta}$ by $(\partial_t)^{1-\beta}$.
- If one asks instead the initial data u_0 to be in the smaller, but complete, space $\mathcal{D}_A(\theta, q)$ then one can drop Assumptions 3.1 and 3.2, and the estimate (3.7) still holds.

Before going further, we want to simplify notations. From now on, we will only consider function spaces on \mathbb{R}_+^n and no longer on \mathbb{R}^n , so that we drop the mention of the open set in the domains of operators, we also drop the mention of degree of differential forms, except when it is necessary. Any discussion involving Dirichlet and Neumann Laplacians will always contain the implicit information that their domains are made of scalar valued functions, whereas talking about Hodge Laplacians and their derived operators will always contain the implicit information we are talking about general differential forms valued functions of any degree, unless it is explicitly stated.

A first aim of this section is about to give an explicit description of homogeneous interpolation spaces, provided $\theta \in (0, 1)$, $q \in [1, +\infty]$,

$$(3.9) \quad (X, D(\mathring{A}))_{\theta, q} = \mathring{D}_A(\theta, q),$$

where $X = \dot{H}^{s,p}, \dot{B}_{p,r}^s$ and $A \in \{-\Delta_{\mathcal{H}}, \mathbb{A}_{\mathcal{H}}, \mathbb{M}_{\mathcal{H}}\}$, with $p \in (1, +\infty)$, $-1 + 1/p < s < 1/p$, $r \in [1, +\infty)$. The main task here will be to compute the space (3.9) above, provided $A = -\Delta_{\mathcal{H}}$. Indeed, for the related Hodge–Stokes operator, due to the commutation relations between the Hodge Laplacian and its Helmholtz–Leray projections, see (2.15), (2.17) and Corollary 2.34, we should have (at least formally or up to a dense subset)

$$\begin{aligned} \dot{\mathcal{D}}_{\mathbb{A}_{\mathcal{H},n}}^{s,p}(\theta, q) &= \left(\dot{H}_{n,\sigma}^{s,p}(\mathbb{R}_+^n), \dot{D}_p^s(\mathring{\mathbb{A}}_{\mathcal{H},n}) \right)_{\theta,q} \\ &= \mathbb{P} \left(\dot{H}^{s,p}(\mathbb{R}_+^n), \dot{D}_p^s(\mathring{\Delta}_{\mathcal{H},n}) \right)_{\theta,q} \\ &= \mathbb{P} \dot{\mathcal{D}}_{-\Delta_{\mathcal{H},n}}^{s,p}(\theta, q). \end{aligned}$$

Obviously similar identities can be obtained with $(\mathfrak{t}, \mathbb{Q})$ instead of $(\mathfrak{n}, \mathbb{P})$, but also for the Hodge–Maxwell operators up to appropriate changes.

Secondly, we will aim to recover global-in-time L^q -maximal regularity estimates for the abstract Cauchy problem (ACP), provided $T \in (0, +\infty]$, $A \in \{-\Delta_{\mathcal{H}}, \mathbb{A}_{\mathcal{H}}, \mathbb{M}_{\mathcal{H}}\}$, so that we will apply Theorems 3.3 and 3.4.

3.1. Interpolation of homogeneous $\dot{H}^{s,p}$ -domains of operators

We start this section claiming that one can reduce the problem to the computation of interpolation spaces

$$\dot{\mathcal{D}}_{-\Delta_{\mathcal{D}}}^{s,p}(\theta, q) \quad \text{and} \quad \dot{\mathcal{D}}_{-\Delta_{\mathcal{N}}}^{s,p}(\theta, q).$$

We recall here for convenience that $-\Delta_{\mathcal{D}}$ and $-\Delta_{\mathcal{N}}$ stands respectively for the (negative) Dirichlet and the Neumann Laplacian on the half-space for which a wide review of their properties in homogeneous function spaces was achieved by the author, see [Gau24b, Section 5].

LEMMA 3.6. — *Let $p \in (1, +\infty)$, $s \in (-1 + 1/p, 1/p)$, $k \in \llbracket 0, n \rrbracket$. For all $u \in \dot{D}_p^s(\Delta_{\mathcal{H},n}, \Lambda^k)$, we have*

$$\Delta_{\mathcal{H},n}u = \sum_{I \in \mathcal{I}_{n-1}^k} \Delta_{\mathcal{N}}u_I dx_I + \sum_{I' \in \mathcal{I}_{n-1}^{k-1}} \Delta_{\mathcal{D}}u_{I',n} dx_{I'} \wedge dx_n.$$

We also have estimates,

$$\begin{aligned} &\|\delta du\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} + \|d\delta u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} \\ &\sim_{p,s,n} \sum_{I \in \mathcal{I}_{n-1}^k} \|\Delta_{\mathcal{N}}u_I\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} + \sum_{I' \in \mathcal{I}_{n-1}^{k-1}} \|\Delta_{\mathcal{D}}u_{I',n}\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} \\ &\sim_{p,s,n} \|\nabla^2 u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} \\ &\sim_{p,s,n} \|u\|_{\dot{H}^{s+2,p}(\mathbb{R}_+^n)}. \end{aligned}$$

The result still holds replacing $(\mathfrak{n}, \mathcal{N}, \mathcal{D})$ by $(\mathfrak{t}, \mathcal{D}, \mathcal{N})$.

Proof. — The results combine [Gau24b, Propositions 5.4 & 5.6], formula (2.10), Proposition 2.28 and Theorem 2.29. Indeed, we have the consequence, that for $u \in [\dot{H}^{s,p} \cap \dot{H}^{s+2,p}](\mathbb{R}_+^n, \Lambda^k)$,

$$u \in \dot{D}_p^s(\Delta_{\mathcal{H},n}, \Lambda^k) \Leftrightarrow \begin{cases} u_I \in \dot{D}_p^s(\Delta_{\mathcal{D}}), & \text{if } n \in I, \\ u_I \in \dot{D}_p^s(\Delta_{\mathcal{N}}), & \text{if } n \notin I. \end{cases}$$

By the identity provided by Proposition 2.28,

$$E_{\mathcal{H},n} \Delta_{\mathcal{H},n} u = \Delta E_{\mathcal{H},n} u,$$

and the boundedness properties of $E_{\mathcal{H},n}$, (2.11), we deduce by the definition of function spaces by restriction

$$\begin{aligned} \|u\|_{\dot{H}^{s+2,p}(\mathbb{R}_+^n)} &\leq \|E_{\mathcal{H},n} u\|_{\dot{H}^{s+2,p}(\mathbb{R}^n)} = \|\Delta E_{\mathcal{H},n} u\|_{\dot{H}^{s,p}(\mathbb{R}^n)} \\ &= \|E_{\mathcal{H},n} \Delta_{\mathcal{H},n} u\|_{\dot{H}^{s,p}(\mathbb{R}^n)} \lesssim_{p,s,n} \|\Delta_{\mathcal{H},n} u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)}. \end{aligned}$$

Now, we close the estimates in two different ways

$$\begin{aligned} \|\Delta_{\mathcal{H},n} u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} &\leq \left\{ \sum_{I \in \mathcal{I}_{n-1}^k} \|\Delta_{\mathcal{N}} u_I\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} + \sum_{I' \in \mathcal{I}_{n-1}^{k-1}} \|\Delta_{\mathcal{D}} u_{I',n}\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)}, \right. \\ &\quad \left. \|\delta du\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} + \|d\delta u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)}, \right. \\ &\lesssim_{p,s,n} \|\nabla^2 u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)}. \end{aligned}$$

This ends the proof of the Lemma 3.6. □

And for the same reasons, one has more generally,

LEMMA 3.7. — *Let $p \in (1, +\infty)$, $s \in (-1 + 1/p, 1/p)$, $\alpha \in [0, 2]$, such that $s + \alpha \neq 1/p, 1 + 1/p$. For all $u \in \dot{D}_p^s((-\Delta_{\mathcal{H}})^{\frac{\alpha}{2}}, \Lambda)$, we have*

$$\|(-\Delta_{\mathcal{H}})^{\frac{\alpha}{2}} u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} \sim_{p,s,\alpha,n} \|u\|_{\dot{H}^{s+\alpha,p}(\mathbb{R}_+^n)} \sim_{p,s,\alpha,n} \|(-\Delta_{\mathcal{H}})^{\frac{s+\alpha}{2}} u\|_{L^p(\mathbb{R}_+^n)}.$$

We recall that, in particular, $\Delta_{\mathcal{H},n}|_{\Lambda^0} = \Delta_{\mathcal{N}}$ and $\Delta_{\mathcal{H},n}|_{\Lambda^0} = \Delta_{\mathcal{D}}$.

In general, explicit description for interpolation spaces with boundary condition may be quite tedious. We mention the work of Guidetti [Gui91a, Gui91b], where such investigation is done. Guidetti’s results were used to make an extensive treatment of elliptic boundary value problem with general Lopatinskii–Shapiro boundary conditions in inhomogeneous Besov spaces on the half-space and on bounded domains with smooth boundary.

Thanks to Lemmas 3.6 and 3.7, the current work will be reduced to Dirichlet and Neumann boundary conditions in the homogeneous case which is unknown to the author’s knowledge yet.

For $p \in (1, +\infty)$, $q \in [1, +\infty]$, $s \in (-1 + 1/p, 2 + 1/p)$, such that $(\mathcal{C}_{s,p,q})$ is satisfied, we set:

$$\begin{aligned} \dot{B}_{p,q,\mathcal{D}}^s(\mathbb{R}_+^n) &:= \begin{cases} \dot{B}_{p,q}^s(\mathbb{R}_+^n), & \text{if } (s < \frac{1}{p}), \\ \left\{ u \in \dot{B}_{p,q}^s(\mathbb{R}_+^n) \mid u|_{\partial\mathbb{R}_+^n} = 0 \right\}, & \text{if } (s > \frac{1}{p}) \text{ or } (s = \frac{1}{p}, q = 1), \end{cases} \\ \dot{B}_{p,q,\mathcal{N}}^s(\mathbb{R}_+^n) &:= \begin{cases} \dot{B}_{p,q}^s(\mathbb{R}_+^n), & \text{if } (s < 1 + \frac{1}{p}), \\ \left\{ u \in \dot{B}_{p,q}^s(\mathbb{R}_+^n) \mid \partial_\nu u|_{\partial\mathbb{R}_+^n} = 0 \right\}, & \text{if } (s > 1 + \frac{1}{p}) \\ & \text{or } (s = 1 + \frac{1}{p}, q = 1). \end{cases} \end{aligned}$$

and similarly if $(\mathcal{C}_{s,p})$ is satisfied, we also set:

$$\begin{aligned} \dot{H}_{\mathcal{D}}^{s,p}(\mathbb{R}_+^n) &:= \begin{cases} \dot{H}^{s,p}(\mathbb{R}_+^n), & \text{if } s < \frac{1}{p}, \\ \left\{ u \in \dot{H}^{s,p}(\mathbb{R}_+^n) \mid u|_{\partial\mathbb{R}_+^n} = 0 \right\}, & \text{if } s > \frac{1}{p}, \end{cases} \\ \dot{H}_{\mathcal{N}}^{s,p}(\mathbb{R}_+^n) &:= \begin{cases} \dot{H}^{s,p}(\mathbb{R}_+^n), & \text{if } s < 1 + \frac{1}{p}, \\ \left\{ u \in \dot{H}^{s,p}(\mathbb{R}_+^n) \mid \partial_\nu u|_{\partial\mathbb{R}_+^n} = 0 \right\}, & \text{if } s > 1 + \frac{1}{p}. \end{cases} \end{aligned}$$

And then, for $\mathcal{J} \in \{\mathcal{D}, \mathcal{N}\}$, we introduce the following subspace

$$Y_{\mathcal{J}} := \bigcap_{\substack{s \in (-1+1/p, 1/p) \\ p \in (1, +\infty) \\ q \in [1, +\infty]}} [\mathbf{D}_p^s \cap \dot{\mathbf{D}}_p^s \cap \mathbf{D}_{p,q}^s \cap \dot{\mathbf{D}}_{p,q}^s](\Delta_{\mathcal{J}}).$$

PROPOSITION 3.8. — *Let $p \in (1, +\infty)$, $q \in [1, +\infty)$, $s \in (-1 + 1/p, 2 + 1/p)$, such that $(\mathcal{C}_{s,p,q})$ is satisfied, we have that*

$$\dot{B}_{p,q,\mathcal{J}}^s(\mathbb{R}_+^n) = \overline{Y_{\mathcal{J}}}^{\|\cdot\|_{\dot{B}_{p,q}^s(\mathbb{R}_+^n)}},$$

whenever

- $\mathcal{J} = \mathcal{D}$, $s \neq 1/p, 1 + 1/p$;
- $\mathcal{J} = \mathcal{N}$, $s \neq 1 + 1/p$.

When $q = +\infty$, we still have weak* density.

Proof. — We recall that for all $\tilde{p} \in (1, +\infty)$, $\tilde{q} \in [1, +\infty)$, $\tilde{s} \in \mathbb{R}$, we have that

$$\dot{B}_{\tilde{p},\tilde{q}}^{\tilde{s}}(\mathbb{R}_+^n) = \overline{\mathcal{S}_0(\mathbb{R}_+^n)}^{\|\cdot\|_{\dot{B}_{\tilde{p},\tilde{q}}^{\tilde{s}}(\mathbb{R}_+^n)}}.$$

- First, assume that $s \in (1 + 1/p, 2 + 1/p)$, for $u \in \dot{B}_{p,q,\mathcal{J}}^s(\mathbb{R}_+^n)$, we set $f := -\Delta_{\mathcal{J}} u \in \dot{B}_{p,q}^{s-2}(\mathbb{R}_+^n)$. For $(f_j)_{j \in \mathbb{N}} \subset \mathcal{S}_0(\mathbb{R}_+^n)$ such that

$$f_j \xrightarrow{j \rightarrow +\infty} f, \quad \text{in } \dot{B}_{p,q}^{s-2}(\mathbb{R}_+^n).$$

Since it is not clear that $u_j := (-\Delta_{\mathcal{J}})^{-1} f_j$ is an element of $Y_{\mathcal{J}}$, we set for all $\lambda > 0$,

$$u_{\lambda,j} := (\lambda I - \Delta_{\mathcal{J}})^{-1} f_j \in Y_{\mathcal{J}}$$

where belonging to the space $Y_{\mathcal{J}}$ is a consequence of [Gau24b, Propositions 5.4 & 5.6]. For $\mu, \lambda > 0$, by [Gau24b, Propositions 5.7 & 5.8] and Proposition 1.4, we have

$$\begin{aligned} \|u_{j,\lambda} - u_{j,\mu}\|_{\dot{B}_{p,q}^s(\mathbb{R}_+^n)} &\lesssim_{s,p,q,n} \left\| -\Delta_{\mathcal{J}}(\lambda\mathbf{I} - \Delta_{\mathcal{J}})^{-1} f_j + \Delta_{\mathcal{J}}(\mu\mathbf{I} - \Delta_{\mathcal{J}})^{-1} f_j \right\|_{\dot{B}_{p,q}^{s-2}(\mathbb{R}_+^n)} \\ &\lesssim_{s,p,q,n} \left\| -\Delta_{\mathcal{J}}(\lambda\mathbf{I} - \Delta_{\mathcal{J}})^{-1} f_j - f_j \right\|_{\dot{B}_{p,q}^{s-2}(\mathbb{R}_+^n)} \\ &\quad + \left\| f_j + \Delta_{\mathcal{J}}(\mu\mathbf{I} - \Delta_{\mathcal{J}})^{-1} f_j \right\|_{\dot{B}_{p,q}^{s-2}(\mathbb{R}_+^n)} \xrightarrow{\lambda,\mu \rightarrow 0} 0. \end{aligned}$$

By uniqueness of the solution for the Neumann (resp. Dirichlet) problem provided by [Gau24b, Proposition 5.8] (resp. [Gau24b, Proposition 5.7]), $(u_{\mu,j})_{\mu>0}$ is a Cauchy net that admits a limit that must be the unique solution u_j . Since as j tends to infinity, $u_j = (-\Delta_{\mathcal{J}})^{-1} f_j$ converges to $u = (-\Delta_{\mathcal{J}})^{-1} f$, it follows that for any $\varepsilon > 0$, one can find j and λ large enough so that

$$\|u - u_{j,\lambda}\|_{\dot{B}_{p,q}^s(\mathbb{R}_+^n)} \leq \|u - u_j\|_{\dot{B}_{p,q}^s(\mathbb{R}_+^n)} + \|u_j - u_{j,\lambda}\|_{\dot{B}_{p,q}^s(\mathbb{R}_+^n)} < 2\varepsilon.$$

This concludes the case $s \in (1 + 1/p, 2 + 1/p)$.

- For $s \in (1/p, 1 + 1/p)$, we consider first the case $\mathcal{J} = \mathcal{N}$. Again, for $u \in \dot{B}_{p,q,\mathcal{N}}^s(\mathbb{R}_+^n) = \dot{B}_{p,q}^s(\mathbb{R}_+^n)$, we can introduce $(u_j)_{j \in \mathbb{N}} \subset \mathcal{S}_0(\overline{\mathbb{R}_+^n})$ such that

$$u_j \xrightarrow{j \rightarrow +\infty} u,$$

and we set for all $\tau > 0$,

$$u_{\tau,j} := (\mathbf{I} - \tau\Delta_{\mathcal{J}})^{-1} u_j \in Y_{\mathcal{J}}$$

where belonging to the space $Y_{\mathcal{J}}$ is a consequence of [Gau24b, Propositions 5.4 & 5.6]. It is direct to see that

$$u_{\tau,j} \xrightarrow{\tau \rightarrow 0} u_j \xrightarrow{j \rightarrow +\infty} u \quad \text{in } \dot{B}_{p,q}^s(\mathbb{R}_+^n).$$

This argument still works for $s \in (-1 + 1/p, 1 + 1/p)$, when $\mathcal{J} = \mathcal{N}$.

For the case $\mathcal{J} = \mathcal{D}$, since $\dot{B}_{p,q,\mathcal{D}}^s(\mathbb{R}_+^n) = \dot{B}_{p,q,0}^s(\mathbb{R}_+^n)$ and $C_c^\infty(\mathbb{R}_+^n) \subset Y_{\mathcal{D}}$, the result follows from [Gau24b, Lemma 3.16].

- Finally, when $s \in (-1 + 1/p, 1/p)$, we notice that $\dot{B}_{p,q,\mathcal{J}}^s(\mathbb{R}_+^n) = \dot{B}_{p,q,0}^s(\mathbb{R}_+^n) = \dot{B}_{p,q}^s(\mathbb{R}_+^n)$. Since $C_c^\infty(\mathbb{R}_+^n) \subset Y_{\mathcal{J}}$, the result follows from [Gau24b, Corollary 3.18]. \square

The next result has a similar proof and is left to the reader.

PROPOSITION 3.9. — *Let $p \in (1, +\infty)$, $s_0, s_1 \in (1/p, 2 + 1/p)$, $\mathcal{J} \in \{\mathcal{D}, \mathcal{N}\}$ such that $(\mathcal{C}_{s_0,p})$ is satisfied, we have*

$$\left[\dot{H}_{\mathcal{J}}^{s_0,p} \cap \dot{H}^{s_1,p} \right] (\mathbb{R}_+^n) = \overline{Y_{\mathcal{J}}}^{\|\cdot\|_{[\dot{H}^{s_0,p} \cap \dot{H}^{s_1,p}](\mathbb{R}_+^n)}},$$

whenever $1/p < s_0, s_1 < 2 + 1/p$, except for $s = 1 + 1/p$ when $\mathcal{J} = \mathcal{N}$.

The next lemma is inspired from [Gui91b, Lemma 2.4].

LEMMA 3.10. — Let $p_j \in (1, +\infty)$, $q_j \in [1, +\infty)$, and $s_j > 1/p_j$, for $j \in \{0, 1\}$. Let T be the map

$$T : f \longmapsto \left[(x', x_n) \mapsto e^{-x_n(-\Delta')^{\frac{1}{2}}} f(x') \right].$$

(i) Assume $s_j \in (1/p_j, 1 + 2/p_j)$, for $j \in \{0, 1\}$. Then the operator defined formally by

$$\mathcal{P}_{\mathcal{D}} u := u - T \left[u|_{\partial\mathbb{R}_+^n} \right],$$

is such that

(a) If (\mathcal{C}_{s_0, p_0}) is satisfied, then $\mathcal{P}_{\mathcal{D}} : [\dot{H}^{s_0, p_0} \cap \dot{H}^{s_1, p_1}](\mathbb{R}_+^n) \longrightarrow [\dot{H}_{\mathcal{D}}^{s_0, p_0} \cap \dot{H}^{s_1, p_1}](\mathbb{R}_+^n)$ is a well-defined linear and bounded projection. For all $u \in [\dot{H}^{s_0, p_0} \cap \dot{H}^{s_1, p_1}](\mathbb{R}_+^n)$ the following estimate is true

$$\|\mathcal{P}_{\mathcal{D}} u\|_{\dot{H}^{s_j, p_j}(\mathbb{R}_+^n)} \lesssim_{p_j, s_j, n} \|u\|_{\dot{H}^{s_j, p_j}(\mathbb{R}_+^n)}, j \in \{0, 1\}.$$

(b) If instead $(\mathcal{C}_{s_0, p_0, q_0})$ is satisfied, then the above statement still holds with $(\dot{B}_{p_0, q_0}^{s_0}, \dot{B}_{p_1, q_1}^{s_1})$ replacing $(\dot{H}^{s_0, p_0}, \dot{H}^{s_1, p_1})$.

We also have that $\mathcal{P}_{\mathcal{D}} : \dot{B}_{p_0, \infty}^{s_0}(\mathbb{R}_+^n) \longrightarrow \dot{B}_{p_0, \infty, \mathcal{D}}^{s_0}(\mathbb{R}_+^n)$ is also well-defined linear and bounded.

(ii) Assume $s_j \in (1 + 1/p_j, 1 + 2/p_j)$, for $j \in \{0, 1\}$. Then the operator defined formally by

$$\mathcal{P}_{\mathcal{N}} u := u + (-\Delta')^{-\frac{1}{2}} T \left[\partial_{x_n} u|_{\partial\mathbb{R}_+^n} \right],$$

satisfies points (ia) and (ib) with \mathcal{N} instead of \mathcal{D} .

Proof. — This is a direct consequence of [Gau24b, Proposition B.2, Corollary B.3]. □

PROPOSITION 3.11. — Let $p \in (1, +\infty)$, $s \in (-1 + 1/p, 1/p)$, $\mathcal{J} \in \{\mathcal{D}, \mathcal{N}, \mathcal{H}\}$, then $(\dot{D}_p^s(\Delta_{\mathcal{J}}), -\Delta_{\mathcal{J}})$ satisfies Assumptions 3.1 and 3.2. In other words, $-\Delta_{\mathcal{J}}$ is injective on $\dot{H}^{s, p}(\mathbb{R}_+^n)$, and we can define

$$\dot{D}_p^s(\mathring{\Delta}_{\mathcal{J}}) := \left\{ u \in \dot{H}^{s+2, p}(\mathbb{R}_+^n) \mid \exists (u_j)_{j \in \mathbb{N}} \subset \dot{D}_p^s(\Delta_{\mathcal{J}}), \|u - u_j\|_{\dot{H}^{s+2, p}(\mathbb{R}_+^n)} \xrightarrow{j \rightarrow +\infty} 0 \right\}$$

such that it also satisfies

$$(3.10) \quad \dot{H}^{s, p}(\mathbb{R}_+^n) \cap \dot{D}_p^s(\mathring{\Delta}_{\mathcal{J}}) = \dot{D}_p^s(\Delta_{\mathcal{J}}).$$

The result holds with either $\mathbb{A}_{\mathcal{H}, \mathbf{n}}$ (resp. $\mathbb{M}_{\mathcal{H}, \mathbf{n}}$) on $\dot{H}_{\mathbf{n}, \sigma}^{s, p}(\mathbb{R}_+^n)$ (resp. $\dot{H}_{\gamma}^{s, p}(\mathbb{R}_+^n)$), and similarly replacing $(\mathbf{n}, \sigma, \gamma, \mathbb{A}, \mathbb{M})$ by $(\mathbf{t}, \gamma, \sigma, \mathbb{M}, \mathbb{A})$.

Proof. — We only show (3.10). The following inclusion is clear

$$\dot{D}_p^s(\Delta_{\mathcal{J}}) \subset \dot{H}^{s, p}(\mathbb{R}_+^n) \cap \dot{D}_p^s(\mathring{\Delta}_{\mathcal{J}}).$$

Now, let $u \in \dot{H}^{s, p}(\mathbb{R}_+^n) \cap \dot{D}_p^s(\mathring{\Delta}_{\mathcal{J}})$, by definition and Lemma 3.6, we obtain

$$u \in \dot{H}^{s, p}(\mathbb{R}_+^n) \cap \dot{H}^{s+2, p}(\mathbb{R}_+^n).$$

It suffices to show that u has appropriate boundary conditions. We assume here that $\mathcal{J} = \mathcal{D}$, other cases would be achieved similarly. Let $(u_j)_{j \in \mathbb{N}} \subset \dot{D}_p^s(\Delta_{\mathcal{D}})$, such that

$$\|u - u_j\|_{\dot{H}^{s+2,p}(\mathbb{R}_+^n)} \xrightarrow{j \rightarrow +\infty} 0.$$

Since $u - u_j \in \dot{H}^{s,p}(\mathbb{R}_+^n) \cap \dot{H}^{s+2,p}(\mathbb{R}_+^n)$, one may apply [Gau24b, Proposition 4.6], and use $u_j|_{\partial\mathbb{R}_+^n} = 0$ to obtain

$$\|u|_{\partial\mathbb{R}_+^n}\|_{\dot{B}_{p,p}^{s+2-1/p}(\mathbb{R}^{n-1})} \lesssim_{s,p,n} \|u - u_j\|_{\dot{H}^{s+2,p}(\mathbb{R}_+^n)} \xrightarrow{j \rightarrow +\infty} 0.$$

Therefore $u|_{\partial\mathbb{R}_+^n} = 0$ so that $u \in \dot{D}_p^s(\Delta_{\mathcal{D}})$. □

The Proposition 3.11 tells us that, for all $p \in (1, +\infty)$, $s \in (-1 + 1/p, 1/p)$, it makes sense to consider the semigroup,

$$e^{t\Delta_{\mathcal{H}}} : \dot{H}^{s,p}(\mathbb{R}_+^n) + \dot{D}_p^s(\dot{\Delta}_{\mathcal{H}}) \longrightarrow \dot{H}^{s,p}(\mathbb{R}_+^n) + \dot{D}_p^s(\dot{\Delta}_{\mathcal{H}}),$$

thanks to [DHMT21, Chapter 2, Section 1].

For convenience of notations, and for later use, one may think about Lemma 3.6, we also set for all $p \in (1, +\infty)$, $q \in [1, +\infty]$, $s \in (-1 + 1/p, 2 + 1/p)$, such that $(\mathcal{C}_{s,p,q})$ is satisfied, and for $k \in \llbracket 0, n \rrbracket$,

$$(3.11) \quad \dot{B}_{p,q,\mathcal{H}_n}^s(\mathbb{R}_+^n, \Lambda^k) := \begin{cases} \dot{B}_{p,q}^s(\mathbb{R}_+^n, \Lambda^k), & s < \frac{1}{p}, \\ \left\{ u \in \dot{B}_{p,q}^s(\mathbb{R}_+^n, \Lambda^k) \mid \mathbf{e}_n \lrcorner u|_{\partial\mathbb{R}_+^n} = 0 \right\}, & 0 < s - \frac{1}{p} < 1, \\ \left\{ u \in \dot{B}_{p,q}^s(\mathbb{R}_+^n, \Lambda^k) \mid \mathbf{e}_n \lrcorner (u, du)|_{\partial\mathbb{R}_+^n} = (0, 0) \right\}, & 1 < s - \frac{1}{p} < 2. \end{cases}$$

It is not difficult to see from point (iii) of Theorem A.2, that,

$$\dot{B}_{p,q,\mathcal{H}_n}^s(\mathbb{R}_+^n, \Lambda^k) \simeq \dot{B}_{p,q,\mathcal{D}}^s(\mathbb{R}_+^n)^{\binom{n-1}{k-1}} \times \dot{B}_{p,q,\mathcal{N}}^s(\mathbb{R}_+^n)^{\binom{n-1}{k}},$$

for which one may check for instance the **Step 3** of Theorem A.2's proof.

One can also build in the same fashion $\dot{B}_{p,q,\mathcal{H}_t}^s(\mathbb{R}_+^n, \Lambda^k)$, with boundary conditions $\nu \wedge u|_{\partial\mathbb{R}_+^n} = 0$ and $\nu \wedge \delta u|_{\partial\mathbb{R}_+^n} = 0$, so that

$$\dot{B}_{p,q,\mathcal{H}_t}^s(\mathbb{R}_+^n, \Lambda^k) \simeq \dot{B}_{p,q,\mathcal{N}}^s(\mathbb{R}_+^n)^{\binom{n-1}{k-1}} \times \dot{B}_{p,q,\mathcal{D}}^s(\mathbb{R}_+^n)^{\binom{n-1}{k}}.$$

We denote by $\dot{B}_{p,q,\mathcal{H}}^s(\mathbb{R}_+^n)$, either $\dot{B}_{p,q,\mathcal{H}_n}^s(\mathbb{R}_+^n, \Lambda)$ or $\dot{B}_{p,q,\mathcal{H}_t}^s(\mathbb{R}_+^n, \Lambda)$.

PROPOSITION 3.12. — *Let $p \in (1, +\infty)$, $q \in [1, +\infty]$, and $s \in (-1 + 1/p, 1/p)$. For all $\theta \in (0, 1)$ such that $(\mathcal{C}_{s+2\theta,p,q})$ is satisfied, provided $\mathcal{J} \in \{\mathcal{D}, \mathcal{N}, \mathcal{H}\}$, one has*

$$\left(\dot{H}^{s,p}(\mathbb{R}_+^n), \dot{D}_p^s(\dot{\Delta}_{\mathcal{J}}) \right)_{\theta,q} = \dot{B}_{p,q,\mathcal{J}}^{s+2\theta}(\mathbb{R}_+^n),$$

with equivalence of norms, whenever

- $s + 2\theta \neq 1/p$ if $\mathcal{J} = \mathcal{D}$,
- $s + 2\theta \neq 1 + 1/p$ if $\mathcal{J} = \mathcal{N}$,
- $s + 2\theta \neq 1/p, 1 + 1/p$ if $\mathcal{J} = \mathcal{H}$.

The proof is heavily inspired from the one of [DHMT21, Proposition 4.12].

Proof. —

Step 1: We start applying [Gau24b, Proposition 3.17] which yields the embedding

$$\left(\dot{H}^{s,p}(\mathbb{R}_+^n), \dot{D}_p^s(\dot{\Delta}_{\mathcal{J}})\right)_{\theta,q} \hookrightarrow \left(\dot{H}^{s,p}(\mathbb{R}_+^n), \dot{H}^{s+2,p}(\mathbb{R}_+^n)\right)_{\theta,q} = \dot{B}_{p,q}^{s+2\theta}(\mathbb{R}_+^n).$$

Now, if $q < +\infty$, we recall that $\dot{H}^{s,p}(\mathbb{R}_+^n) \cap \dot{D}_p^s(\dot{\Delta}_{\mathcal{J}}) = \dot{D}_p^s(\Delta_{\mathcal{J}})$ is a dense subspace of $(\dot{H}^{s,p}(\mathbb{R}_+^n), \dot{D}_p^s(\dot{\Delta}_{\mathcal{J}}))_{\theta,q}$ by [BL76, Theorem 3.4.2], so that by continuity of the trace operator,

$$\left(\dot{H}^{s,p}(\mathbb{R}_+^n), \dot{D}_p^s(\dot{\Delta}_{\mathcal{J}})\right)_{\theta,q} \hookrightarrow \dot{B}_{p,q,\mathcal{J}}^{s+2\theta}(\mathbb{R}_+^n).$$

The case $q = +\infty$ will be done in later steps.

Step 2: The reverse embedding when $s + 2\theta \in (-1 + 1/p, 1/p)$. Let $f \in \dot{D}_p^s(\Delta_{\mathcal{J}})$, then for all $t > 0$

$$(3.12) \quad f = e^{t\Delta_{\mathcal{J}}} f + \dot{\Delta}_{\mathcal{J}} \int_0^t \tau e^{\tau\Delta_{\mathcal{J}}} f \frac{d\tau}{\tau} =: b + a$$

with obviously $f \in \dot{D}_p^s(\Delta_{\mathcal{J}}) \subset \dot{H}^{s,p}(\mathbb{R}_+^n) + \dot{D}_p^s(\dot{\Delta}_{\mathcal{J}})$ and by definition of the K -functional, we obtain

$$K\left(t, f, \dot{H}^{s,p}(\mathbb{R}_+^n), \dot{D}_p^s(\dot{\Delta}_{\mathcal{J}})\right) \leq \|a\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} + t \|\Delta_{\mathcal{J}} b\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)}.$$

So, as in the proof of [DHMT21, Proposition 4.12], we apply [DHMT21, Lemma 2.11] so that

$$(3.13) \quad \|f\|_{(\dot{H}^{s,p}(\mathbb{R}_+^n), \dot{D}_p^s(\dot{\Delta}_{\mathcal{J}}))_{\theta,q}} \leq \frac{1 + \theta}{\theta} \left(\int_0^{+\infty} \|t^{1-\theta} \Delta_{\mathcal{J}} e^{t\Delta_{\mathcal{J}}} f\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)}^q \frac{dt}{t} \right)^{\frac{1}{q}}.$$

Now, for $0 < \theta < \varepsilon$, such that $s + 2\theta < s + 2\varepsilon < 1/p$ we want to bound the L_*^q -norm of $\|t^{1-\theta} \Delta_{\mathcal{J}} e^{t\Delta_{\mathcal{J}}} f\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)}$ by the L_*^q -norm of the K -functional associated with the real interpolation space $(\dot{H}^{s,p}(\mathbb{R}_+^n), \dot{H}^{s+2\varepsilon,p}(\mathbb{R}_+^n))_{\frac{\theta}{\varepsilon},q}$.

Notice that Lemmas 1.8 and 3.6 do imply that $e^{t\Delta_{\mathcal{J}}} f \in \dot{H}^{s+2\varepsilon,p}(\mathbb{R}_+^n)$ for $0 < \tilde{\varepsilon} < 1$:

$$\begin{aligned} \|e^{t\Delta_{\mathcal{J}}} f\|_{\dot{H}^{s+2\varepsilon,p}(\mathbb{R}_+^n)} &\lesssim_{p,s,n}^{\tilde{\varepsilon}} \|e^{t\Delta_{\mathcal{J}}} f\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)}^{(1-\tilde{\varepsilon})} \|e^{t\Delta_{\mathcal{J}}} f\|_{\dot{H}^{s+2,p}(\mathbb{R}_+^n)}^{\tilde{\varepsilon}} \\ &\lesssim_{p,s,n}^{\tilde{\varepsilon}} \|f\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)}^{(1-\tilde{\varepsilon})} \|\Delta_{\mathcal{J}} f\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)}^{\tilde{\varepsilon}}. \end{aligned}$$

Hence, by (3.12), $f \in \dot{H}^{s,p}(\mathbb{R}_+^n) + \dot{H}^{s+2\varepsilon,p}(\mathbb{R}_+^n)$ for all $\tilde{\varepsilon} \in (0, 1)$.

Let $(\tilde{a}, \tilde{b}) \in \dot{H}^{s,p}(\mathbb{R}_+^n) \times \dot{H}^{s+2\varepsilon,p}(\mathbb{R}_+^n)$, such that $f = \tilde{a} + \tilde{b}$, the fact that $f \in \dot{D}_p^s(\Delta_{\mathcal{J}})$ implies $\tilde{a}, \tilde{b} \in [\dot{H}^{s,p} \cap \dot{H}^{s+2\varepsilon,p}](\mathbb{R}_+^n)$. By Lemma 3.7, since $s + 2\varepsilon < 1/p$, we have

$$\tilde{b} \in \dot{D}_p^s((-\Delta_{\mathcal{J}})^\varepsilon).$$

Therefore, since the semigroup $(e^{t\Delta_{\mathcal{J}}})_{t>0}$ is analytic, by the use of Lemma 3.7, we have

$$\begin{aligned} \|t^{1-\theta} \Delta_{\mathcal{J}} e^{t\Delta_{\mathcal{J}}} f\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} &\leq \|t^{1-\theta} \Delta_{\mathcal{J}} e^{t\Delta_{\mathcal{J}}} \tilde{a}\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} + \|t^{1-\theta} \Delta_{\mathcal{J}} e^{t\Delta_{\mathcal{J}}} \tilde{b}\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} \\ &\lesssim_{n,p,s} t^{-\theta} \left(\|\tilde{a}\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} + t^\varepsilon \|\tilde{b}\|_{\dot{H}^{s+2\varepsilon,p}(\mathbb{R}_+^n)} \right). \end{aligned}$$

Taking the infimum of all such \tilde{a}, \tilde{b} , yields

$$\|t^{1-\theta} \Delta_{\mathcal{J}} e^{t\Delta_{\mathcal{J}}} f\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} \lesssim_{p,n,s} t^{-\theta} K(t^\varepsilon, f, \dot{H}^{s,p}(\mathbb{R}_+^n), \dot{H}^{s+2\varepsilon,p}(\mathbb{R}_+^n)).$$

Therefore one may take the L^q_* -norm on both sides, and use [Gau24b, Proposition 3.17], to obtain for all $f \in \dot{D}_p^s(\Delta_{\mathcal{J}})$,

$$\|f\|_{(\dot{H}^{s,p}(\mathbb{R}_+^n), \dot{D}_p^s(\Delta_{\mathcal{J}}))_{\theta,q}} \lesssim_{p,s,n,\theta,\varepsilon} \|f\|_{(\dot{H}^{s,p}(\mathbb{R}_+^n), \dot{H}^{s+2\varepsilon,p}(\mathbb{R}_+^n))_{\frac{\theta}{\varepsilon},q}} \sim_{p,s,n,\theta,\varepsilon} \|f\|_{\dot{B}_{p,q}^{s+2\theta}(\mathbb{R}_+^n)}.$$

Then, thanks to Step 1, one has for all $f \in \dot{D}_p^s(\Delta_{\mathcal{J}})$,

$$\|f\|_{(\dot{H}^{s,p}(\mathbb{R}_+^n), \dot{D}_p^s(\Delta_{\mathcal{J}}))_{\theta,q}} \sim_{p,s,n,\theta,\varepsilon} \|f\|_{\dot{B}_{p,q}^{s+2\theta}(\mathbb{R}_+^n)}.$$

If $q \in [1, +\infty)$, the result follows from [Gau24b, Corollary 3.18], since $C_c^\infty(\mathbb{R}_+^n)$ is a subspace of $\dot{D}_p^s(\Delta_{\mathcal{J}})$. The case $q = +\infty$ is obtained via the reiteration theorem [BL76, Theorem 3.5.3].

Step 3: The reverse embedding when $s + 2\theta \in (1/p, 2 + 1/p)$, $\mathcal{J} = \mathcal{D}$. Provided $f \in Y_{\mathcal{D}}$, as introduced before Proposition 3.8, we may reproduce above Step 2 up to (3.13). From there, for $0 < \eta < \theta$ such that $1/p < s + 2\eta < s + 2\theta$, we want to prove that one can bound (3.13) by the L^q_* -norm of the K -functional associated with the real interpolation space $(\dot{H}^{s+2\eta,p}(\mathbb{R}_+^n), \dot{H}^{s+2,p}(\mathbb{R}_+^n))_{\frac{\theta-\eta}{1-\eta},q}$.

Since $f \in Y_{\mathcal{D}} \subset \dot{H}^{s+2\eta,p}(\mathbb{R}_+^n) + \dot{H}^{s+2,p}(\mathbb{R}_+^n)$, for $(a, b) \in \dot{H}^{s+2\eta,p}(\mathbb{R}_+^n) \times \dot{H}^{s+2,p}(\mathbb{R}_+^n)$ such that $f = a + b$, we get

$$a = f - b \in \dot{H}^{s+2\eta,p}(\mathbb{R}_+^n) \cap (\dot{H}^{s+2,p}(\mathbb{R}_+^n) + Y_{\mathcal{D}}) \subset \dot{H}^{s+2\eta,p}(\mathbb{R}_+^n) \cap \dot{H}^{s+2,p}(\mathbb{R}_+^n),$$

and the same argument leads to $b \in \dot{H}^{s+2\eta,p}(\mathbb{R}_+^n) \cap \dot{H}^{s+2,p}(\mathbb{R}_+^n)$. From Lemma 3.10, we have

$$f = \mathcal{P}_{\mathcal{D}} f = \mathcal{P}_{\mathcal{D}} a + \mathcal{P}_{\mathcal{D}} b$$

where $\mathcal{P}_{\mathcal{D}} a, \mathcal{P}_{\mathcal{D}} b \in \dot{H}_{\mathcal{D}}^{s+2\eta,p}(\mathbb{R}_+^n) \cap \dot{H}^{s+2,p}(\mathbb{R}_+^n)$ with the estimates

$$\|\mathcal{P}_{\mathcal{D}} a\|_{\dot{H}^{s+2\eta,p}(\mathbb{R}_+^n)} \lesssim_{p,s,\eta,n} \|a\|_{\dot{H}^{s+2\eta,p}(\mathbb{R}_+^n)} \quad \text{and} \quad \|\mathcal{P}_{\mathcal{D}} b\|_{\dot{H}^{s+2,p}(\mathbb{R}_+^n)} \lesssim_{p,s,n} \|b\|_{\dot{H}^{s+2,p}(\mathbb{R}_+^n)}.$$

Therefore, by above estimate, analyticity of the semigroup $(e^{t\Delta_{\mathcal{D}}})_{t>0}$, and Lemma 3.7, we are able to obtain

$$\begin{aligned} \|t^{1-\theta} \Delta_{\mathcal{D}} e^{t\Delta_{\mathcal{D}}} f\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} &\leq \|t^{1-\theta} \Delta_{\mathcal{D}} e^{t\Delta_{\mathcal{D}}} \mathcal{P}_{\mathcal{D}} a\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} + \|t^{1-\theta} \Delta_{\mathcal{D}} e^{t\Delta_{\mathcal{D}}} \mathcal{P}_{\mathcal{D}} b\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} \\ &\lesssim_{n,p,s} t^{-(\theta-\eta)} \|\mathcal{P}_{\mathcal{D}} a\|_{\dot{H}^{s+2\eta,p}(\mathbb{R}_+^n)} + t^{1-\theta} \|\mathcal{P}_{\mathcal{D}} b\|_{\dot{H}^{s+2,p}(\mathbb{R}_+^n)} \\ &\lesssim_{n,p,s} t^{-(\theta-\eta)} \left(\|a\|_{\dot{H}^{s+2\eta,p}(\mathbb{R}_+^n)} + t^{1-\eta} \|b\|_{\dot{H}^{s+2,p}(\mathbb{R}_+^n)} \right). \end{aligned}$$

Taking the infimum of all such couples (a, b) , yields

$$\left\| t^{1-\theta} \Delta_{\mathcal{D}} e^{t\Delta_{\mathcal{D}}} f \right\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} \lesssim_{p,n,s} t^{-(\theta-\eta)} K \left(t^{1-\eta}, f, \dot{H}^{s+2\eta,p}(\mathbb{R}_+^n), \dot{H}^{s+2,p}(\mathbb{R}_+^n) \right).$$

As in the Step 2, one may take the L_*^q -norm on both sides, and use [Gau24b, Proposition 3.17], to obtain for all $f \in Y_{\mathcal{D}}$,

$$\|f\|_{(\dot{H}^{s,p}(\mathbb{R}_+^n), \dot{D}_p^s(\Delta_{\mathcal{D}}))_{\theta,q}} \sim_{p,s,n,\theta,\eta} \|f\|_{(\dot{H}^{s+2\eta,p}(\mathbb{R}_+^n), \dot{H}^{s+2,p}(\mathbb{R}_+^n))_{\frac{\theta-\eta}{1-\eta},q}} \sim_{p,s,n,\theta,\eta} \|f\|_{\dot{B}_{p,q}^{s+2\theta}(\mathbb{R}_+^n)}.$$

If $q \in [1, +\infty)$, the result follows from Proposition 3.8. The case $q = +\infty$ is obtained via the application of the reiteration theorem [BL76, Theorem 3.5.3], by means of Lemma 3.10.

Step 4: The reverse embedding $s + 2\theta \in [1/p, 1 + 1/p)$, $\mathcal{J} = \mathcal{N}$. One may pick $f \in Y_{\mathcal{N}}$ so that, as before, we can reproduce above Step 2 up to (3.13). From there, for $0 < \eta < \theta < \varepsilon$ such that $1/p < s + 2\eta < s + 2\theta < s + 2\varepsilon < 1 + 1/p$, we want to prove that one can bound (3.13) by the L_*^q -norm of the K -functional associated with the interpolation space $(\dot{H}^{s+2\eta,p}(\mathbb{R}_+^n), \dot{H}^{s+2\varepsilon,p}(\mathbb{R}_+^n))_{\frac{\theta-\eta}{\varepsilon-\eta},q}$.

Since $f \in Y_{\mathcal{D}} \subset \dot{H}^{s+2\eta,p}(\mathbb{R}_+^n) + \dot{H}^{s+2\varepsilon,p}(\mathbb{R}_+^n)$, for $(a, b) \in \dot{H}^{s+2\eta,p}(\mathbb{R}_+^n) \times \dot{H}^{s+2\varepsilon,p}(\mathbb{R}_+^n)$ such that $f = a + b$, we get

$$b = f - a \in \dot{H}^{s+2\eta,p}(\mathbb{R}_+^n) \cap (\dot{H}^{s+2\varepsilon,p}(\mathbb{R}_+^n) + Y_{\mathcal{N}}) \subset \dot{H}^{s+2\eta,p}(\mathbb{R}_+^n) \cap \dot{H}^{s+2\varepsilon,p}(\mathbb{R}_+^n).$$

By Proposition 3.9, there exists sequences $(a_j)_{j \in \mathbb{N}}, (b_j)_{j \in \mathbb{N}} \subset Y_{\mathcal{N}}$ such that

$$\|a_j - a\|_{\dot{H}^{s+2\eta,p}(\mathbb{R}_+^n)} + \|b_j - b\|_{[\dot{H}^{s+2\eta,p} \cap \dot{H}^{s+2\varepsilon,p}](\mathbb{R}_+^n)} \xrightarrow{j \rightarrow +\infty} 0.$$

Therefore, the analyticity of the semigroup $(e^{t\Delta_{\mathcal{N}}})_{t>0}$ and Lemma 3.7 works together to deliver

$$\begin{aligned} \left\| t\Delta_{\mathcal{N}} e^{t\Delta_{\mathcal{N}}} a_j \right\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} &\lesssim_{p,s,n,\eta} t^\eta \|a_j\|_{\dot{H}^{s+2\eta,p}(\mathbb{R}_+^n)}, \\ \left\| t\Delta_{\mathcal{N}} e^{t\Delta_{\mathcal{N}}} b_j \right\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} &\lesssim_{p,s,n,\varepsilon} t^\varepsilon \|b_j\|_{\dot{H}^{s+2\varepsilon,p}(\mathbb{R}_+^n)}, \end{aligned}$$

so that taking limits, it yields

$$(3.14) \quad \begin{aligned} \left\| t\Delta_{\mathcal{N}} e^{t\Delta_{\mathcal{N}}} a \right\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} &\lesssim_{p,s,n,\eta} t^\eta \|a\|_{\dot{H}^{s+2\eta,p}(\mathbb{R}_+^n)}, \\ \left\| t\Delta_{\mathcal{N}} e^{t\Delta_{\mathcal{N}}} b \right\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} &\lesssim_{p,s,n,\varepsilon} t^\varepsilon \|b\|_{\dot{H}^{s+2\varepsilon,p}(\mathbb{R}_+^n)}. \end{aligned}$$

Therefore, by the estimates (3.14), the following holds

$$\left\| t^{1-\theta} \Delta_{\mathcal{N}} e^{t\Delta_{\mathcal{N}}} f \right\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} \lesssim_{n,p,s,\eta,\varepsilon} t^{-(\theta-\eta)} \left(\|a\|_{\dot{H}^{s+2\eta,p}(\mathbb{R}_+^n)} + t^{\varepsilon-\eta} \|b\|_{\dot{H}^{s+2\varepsilon,p}(\mathbb{R}_+^n)} \right).$$

From there, we can take the infimum of all such couples (a, b) , and we see that

$$\left\| t^{1-\theta} \Delta_{\mathcal{N}} e^{t\Delta_{\mathcal{N}}} f \right\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} \lesssim_{p,n,s} t^{-(\theta-\eta)} K \left(t^{\varepsilon-\eta}, f, \dot{H}^{s+2\eta,p}(\mathbb{R}_+^n), \dot{H}^{s+2\varepsilon,p}(\mathbb{R}_+^n) \right).$$

As in the Step 2, one may take the L^q_* -norm on both sides, and use [Gau24b, Proposition 3.17], to obtain for all $f \in Y_{\mathcal{N}}$,

$$\begin{aligned} \|f\|_{(\dot{H}^{s,p}(\mathbb{R}_+^n), \dot{D}_p^s(\dot{\Delta}_{\mathcal{N}}))_{\theta,q}} &\sim_{p,s,n,\theta,\eta,\varepsilon} \|f\|_{(\dot{H}^{s+2\eta,p}(\mathbb{R}_+^n), \dot{H}^{s+2\varepsilon,p}(\mathbb{R}_+^n))_{\frac{\theta-\eta}{\varepsilon-\eta},q}} \\ &\sim_{p,s,n,\theta,\eta,\varepsilon} \|f\|_{\dot{B}_{p,q}^{s+2\theta}(\mathbb{R}_+^n)}. \end{aligned}$$

If $q \in [1, +\infty)$, the result follows from Proposition 3.8. The case $q = +\infty$ is obtained via the application of the reiteration theorem [BL76, Theorem 3.5.3]. The case $s = 1/p$ follows from reiteration theorem [BL76, Theorem 3.5.3] between Step 2 and this one.

Step 5: The reverse embedding $s + 2\theta \in (1 + 1/p, 2 + 1/p)$, $\mathcal{J} = \mathcal{N}$. For $f \in Y_{\mathcal{N}}$, we reproduce again the Step 2 up to (3.13). Now let $0 < \eta < \theta$ such that $1 + 1/p < s + 2\eta < s + 2\theta < 2 + 1/p$, we want to achieve the same estimate obtained at the end of Step 3.

Since $f \in Y_{\mathcal{N}} \subset \dot{H}^{s+2\eta,p}(\mathbb{R}_+^n) + \dot{H}^{s+2,p}(\mathbb{R}_+^n)$, for $(a, b) \in \dot{H}^{s+2\eta,p}(\mathbb{R}_+^n) \times \dot{H}^{s+2,p}(\mathbb{R}_+^n)$ such that $f = a + b$, we get

$$b = f - a \in \dot{H}^{s+2\eta,p}(\mathbb{R}_+^n) \cap (\dot{H}^{s+2,p}(\mathbb{R}_+^n) + Y_{\mathcal{N}}) \subset \dot{H}^{s+2\eta,p}(\mathbb{R}_+^n) \cap \dot{H}^{s+2,p}(\mathbb{R}_+^n).$$

We want to fall in the expected homogeneous domains, i.e. to get back the Neumann boundary condition, to do so, we use Lemma 3.10, and we get

$$f = \mathcal{P}_{\mathcal{N}}f = \mathcal{P}_{\mathcal{N}}a + \mathcal{P}_{\mathcal{N}}b,$$

with estimates

$$\|\mathcal{P}_{\mathcal{N}}a\|_{\dot{H}^{s+2\eta,p}(\mathbb{R}_+^n)} \lesssim_{p,n,s,\eta} \|a\|_{\dot{H}^{s+2\eta,p}(\mathbb{R}_+^n)} \quad \text{and} \quad \|\mathcal{P}_{\mathcal{N}}b\|_{\dot{H}^{s+2,p}(\mathbb{R}_+^n)} \lesssim_{p,n,s,\eta} \|b\|_{\dot{H}^{s+2,p}(\mathbb{R}_+^n)}.$$

By Proposition 3.9, there exists sequences $(\mathbf{a}_j)_{j \in \mathbb{N}}, (\mathbf{b}_j)_{j \in \mathbb{N}} \subset Y_{\mathcal{N}}$ such that

$$\|\mathbf{a}_j - \mathcal{P}_{\mathcal{N}}a\|_{\dot{H}^{s+2\eta,p}(\mathbb{R}_+^n)} + \|\mathbf{b}_j - \mathcal{P}_{\mathcal{N}}b\|_{[\dot{H}^{s+2\eta,p} \cap \dot{H}^{s+2,p}](\mathbb{R}_+^n)} \xrightarrow{j \rightarrow +\infty} 0.$$

As in Step 4, we obtain

$$\begin{aligned} \|t\Delta_{\mathcal{N}}e^{t\Delta_{\mathcal{N}}}\mathcal{P}_{\mathcal{N}}a\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} &\lesssim_{p,s,n,\eta} t^\eta \|a\|_{\dot{H}^{s+2\eta,p}(\mathbb{R}_+^n)}, \\ \|t\Delta_{\mathcal{N}}e^{t\Delta_{\mathcal{N}}}\mathcal{P}_{\mathcal{N}}b\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} &\lesssim_{p,s,n} t \|b\|_{\dot{H}^{s+2,p}(\mathbb{R}_+^n)}. \end{aligned}$$

Therefore, by the estimates above, the following estimate holds

$$\|t^{1-\theta}\Delta_{\mathcal{N}}e^{t\Delta_{\mathcal{N}}}f\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} \lesssim_{n,p,s} t^{-(\theta-\eta)} \left(\|a\|_{\dot{H}^{s+2\eta,p}(\mathbb{R}_+^n)} + t^{1-\eta} \|b\|_{\dot{H}^{s+2,p}(\mathbb{R}_+^n)} \right).$$

Finally, one may finish the present Step 5 with the same arguments present in Step 3.

Step 6: The case $\mathcal{J} = \mathcal{H}$. Let $k \in \llbracket 0, n \rrbracket$, from Lemma 3.6, we deduce that the following holds with equivalence of norms

$$\dot{D}_p^s(\dot{\Delta}_{\mathcal{H}}, \mathbf{n}, \Lambda^k) \simeq \dot{D}_p^s(\dot{\Delta}_{\mathcal{D}})^{\binom{n-1}{k-1}} \times \dot{D}_p^s(\dot{\Delta}_{\mathcal{N}})^{\binom{n-1}{k}}$$

The result is then immediate, by all above steps. The case of the Hodge Laplacian with generalized tangential boundary conditions admits a similar proof. \square

Finally, we want to compute interpolation spaces for the Hodge–Stokes and the Hodge–Maxwell operators. To do so, we set for all $p \in (1, +\infty)$, $q \in [1, +\infty]$, $s \in (-1 + 1/p, 2 + 1/p)$, such that $(\mathcal{C}_{s,p,q})$ is satisfied, provided, $k \in \llbracket 0, n \rrbracket$,

$$(3.15) \quad \dot{B}_{p,q,\mathcal{H}_n}^{s,\sigma}(\mathbb{R}_+^n, \Lambda^k) := \begin{cases} \dot{B}_{p,q,n}^{s,\sigma}(\mathbb{R}_+^n, \Lambda^k), & s < \frac{1}{p}, \\ \left\{ u \in \dot{B}_{p,q,\mathcal{H}_n}^s(\mathbb{R}_+^n, \Lambda^k) \mid \delta u = 0 \right\}, & \frac{1}{p} < s < 2 + \frac{1}{p}, s \neq 1 + 1/p. \end{cases}$$

$$(3.16) \quad \dot{B}_{p,q,\mathcal{H}_n}^{s,\gamma}(\mathbb{R}_+^n, \Lambda^k) := \begin{cases} \dot{B}_{p,q}^{s,\gamma}(\mathbb{R}_+^n, \Lambda^k), & s < \frac{1}{p}, \\ \left\{ u \in \dot{B}_{p,q,\mathcal{H}_n}^s(\mathbb{R}_+^n, \Lambda^k) \mid du = 0 \right\}, & \frac{1}{p} < s < 2 + \frac{1}{p}, s \neq 1 + 1/p. \end{cases}$$

One may build similarly $\dot{B}_{p,q,\mathcal{H}_t}^{s,\sigma}(\mathbb{R}_+^n, \Lambda^k)$ and $\dot{B}_{p,q,\mathcal{H}_t}^{s,\gamma}(\mathbb{R}_+^n, \Lambda^k)$ replacing $(\mathbf{n}, \sigma, \gamma, d, \delta)$ by $(\mathbf{t}, \gamma, \sigma, \delta, d)$.

PROPOSITION 3.13. — *Let $p \in (1, +\infty)$, $q \in [1, +\infty]$, and $s \in (-1 + 1/p, 1/p)$. For all $\theta \in (0, 1)$ such that $(\mathcal{C}_{s+2\theta,p,q})$ is satisfied, one has*

$$(3.17) \quad \left(\dot{H}_{\mathbf{n},\sigma}^{s,p}(\mathbb{R}_+^n), \dot{D}_p^s(\mathring{\mathbb{A}}_{\mathcal{H},\mathbf{n}}) \right)_{\theta,q} = \dot{B}_{p,q,\mathcal{H}_n}^{s+2\theta,\sigma}(\mathbb{R}_+^n),$$

$$(3.18) \quad \left(\dot{H}_\gamma^{s,p}(\mathbb{R}_+^n), \dot{D}_p^s(\mathring{\mathbb{M}}_{\mathcal{H},\mathbf{n}}) \right)_{\theta,q} = \dot{B}_{p,q,\mathcal{H}_n}^{s+2\theta,\gamma}(\mathbb{R}_+^n),$$

with equivalence of norms, whenever $s + 2\theta \neq 1/p, 1 + 1/p$.

The same result holds replacing $(\mathbf{n}, \sigma, \gamma, \mathbb{A}, \mathbb{M})$ by $(\mathbf{t}, \gamma, \sigma, \mathbb{M}, \mathbb{A})$.

Proof. — We only prove (3.17), other equalities have the same proof.

Step 1: We start with [Gau24b, Proposition 3.17] which yields the embedding

$$\left(\dot{H}_{\mathbf{n},\sigma}^{s,p}(\mathbb{R}_+^n), \dot{D}_p^s(\mathring{\mathbb{A}}_{\mathcal{H},\mathbf{n}}) \right)_{\theta,q} \hookrightarrow \left(\dot{H}^{s,p}(\mathbb{R}_+^n), \dot{H}^{s+2,p}(\mathbb{R}_+^n) \right)_{\theta,q} = \dot{B}_{p,q}^{s+2\theta}(\mathbb{R}_+^n).$$

Now, if $q < +\infty$, we recall that $\dot{H}_{\mathbf{n},\sigma}^{s,p}(\mathbb{R}_+^n) \cap \dot{d}_p^s(\mathring{\mathbb{A}}_{\mathcal{H},\mathbf{n}})$ is a dense subspace of $(\dot{H}_{\mathbf{n},\sigma}^{s,p}(\mathbb{R}_+^n), \dot{d}_p^s(\mathring{\mathbb{A}}_{\mathcal{H},\mathbf{n}}))_{\theta,q}$ by [BL76, Theorem 3.4.2], so that by continuity of traces,

$$\left(\dot{H}_{\mathbf{n},\sigma}^{s,p}(\mathbb{R}_+^n), \dot{D}_p^s(\mathring{\mathbb{A}}_{\mathcal{H},\mathbf{n}}) \right)_{\theta,q} \hookrightarrow \dot{B}_{p,q,\mathcal{H}_n}^{s+2\theta,\sigma}(\mathbb{R}_+^n) \hookrightarrow \dot{B}_{p,q,\mathcal{H}_n}^{s+2\theta}(\mathbb{R}_+^n).$$

Again density of $\dot{H}_{\mathbf{n},\sigma}^{s,p}(\mathbb{R}_+^n) \cap \dot{D}_p^s(\mathring{\mathbb{A}}_{\mathcal{H},\mathbf{n}})$ yields $\delta f = 0$ for all $f \in (\dot{H}_{\mathbf{n},\sigma}^{s,p}(\mathbb{R}_+^n), \dot{D}_p^s(\mathring{\mathbb{A}}_{\mathcal{H},\mathbf{n}}))_{\theta,q}$. The case $q = +\infty$ is left to the end of Step 3.

Step 2: We want to extend the range of exponents for the boundedness of \mathbb{P} , and get a density result.

Let $f \in \dot{D}_p^s(\Delta_{\mathcal{H},n}) \subset \dot{B}_{p,q,\mathcal{H}_n}^{s+2\theta}(\mathbb{R}_+^n)$, we have $\mathbb{P}f \in \dot{D}_p^s(\Delta_{\mathcal{H},n})$ and by Proposition 3.12, [DHMT21, Proposition 2.12], Corollary 2.34 and Theorem 2.33, we obtain successively

$$\begin{aligned} \|\mathbb{P}f\|_{\dot{B}_{p,q}^{s+2\theta}(\mathbb{R}_+^n)} &\lesssim_{p,s,n,\theta} \left(\int_0^{+\infty} \|t^{1-\theta} \Delta_{\mathcal{H},n} e^{t\Delta_{\mathcal{H},n}} \mathbb{P}f\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)}^q \frac{dt}{t} \right)^{\frac{1}{q}} \\ &\lesssim_{p,s,n,\theta} \left(\int_0^{+\infty} \|t^{1-\theta} \mathbb{P} \Delta_{\mathcal{H},n} e^{t\Delta_{\mathcal{H},n}} f\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)}^q \frac{dt}{t} \right)^{\frac{1}{q}} \\ &\lesssim_{p,s,n,\theta} \left(\int_0^{+\infty} \|t^{1-\theta} \Delta_{\mathcal{H},n} e^{t\Delta_{\mathcal{H},n}} f\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)}^q \frac{dt}{t} \right)^{\frac{1}{q}} \\ &\lesssim_{p,s,n,\theta} \|f\|_{\dot{B}_{p,q}^{s+2\theta}(\mathbb{R}_+^n)}. \end{aligned}$$

From above estimates, if $q < +\infty$, by density of $\dot{D}_p^s(\Delta_{\mathcal{H},n})$ in $\dot{B}_{p,q,\mathcal{H}_n}^{s+2\theta}(\mathbb{R}_+^n)$, we have that

$$\mathbb{P} : \dot{B}_{p,q,\mathcal{H}_n}^{s+2\theta}(\mathbb{R}_+^n) \longrightarrow \dot{B}_{p,q,\mathcal{H}_n}^{s+2\theta}(\mathbb{R}_+^n)$$

extends uniquely as a bounded projection on $\dot{B}_{p,q,\mathcal{H}_n}^{s+2\theta}(\mathbb{R}_+^n)$ with range $\dot{B}_{p,q,\mathcal{H}_n}^{s+2\theta,\sigma}(\mathbb{R}_+^n)$. The result still holds for $q = +\infty$, by above Step 1, the reiteration theorem [BL76, Theorem 3.5.3] and Proposition 3.12.

In particular, $\dot{D}_p^s(\mathbb{A}_{\mathcal{H},n}) = \mathbb{P}\dot{D}_p^s(\Delta_{\mathcal{H},n})$ is dense in $\dot{B}_{p,q,\mathcal{H}_n}^{s+2\theta,\sigma}(\mathbb{R}_+^n)$, when $q < +\infty$.

Step 3: For the reverse embedding. Let $f \in \dot{D}_p^s(\mathbb{A}_{\mathcal{H},n}) \subset \dot{B}_{p,q,\mathcal{H}_n}^{s+2\theta}(\mathbb{R}_+^n)$, and note that $\dot{B}_{p,q,\mathcal{H}_n}^{s+2\theta}(\mathbb{R}_+^n) = (\dot{H}^{s,p}(\mathbb{R}_+^n), \dot{D}_p^s(\mathring{\Delta}_{\mathcal{H},n}))_{\theta,q} \subset \dot{H}^{s,p}(\mathbb{R}_+^n) + \dot{D}_p^s(\mathring{\Delta}_{\mathcal{H},n})$

If we let $(a, b) \in \dot{H}^{s,p}(\mathbb{R}_+^n) \times \dot{D}_p^s(\mathring{\Delta}_{\mathcal{H},n})$ such that $f = a + b$, by Proposition 3.11, it is given that

$$b = f - a \in (\dot{D}_p^s(\Delta_{\mathcal{H},n}) + \dot{H}^{s,p}(\mathbb{R}_+^n)) \cap \dot{D}_p^s(\mathring{\Delta}_{\mathcal{H},n}) \subset \dot{D}_p^s(\Delta_{\mathcal{H},n})$$

and for the same reason $a \in \dot{D}_p^s(\Delta_{\mathcal{H},n})$. Therefore,

$$f = \mathbb{P}f = \mathbb{P}a + \mathbb{P}b \in \dot{H}_{n,\sigma}^{s,p}(\mathbb{R}_+^n) + \dot{D}_p^s(\mathring{\mathbb{A}}_{\mathcal{H},n}).$$

By (2.15) and Corollary 2.34, we have

$$\begin{aligned} &\|t^{1-\theta} \mathring{\mathbb{A}}_{\mathcal{H},n} e^{-t\mathring{\mathbb{A}}_{\mathcal{H},n}} f\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} \\ &\leq \|t^{1-\theta} \mathring{\mathbb{A}}_{\mathcal{H},n} e^{-t\mathring{\mathbb{A}}_{\mathcal{H},n}} \mathbb{P}a\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} \|t^{1-\theta} \mathring{\mathbb{A}}_{\mathcal{H},n} e^{-t\mathring{\mathbb{A}}_{\mathcal{H},n}} \mathbb{P}b\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} \\ &\lesssim_{p,n,s} \|t^{1-\theta} \mathbb{P} \Delta_{\mathcal{H},n} e^{t\Delta_{\mathcal{H},n}} a\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} + \|t^{1-\theta} \mathbb{P} \Delta_{\mathcal{H},n} e^{t\Delta_{\mathcal{H},n}} b\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)}. \end{aligned}$$

From there, we use analyticity of the semigroup, boundedness of \mathbb{P} given by Theorem 2.33, to obtain

$$\|t^{1-\theta} \mathring{\mathbb{A}}_{\mathcal{H},n} e^{-t\mathring{\mathbb{A}}_{\mathcal{H},n}} f\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} \lesssim_{p,n,s} t^{-\theta} \|a\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} + t^{1-\theta} \|\Delta_{\mathcal{H},n} b\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)}.$$

Taking the infimum on all such pairs (a, b) yields

$$\left\| t^{1-\theta} \mathring{\mathbb{A}}_{\mathcal{H},n} e^{-t\mathbb{A}_{\mathcal{H},n}} f \right\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} \lesssim_{p,n,s} t^{-\theta} K(t, f, \dot{H}^{s,p}(\mathbb{R}_+^n), \dot{D}_p^s(\mathring{\Delta}_{\mathcal{H},n})).$$

One may take the L_*^q -norm of the inequality above, then applies [DHMT21, Proposition 2.12] and Proposition 3.12, to deduce that

$$\begin{aligned} \|f\|_{(\dot{H}_{n,\sigma}^{s,p}(\mathbb{R}_+^n), \dot{d}_p^s(\mathring{\mathbb{A}}_{\mathcal{H},n}))_{\theta,q}} &\lesssim_{\theta} \left(\int_0^{+\infty} \|t^{1-\theta} \Delta_{\mathcal{H},n} e^{t\Delta_{\mathcal{H},n}} \mathbb{P} f\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)}^q \frac{dt}{t} \right) \\ &\lesssim_{p,n,s,\theta} \|f\|_{(\dot{H}^{s,p}(\mathbb{R}_+^n), \dot{D}_p^s(\mathring{\Delta}_{\mathcal{H},n}))_{\theta,q}} \\ &\lesssim_{p,n,s,\theta} \|f\|_{\dot{B}_{p,q}^{s+2\theta}(\mathbb{R}_+^n)}. \end{aligned}$$

With Step 1, one has for all $f \in \dot{D}_p^s(\mathbb{A}_{\mathcal{H},n})$,

$$\|f\|_{(\dot{H}_{n,\sigma}^{s,p}(\mathbb{R}_+^n), \dot{D}_p^s(\mathring{\mathbb{A}}_{\mathcal{H},n}))_{\theta,q}} \sim_{p,n,s,\theta} \|f\|_{\dot{B}_{p,q}^{s+2\theta}(\mathbb{R}_+^n)}.$$

If $q < +\infty$, then the end of the Step 2 above, and [DHMT21, Lemma 2.10] allows to conclude by density. If $q = +\infty$, the result follows from the reiteration theorem [BL76, Theorem 3.5.3] and the boundedness and the range of \mathbb{P} in Step 2 (use a retraction argument [BL76, Theorem 6.4.2]). \square

The Step 2 from the proof above leads to the immediate following corollary.

COROLLARY 3.14. — *Let $p \in (1, +\infty)$, $q \in [1, +\infty]$, $s \in (-1 + 1/p, 2 + 1/p)$, such that $s \notin \mathbb{N} + \frac{1}{p}$, and $(\mathcal{C}_{s,p,q})$ is satisfied. Then,*

$$\begin{aligned} \mathbb{P} : \dot{B}_{p,q,\mathcal{H}_n}^s(\mathbb{R}_+^n) &\longrightarrow \dot{B}_{p,q,\mathcal{H}_n}^{s,\sigma}(\mathbb{R}_+^n), \\ [\mathbb{I} - \mathbb{P}] : \dot{B}_{p,q,\mathcal{H}_n}^s(\mathbb{R}_+^n) &\longrightarrow \dot{B}_{p,q,\mathcal{H}_n}^{s,\gamma}(\mathbb{R}_+^n), \end{aligned}$$

are both well-defined bounded linear projections, so that the following Hodge decomposition holds

$$\dot{B}_{p,q,\mathcal{H}_n}^s(\mathbb{R}_+^n) = \dot{B}_{p,q,\mathcal{H}_n}^{s,\sigma}(\mathbb{R}_+^n) \oplus \dot{B}_{p,q,\mathcal{H}_n}^{s,\gamma}(\mathbb{R}_+^n).$$

The result still holds if we replace $(\mathfrak{n}, \mathbb{P})$ by $(\mathfrak{t}, \mathbb{Q})$.

Finally, we mention without its proofs, that follows exactly the same lines, the result for interpolation spaces of the homogeneous domains with Besov spaces as an ambient function space, say, for $\theta \in (0, 1)$, $r, q \in [1, +\infty]$, $p \in (1, +\infty)$, $-1 + 1/p < s < 1/p$,

$$\dot{D}_{-\Delta_{\mathcal{H}}}^{s,p,r}(\theta, q) = \left(\dot{B}_{p,r}^s(\mathbb{R}_+^n), \dot{D}_{p,r}^s(\mathring{\Delta}_{\mathcal{H}}) \right)_{\theta,q}.$$

We are able to obtain,

PROPOSITION 3.15. — *Let $p \in (1, +\infty)$, $r \in [1, +\infty]$, $q \in [1, +\infty]$, and $s \in (-1 + 1/p, 1/p)$. For all $\theta \in (0, 1)$ such that $(\mathcal{C}_{s+2\theta,p,q})$ is satisfied, provided $\mathcal{J} \in \{\mathcal{D}, \mathcal{N}, \mathcal{H}\}$, one has*

$$\left(\dot{B}_{p,r}^s(\mathbb{R}_+^n), \dot{D}_{p,r}^s(\mathring{\Delta}_{\mathcal{J}}) \right)_{\theta,q} = \dot{B}_{p,q,\mathcal{J}}^{s+2\theta}(\mathbb{R}_+^n),$$

with equivalence of norms, whenever

- $s + 2\theta \neq 1/p$ if $\mathcal{J} = \mathcal{D}$,

- $s + 2\theta \neq 1 + 1/p$ if $\mathcal{J} = \mathcal{N}$,
- $s + 2\theta \neq 1/p, 1 + 1/p$ if $\mathcal{J} = \mathcal{H}$.

Then by means of Corollary 3.14,

PROPOSITION 3.16. — *Let $p \in (1, +\infty)$, $r \in [1, +\infty)$, $q \in [1, +\infty]$, and $s \in (-1 + 1/p, 1/p)$. For all $\theta \in (0, 1)$ such that $(\mathcal{C}_{s+2\theta,p,q})$ is satisfied, one has*

$$\begin{aligned} \left(\dot{B}_{p,r,n}^{s,\sigma}(\mathbb{R}_+^n), \dot{D}_{p,r}^s(\mathring{A}_{\mathcal{H},n}) \right)_{\theta,q} &= \dot{B}_{p,q,\mathcal{H}_n}^{s+2\theta,\sigma}(\mathbb{R}_+^n), \\ \left(\dot{B}_{p,r}^{s,\gamma}(\mathbb{R}_+^n), \dot{D}_{p,r}^s(\mathring{M}_{\mathcal{H},n}) \right)_{\theta,q} &= \dot{B}_{p,q,\mathcal{H}_n}^{s+2\theta,\gamma}(\mathbb{R}_+^n), \end{aligned}$$

with equivalence of norms, whenever $s + 2\theta \neq 1/p, 1 + 1/p$.

The same result holds replacing $(\mathbf{n}, \sigma, \gamma, \mathbb{A}, \mathbb{M})$ by $(\mathbf{t}, \gamma, \sigma, \mathbb{M}, \mathbb{A})$.

3.2. Maximal regularity for Hodge Laplacians and related operators

We will present here a direct application of Theorem 3.4, and [DHMT21, Theorem 2.20] with appropriate identification of real interpolation spaces, provided $p, r \in (1, +\infty)$, $s \in (-1 + 1/p, 1/p)$,

$$\mathring{D}_A^{s,p}(\theta, q), \mathring{D}_A^{s,p,r}(\theta, q), \theta \in (0, 1), q \in [1, +\infty] \text{ and } A \in \{-\Delta_{\mathcal{H}}, \mathbb{A}_{\mathcal{H}}, \mathbb{M}_{\mathcal{H}}\}$$

subject to either normal or tangential boundary conditions, see Propositions 3.12, 3.13, 3.15 and 3.16.

We recall that the definition of involved spaces are given in Notations 2.32, see also (3.11) and (3.15). To alleviate notations in inequalities, we drop the references to the open set \mathbb{R}_+^n .

We give first two theorems in the case where the ambient spaces is a UMD Banach space which is the case of $\dot{H}^{s,p}$ and $\dot{B}_{p,r}^s$, provided $p, r \in (1, +\infty)$, $s \in (-1 + 1/p, 2 + 1/p)$.

THEOREM 3.17. — *Let $p, q, r \in (1, +\infty)$, and for $\alpha \in (-1 + 1/q, 1/q)$ fixed, we set $\alpha_q := 1 + \alpha - 1/q$.*

Let $s \in (-1 + 1/p, 2 + 1/p)$ such that $s, s + 2\alpha_q \notin \mathbb{N} + \frac{1}{p}$, $(\mathcal{C}_{s+2\alpha_q,p,q})$ is satisfied, and let $T \in (0, +\infty]$.

For any $f \in \dot{H}^{\alpha,q}((0, T), \dot{B}_{p,r,\mathcal{H}_n}^s(\mathbb{R}_+^n, \Lambda))$, $u_0 \in \dot{B}_{p,q,\mathcal{H}_n}^{s+2\alpha_q}(\mathbb{R}_+^n, \Lambda)$, there exists a unique mild solution $u \in C_b^0([0, T], \dot{B}_{p,q,\mathcal{H}_n}^{s+2\alpha_q}(\mathbb{R}_+^n, \Lambda))$ of

$$(HHS_n) \quad \begin{cases} \partial_t u - \Delta u = f, & \text{on } (0, T) \times \mathbb{R}_+^n, \\ \nu \lrcorner du|_{\partial\mathbb{R}_+^n} = 0, & \text{on } (0, T) \times \partial\mathbb{R}_+^n, \\ \nu \lrcorner u|_{\partial\mathbb{R}_+^n} = 0, & \text{on } (0, T) \times \partial\mathbb{R}_+^n, \\ u(0) = u_0, & \text{in } \dot{B}_{p,q}^{2+s-\frac{2}{q}}(\mathbb{R}_+^n, \Lambda), \end{cases}$$

with estimate

$$\begin{aligned} \|u\|_{L^\infty((0,T), \dot{B}_{p,q}^{s+2\alpha_q})} &\lesssim_{p,q,n}^{\alpha,s} \left\| (\partial_t u, \nabla^2 u) \right\|_{\dot{H}^{\alpha,q}((0,T), \dot{B}_{p,r}^s)} \\ &\lesssim_{p,q,n}^{\alpha,s} \|f\|_{\dot{H}^{\alpha,q}((0,T), \dot{B}_{p,r}^s)} + \|u_0\|_{\dot{B}_{p,q}^{s+2\alpha_q}}. \end{aligned}$$

For all $\beta \in [0, 1]$, we also have

$$(3.19) \quad \left\| (-\partial_t)^\beta (-\Delta_{\mathcal{H},\mathbf{n}})^{1-\beta} u \right\|_{\dot{H}^{\alpha,q}((0,T),\dot{B}_{p,r}^s)} \lesssim_{p,q,n}^{s,\alpha} \|f\|_{\dot{H}^{\alpha,q}((0,T),\dot{B}_{p,r}^s)} + \|u_0\|_{\dot{B}_{p,q}^{s+2\alpha_q}}.$$

Proof. — From Theorem 2.29 we have the bounded holomorphic calculus of $-\Delta_{\mathcal{H}, \mathbf{n}}$ on $\dot{B}_{p,r,\mathcal{H}_n}^s(\mathbb{R}_+^n)$, so that we may apply Theorem 3.4 to obtain maximal regularity estimates, whereas Proposition 3.12 gives an exact description of interpolation spaces. \square

THEOREM 3.18. — *Let $p, q \in (1, +\infty)$, $s \in (-1 + 1/p, 1/p)$, and $\alpha \in (-1 + 1/q, 1/q)$ fixed, we set $\alpha_q := 1 + \alpha - 1/q$. We assume that $s + 2\alpha_q \notin \mathbb{N} + \frac{1}{p}$, $(\mathcal{C}_{s+2\alpha_q,p,q})$ is satisfied, and let $T \in (0, +\infty]$.*

For any $f \in \dot{H}^{\alpha,q}((0, T), \dot{H}_{t,\gamma}^{s,p}(\mathbb{R}_+^n, \Lambda))$, $u_0 \in \dot{B}_{p,q,\mathcal{H}_t}^{s+2\alpha_q,\gamma}(\mathbb{R}_+^n, \Lambda)$, there exists a unique mild solution $u \in C_b^0([0, T], \dot{B}_{p,q,\mathcal{H}_t}^{s+2\alpha_q,\gamma}(\mathbb{R}_+^n, \Lambda))$ of

$$(HMS_t) \quad \begin{cases} \partial_t u - \Delta u = f, & \text{on } (0, T) \times \mathbb{R}_+^n, \\ du = 0, & \text{on } (0, T) \times \mathbb{R}_+^n, \\ \nu \wedge \delta u|_{\partial \mathbb{R}_+^n} = 0, & \text{on } (0, T) \times \partial \mathbb{R}_+^n, \\ \nu \wedge u|_{\partial \mathbb{R}_+^n} = 0, & \text{on } (0, T) \times \partial \mathbb{R}_+^n, \\ u(0) = u_0, & \text{in } \dot{B}_{p,q}^{s+2\alpha_q}(\mathbb{R}_+^n, \Lambda), \end{cases}$$

with estimate

$$\begin{aligned} \|u\|_{L^\infty((0,T),\dot{B}_{p,q}^{s+2\alpha_q})} &\lesssim_{p,q,n}^{s,\alpha} \left\| (\partial_t u, \nabla^2 u) \right\|_{\dot{H}^{\alpha,q}((0,T),\dot{H}^{s,p})} \\ &\lesssim_{p,q,n}^{s,\alpha} \|f\|_{\dot{H}^{\alpha,q}((0,T),\dot{H}^{s,p})} + \|u_0\|_{\dot{B}_{p,q}^{s+2\alpha_q}}. \end{aligned}$$

For all $\beta \in [0, 1]$, we also have

$$(3.20) \quad \left\| (-\partial_t)^\beta (\mathbb{M}_{\mathcal{H},t})^{1-\beta} u \right\|_{\dot{H}^{\alpha,q}((0,T),\dot{H}^{s,p})} \lesssim_{p,q,n}^{s,\alpha} \|f\|_{\dot{H}^{\alpha,q}((0,T),\dot{H}^{s,p})} + \|u_0\|_{\dot{B}_{p,q}^{s+2\alpha_q}}.$$

Proof. — From Theorem 2.35, we have the bounded holomorphic calculus of $\mathbb{M}_{\mathcal{H},t}$ on $\dot{H}^{s,p}(\mathbb{R}_+^n)$, so that we may apply Theorem 3.4 to obtain maximal regularity estimates, whereas Proposition 3.15 gives an exact description of interpolation spaces. \square

Finally, the homogeneous Da Prato–Grisvard Theorem 3.19 yields our (almost) last L^q -maximal regularity theorem.

THEOREM 3.19. — *Let $p \in (1, +\infty)$, $q \in [1, +\infty)$, $s \in (-1 + 1/p, 1/p + 2/q)$, such that $s, s + 2 - 2/q \notin \mathbb{N} + \frac{1}{p}$ and $(\mathcal{C}_{s+2-2/q,p,q})$ is satisfied and let $T \in (0, +\infty]$.*

For any

$$f \in L^q \left((0, T), \dot{B}_{p,q,\mathcal{H}_n}^{s,\sigma}(\mathbb{R}_+^n, \Lambda) \right), \quad u_0 \in \dot{B}_{p,q,\mathcal{H}_n}^{2+s-\frac{2}{q},\sigma}(\mathbb{R}_+^n, \Lambda),$$

there exists a unique mild solution $u \in C_b^0([0, T], \dot{B}_{p,q,\mathcal{H}_n}^{2+s-\frac{2}{q},\sigma}(\mathbb{R}_+^n, \Lambda))$ of

$$(HSS_n) \quad \begin{cases} \partial_t u - \Delta u = f, & \text{on } (0, T) \times \mathbb{R}_+^n, \\ \delta u = 0, & \text{on } (0, T) \times \mathbb{R}_+^n, \\ \nu \lrcorner du|_{\partial\mathbb{R}_+^n} = 0, & \text{on } (0, T) \times \partial\mathbb{R}_+^n, \\ \nu \lrcorner u|_{\partial\mathbb{R}_+^n} = 0, & \text{on } (0, T) \times \partial\mathbb{R}_+^n, \\ u(0) = u_0, & \text{in } \dot{B}_{p,q}^{2+s-\frac{2}{q}}(\mathbb{R}_+^n, \Lambda), \end{cases}$$

with estimate

$$\|u\|_{L^\infty([0,T], \dot{B}_{p,q}^{2+s-\frac{2}{q}})} + \|(\partial_t u, \nabla^2 u)\|_{L^q((0,T), \dot{B}_{p,q}^s)} \lesssim_{p,q,n}^s \|f\|_{L^q((0,T), \dot{B}_{p,q}^s)} + \|u_0\|_{\dot{B}_{p,q}^{2+s-\frac{2}{q}}}.$$

In the case $q = +\infty$, if we assume in addition $u_0 \in \dot{A}_p^s(\mathbb{A}_{\mathcal{H},n}^2)$, we have

$$\|(\partial_t u, \nabla^2 u)\|_{L^\infty([0,T], \dot{B}_{p,\infty}^s)} \lesssim_{p,s,n} \|f\|_{L^\infty((0,T), \dot{B}_{p,\infty}^s)} + \|\mathbb{A}_{\mathcal{H},n} u_0\|_{\dot{B}_{p,\infty}^s}.$$

Remark 3.20. — Notice that above Theorem 3.19 is the only one presented here that allows L^1 and L^∞ in time maximal regularity estimates. In particular, one should notice that in the case $q = 1$, that above solution u satisfies for almost every $t \in \mathbb{R}_+$,

$$u(t), \partial_t u(t), \nabla^2 u(t) \in \dot{B}_{p,1}^s(\mathbb{R}_+^n).$$

Proof. — We may apply Theorem 3.3 to obtain maximal regularity estimates, since Proposition 3.13 gives an exact description of interpolation spaces. \square

Remark 3.21. — One may perform a cyclic permutation of systems (HHS_n) , (HMS_t) and (HSS_n) , but also exchange n and t , up to appropriate modification on boundary conditions and considered function spaces, to obtain each type of results for each operator

$$\{-\Delta_{\mathcal{H},n}, \mathbb{A}_{\mathcal{H},n}, \mathbb{M}_{\mathcal{H},n}, -\Delta_{\mathcal{H},t}, \mathbb{A}_{\mathcal{H},t}, \mathbb{M}_{\mathcal{H},t}\}.$$

3.3. Maximal regularity for the Stokes system with Navier-slip boundary conditions

The flatness of $\partial\mathbb{R}_+^n$ has the interesting consequence that, for $u : \mathbb{R}_+^n \rightarrow \mathbb{C}^n \simeq \Lambda^1$ regular enough, the normal Hodge boundary conditions

$$\begin{cases} \nu \lrcorner u|_{\partial\mathbb{R}_+^n} = 0, \\ \nu \lrcorner du|_{\partial\mathbb{R}_+^n} = 0, \end{cases}$$

are equivalent to the Navier-slip boundary conditions

$$(3.21) \quad \begin{cases} \nu \cdot u|_{\partial\mathbb{R}_+^n} = 0, \\ [({}^t \nabla u + \nabla u) \nu]_{\tan|_{\partial\mathbb{R}_+^n}} = 0. \end{cases}$$

Indeed, recalling that $\nu = -\mathbf{e}_n$, one has

$$\begin{aligned} \left[\left({}^t \nabla u + \nabla u \right) \nu \right]_{\tan} &= \left({}^t \nabla u + \nabla u \right) (-\mathbf{e}_n) - \left[\left({}^t \nabla u + \nabla u \right) (-\mathbf{e}_n) \cdot (-\mathbf{e}_n) \right] (-\mathbf{e}_n) \\ &= - \sum_{k=1}^{n-1} \left(\partial_{x_k} u_n + \partial_{x_n} u_k \right) \mathbf{e}_k. \end{aligned}$$

We may use $-\mathbf{e}_n \cdot u|_{\partial \mathbb{R}_+^n} = u_n(\cdot, 0) = 0$, yielding for all $k \in \llbracket 1, n-1 \rrbracket$

$$0 = \sum_{k=1}^{n-1} \left(\partial_{x_k} u_n(\cdot, 0) + \partial_{x_n} u_k(\cdot, 0) \right) \mathbf{e}_k = \sum_{k=1}^{n-1} \partial_{x_n} u_k(\cdot, 0) \mathbf{e}_k.$$

This implies that u satisfies exactly $n - 1$ Neumann boundary conditions, and a single Dirichlet boundary condition on u_n . This stands exactly as in Lemma 3.6. The converse also holds.

As long as one has enough regularity on u , at least in the Sobolev / Besov sense on \mathbb{R}_+^n , one is still able to perform the same decoupling for the boundary values.

This occurs when $s \in (-1 + 1/p, 1 + 1/p)$, $p \in (1, +\infty)$ for the spaces $H^{s+2,p}$, $\dot{H}^{s,p} \cap \dot{H}^{s+2,p}$, $\dot{H}^{s+2,p}$ when those are complete. It still occurs when we replace $H^{\cdot,p}$ by $B_{p,q}^{\cdot}$, $q \in [1, +\infty]$.

Therefore, in each of the previous definitions restricted to $\Lambda^1 \simeq \mathbb{C}^n$, such as e.g. (3.11), one may replace the boundary condition $\nu \lrcorner du|_{\partial \mathbb{R}_+^n} = 0$ by

$$\left[\left({}^t \nabla u + \nabla u \right) \nu \right]_{\tan|_{\partial \mathbb{R}_+^n}} = 0.$$

However, we mention the fact that, as exhibited in [MM09a, Section 2], such an identification is no longer true for (even smooth) domains Ω with non-flat boundary. In this case, the equivalence holds up to a correction term $\mathcal{W}u$, i.e. (3.21) is equivalent to

$$\begin{cases} \nu \cdot u|_{\partial \Omega} = 0, \\ \nu \lrcorner du + \mathcal{W}u|_{\partial \Omega} = 0. \end{cases}$$

Here, \mathcal{W} is the *Weingarten map*. It is linear in u and its coefficients depend linearly on the first derivatives of the outer unit normal ν , requiring then some smoothness on the boundary $\partial \Omega$. With the flat boundary $\partial \mathbb{R}_+^n$, the outer unit normal $\nu = -\mathbf{e}_n$ is constant, which explains why the map \mathcal{W} vanishes.

We can then exhibit the following maximal regularity result, where we identify Λ^1 with \mathbb{C}^n . Similar results such as Theorems 3.17 and 3.18 are also available.

THEOREM 3.22. — *Let $p \in (1, +\infty)$, $q \in [1, +\infty)$, $s \in (-1 + 1/p, 1/p + 2/q)$, such that $s, s + 2 - 2/q \notin \mathbb{N} + \frac{1}{p}$ and $(\mathcal{C}_{s+2-2/q,p,q})$ is satisfied and let $T \in (0, +\infty]$.*

For any

$$f \in L^q \left((0, T), \dot{B}_{p,q,\mathcal{H}_n}^s \left(\mathbb{R}_+^n, \mathbb{C}^n \right) \right), \quad u_0 \in \dot{B}_{p,q,\mathcal{H}_n}^{2+s-\frac{2}{q},\sigma} \left(\mathbb{R}_+^n, \mathbb{C}^n \right),$$

there exists a unique mild solution

$$(u, \nabla \mathbf{p}) \in C_b^0 \left([0, T], \dot{B}_{p,q,\mathcal{H}_n}^{2+s-\frac{2}{q},\sigma} \left(\mathbb{R}_+^n, \mathbb{C}^n \right) \right) \times L^q \left((0, T), \dot{B}_{p,q}^s \left(\mathbb{R}_+^n, \mathbb{C}^n \right) \right) \text{ of}$$

$$(NSS) \quad \begin{cases} \partial_t u - \Delta u + \nabla \mathbf{p} = f, & \text{on } (0, T) \times \mathbb{R}_+^n, \\ \operatorname{div} u = 0, & \text{on } (0, T) \times \mathbb{R}_+^n, \\ \left[\left({}^t \nabla u + \nabla u \right) \nu \right]_{\tan|_{\partial \mathbb{R}_+^n}} = 0, & \text{on } (0, T) \times \partial \mathbb{R}_+^n, \\ \nu \cdot u|_{\partial \mathbb{R}_+^n} = 0, & \text{on } (0, T) \times \partial \mathbb{R}_+^n, \\ u(0) = u_0, & \text{in } \dot{B}_{p,q}^{2+s-\frac{2}{q}}(\mathbb{R}_+^n, \mathbb{C}^n), \end{cases}$$

with estimate

$$\begin{aligned} \|u\|_{L^\infty([0,T], \dot{B}_{p,q}^{2+s-\frac{2}{q}})} + \|(\partial_t u, \nabla^2 u, \nabla \mathbf{p})\|_{L^q((0,T), \dot{B}_{p,q}^s)} \\ \lesssim_{p,q,n}^s \|f\|_{L^q((0,T), \dot{B}_{p,q}^s)} + \|u_0\|_{\dot{B}_{p,q}^{2+s-\frac{2}{q}}}. \end{aligned}$$

In the case $q = +\infty$, if we assume in addition $u_0 \in \dot{D}_p^s(\mathbb{A}_{\mathcal{H},n}^2)$, we have

$$\|(\partial_t u, \nabla^2 u, \nabla \mathbf{p})\|_{L^\infty([0,T], \dot{B}_{p,\infty}^s)} \lesssim_{p,s,n} \|f\|_{L^\infty((0,T), \dot{B}_{p,\infty}^s)} + \|\mathbb{A}_{\mathcal{H},n} u_0\|_{\dot{B}_{p,\infty}^s}.$$

Appendix A. Partial traces of differential forms

We state here a trace theorem for generalized tangential and normal traces of differential forms. The general case for vector fields in inhomogeneous function spaces is well-known, also is the case of differential forms in the setting of inhomogeneous function spaces on bounded Lipschitz domains of a Riemannian manifold, see [MMS08, Section 4] and the references therein.

We recall that $\nu = -\mathbf{e}_n$ is the outer unit normal at the boundary $\partial \mathbb{R}_+^n = \mathbb{R}^{n-1} \times \{0\}$.

THEOREM A.1. — *Let $p \in (1, +\infty)$, $q \in [1, +\infty]$, $s \in (-1 + \frac{1}{p}, \frac{1}{p})$ and let $k \in [0, n]$.*

(i) *For all $u \in D_p^s(\delta, \mathbb{R}_+^n, \Lambda^k)$. Then there exists a unique function*

$$\nu \lrcorner u|_{\partial \mathbb{R}_+^n} \in B_{p,p}^{s-\frac{1}{p}}(\mathbb{R}^{n-1}, \Lambda^{k-1})$$

*called the generalized **normal trace**, such that*

$$(A.1) \quad \begin{aligned} \int_{\mathbb{R}^{n-1}} \left\langle \nu \lrcorner u|_{\partial \mathbb{R}_+^n}(x'), \Psi|_{\partial \mathbb{R}_+^n}(x') \right\rangle dx' \\ = \int_{\mathbb{R}_+^n} \langle u(x), d\Psi(x) \rangle dx - \int_{\mathbb{R}_+^n} \langle \delta u(x), \Psi(x) \rangle dx \end{aligned}$$

for all $\Psi \in H^{1-s,p'}(\mathbb{R}_+^n, \Lambda^{k-1})$, with estimates

$$\left\| \nu \lrcorner u|_{\partial \mathbb{R}_+^n} \right\|_{B_{p,p}^{s-\frac{1}{p}}(\mathbb{R}^{n-1})} \lesssim_{p,s,n} \|u\|_{H^{s,p}(\mathbb{R}_+^n)} + \|\delta u\|_{H^{s,p}(\mathbb{R}_+^n)}.$$

The same result holds with corresponding estimate, for $u \in d_p^s(D, \mathbb{R}_+^n, \Lambda^k)$ we have a partial trace

$$\nu \wedge u|_{\partial \mathbb{R}_+^n} \in B_{p,p}^{s-\frac{1}{p}}(\mathbb{R}^{n-1}, \Lambda^{k+1})$$

called the generalized **tangential trace**, satisfying the identity

$$(A.2) \quad \int_{\mathbb{R}^{n-1}} \left\langle \nu \wedge u|_{\partial\mathbb{R}_+^n}(x'), \Psi|_{\partial\mathbb{R}_+^n}(x') \right\rangle dx' = \int_{\mathbb{R}_+^n} \langle du(x), \Psi(x) \rangle dx - \int_{\mathbb{R}_+^n} \langle u(x), \delta\Psi(x) \rangle dx.$$

for all $\Psi \in H^{1-s,p'}(\mathbb{R}_+^n, \Lambda^{k+1})$.

- (ii) For all $u \in D_{p,q}^s(\delta, \mathbb{R}_+^n, \Lambda^k)$ we have $\nu \lrcorner u|_{\partial\mathbb{R}_+^n} \in B_{p,q}^{s-\frac{1}{p}}(\mathbb{R}^{n-1}, \Lambda^{k-1})$, such that formula (A.1) holds for all $\Psi \in B_{p',q'}^{1-s}(\mathbb{R}_+^n, \Lambda)$. Moreover, we have the estimates

$$\left\| \nu \lrcorner u|_{\partial\mathbb{R}_+^n} \right\|_{B_{p,q}^{s-\frac{1}{p}}(\mathbb{R}^{n-1})} \lesssim_{p,s,n} \|u\|_{B_{p,q}^s(\mathbb{R}_+^n)} + \|\delta u\|_{B_{p,q}^s(\mathbb{R}_+^n)}.$$

The same results holds with the corresponding estimate, for $u \in d_{p,q}^s(d, \mathbb{R}_+^n, \Lambda^k)$ we have a partial trace

$$\nu \wedge u|_{\partial\mathbb{R}_+^n} \in B_{p,q}^{s-\frac{1}{p}}(\mathbb{R}^{n-1}, \Lambda^{k+1})$$

such that (A.2) is satisfied for all $\Psi \in B_{p',q'}^{1-s}(\mathbb{R}_+^n, \Lambda^{k+1})$.

- (iii) For all $u \in B_{p,q}^{s+1}(\mathbb{R}_+^n, \Lambda^k)$, we have

$$(\nu \lrcorner u \oplus \nu \wedge u)|_{\partial\mathbb{R}_+^n} \in B_{p,q}^{s+1-\frac{1}{p}}(\mathbb{R}^{n-1}, \Lambda^{k-1} \oplus \Lambda^{k+1})$$

with estimate

$$\left\| (\nu \lrcorner u \oplus \nu \wedge u)|_{\partial\mathbb{R}_+^n} \right\|_{B_{p,q}^{s+1-\frac{1}{p}}(\mathbb{R}^{n-1})} \lesssim_{p,s,n} \|u\|_{B_{p,q}^{s+1}(\mathbb{R}_+^n)},$$

and everything still hold with $H^{s+1,p}$ instead of $B_{p,q}^{s+1}$, when $q = p$.

THEOREM A.2. — Let $p \in (1, +\infty)$, $q \in [1, +\infty]$, $s \in (-1 + \frac{1}{p}, \frac{1}{p})$ and let $k \in \llbracket 0, n \rrbracket$.

- (i) For all $u \in \dot{D}_p^s(\delta, \mathbb{R}_+^n, \Lambda^k)$,

- If $s \leq 0$, then there exists a unique function $\nu \lrcorner u|_{\partial\mathbb{R}_+^n} \in B_{p,p}^{s-\frac{1}{p}}(\mathbb{R}^{n-1}, \Lambda^{k-1})$ such that the formula (A.1) holds for all $\Psi \in H^{1-s,p'}(\mathbb{R}_+^n, \Lambda)$, with estimate

$$\left\| \nu \lrcorner u|_{\partial\mathbb{R}_+^n} \right\|_{B_{p,p}^{s-\frac{1}{p}}(\mathbb{R}^{n-1})} \lesssim_{p,s,n} \|u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} + \|\delta u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)}.$$

- If $s > 0$, for $\frac{1}{r} = \frac{1}{p} - \frac{s}{n} \in (\frac{n-1}{pn}, \frac{1}{p})$, there exists a unique function

$$\nu \lrcorner u|_{\partial\mathbb{R}_+^n} \in B_{r,r}^{-\frac{1}{r}}(\mathbb{R}^{n-1}, \Lambda^{k-1}),$$

such that the formula (A.1) holds for all $\Psi \in H^{1,r'}(\mathbb{R}_+^n, \Lambda^{k-1})$ with estimate

$$\left\| \nu \lrcorner u|_{\partial\mathbb{R}_+^n} \right\|_{B_{r,r}^{-\frac{1}{r}}(\mathbb{R}^{n-1})} \lesssim_{r,p,s,n} \|u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} + \|\delta u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)}.$$

The same result, up to appropriate changes, still holds for $u \in \dot{D}_p^s(d, \mathbb{R}_+^n, \Lambda^k)$ with partial trace $\nu \wedge u|_{\partial\mathbb{R}_+^n}$ satisfying the formula (A.2).

(ii) For all $u \in \dot{D}_{p,q}^s(\delta, \mathbb{R}_+^n, \Lambda^k)$,

- If $s < 0$, there exists a unique $\nu \lrcorner u|_{\partial\mathbb{R}_+^n} \in B_{p,q}^{s-\frac{1}{p}}(\mathbb{R}^{n-1}, \Lambda^{k-1})$ such that the formula (A.1) holds for all

$$\Psi \in [\mathcal{S} \cap B_{p',q'}^{1-s}](\mathbb{R}_+^n, \Lambda^{k-1} \oplus \Lambda^{k+1}),$$

with estimates

$$\left\| \nu \lrcorner u|_{\partial\mathbb{R}_+^n} \right\|_{B_{p,q}^{s-\frac{1}{p}}(\mathbb{R}^{n-1})} \lesssim_{p,s,n} \|u\|_{\dot{B}_{p,q}^s(\mathbb{R}_+^n)} + \|\delta u\|_{\dot{B}_{p,q}^s(\mathbb{R}_+^n)}.$$

- If $s > 0$, for $\frac{1}{r} = \frac{1}{p} - \frac{s}{n} \in (\frac{n-1}{pn}, \frac{1}{p})$, there exists a unique

$$\nu \lrcorner u|_{\partial\mathbb{R}_+^n} \in B_{r,q}^{-\frac{1}{r}-\varepsilon}(\mathbb{R}^{n-1}, \Lambda^{k-1}),$$

for any sufficiently small $\varepsilon > 0$, with $\frac{1}{r} - \frac{\varepsilon}{n} = \frac{1}{r}$, such that the formula (A.1) holds for all $\Psi \in [\mathcal{S} \cap B_{r',q'}^{1+\varepsilon}](\mathbb{R}_+^n, \Lambda)$ with estimate

$$\left\| \nu \lrcorner u|_{\partial\mathbb{R}_+^n} \right\|_{B_{r,q}^{-\frac{1}{r}-\varepsilon}(\mathbb{R}^{n-1})} \lesssim_{p,s,n,\varepsilon} \|u\|_{\dot{B}_{p,q}^s(\mathbb{R}_+^n)} + \|\delta u\|_{\dot{B}_{p,q}^s(\mathbb{R}_+^n)}.$$

- If $s = 0$, there exists a unique

$$\nu \lrcorner u|_{\partial\mathbb{R}_+^n} \in B_{r,q}^{-\frac{1}{r}-\varepsilon}(\mathbb{R}^{n-1}, \Lambda^{k-1}),$$

where $\frac{1}{r} = \frac{1}{p} - \frac{\varepsilon}{n}$, for any sufficiently small $\varepsilon > 0$, such that the formula (A.1) holds for all $\Psi \in [\mathcal{S} \cap B_{r',q'}^{1+\varepsilon}](\mathbb{R}_+^n, \Lambda)$, with estimates

$$\left\| \nu \lrcorner u|_{\partial\mathbb{R}_+^n} \right\|_{B_{r,q}^{-\frac{1}{r}-\varepsilon}(\mathbb{R}^{n-1})} \lesssim_{p,s,n,\varepsilon} \|u\|_{\dot{B}_{p,q}^0(\mathbb{R}_+^n)} + \|\delta u\|_{\dot{B}_{p,q}^0(\mathbb{R}_+^n)}.$$

The same result, up to appropriate changes, still holds for $u \in \dot{d}_{p,q}^s(d, \mathbb{R}_+^n, \Lambda^k)$ with partial trace $\nu \wedge u|_{\partial\mathbb{R}_+^n}$ satisfying the formula (A.2).

(iii) For all $u \in [\dot{B}_{p,q}^s \cap \dot{B}_{p,q}^{s+1}](\mathbb{R}_+^n, \Lambda^k)$, if $q \neq +\infty$, we have

$$(\nu \lrcorner u \oplus \nu \wedge u)|_{\partial\mathbb{R}_+^n} \in B_{p,q}^{s+1-\frac{1}{p}}(\mathbb{R}^{n-1}, \Lambda^{k-1} \oplus \Lambda^{k+1})$$

with estimate

$$\left\| (\nu \lrcorner u \oplus \nu \wedge u)|_{\partial\mathbb{R}_+^n} \right\|_{B_{p,q}^{s+1-\frac{1}{p}}(\mathbb{R}^{n-1})} \lesssim_{p,s,n} \|u\|_{[\dot{B}_{p,q}^s \cap \dot{B}_{p,q}^{s+1}](\mathbb{R}_+^n)},$$

and everything still hold with $(\dot{H}^{s,p}, \dot{H}^{s+1,p})$ instead of $(\dot{B}_{p,q}^s, \dot{B}_{p,q}^{s+1})$, when $q = p$.

If $q = +\infty$, we have

$$(\nu \lrcorner u \oplus \nu \wedge u)|_{\partial\mathbb{R}_+^n} \in L^p(\mathbb{R}^{n-1}, \Lambda^{k-1} \oplus \Lambda^{k+1})$$

with corresponding estimate.

Remark A.3. — The proof of Theorem A.1 in case of inhomogeneous function spaces follows straightforward the same proof provided for corresponding results in [MMS08, Section 4] and is somewhat sharp.

Notice that Theorem A.2 is certainly not sharp, and investigation of sharp range for partial traces could be of great interest in the treatment of inhomogeneous boundary value problems in homogeneous function spaces.

Proof. — Without loss of generality, we only investigate the case of normal traces $\nu \lrcorner u|_{\partial\mathbb{R}_+^n}$.

Step 1.1: Proof of (i), for $s \leq 0$. Let $u \in \dot{D}_p^s(\delta, \mathbb{R}_+^n, \Lambda^k)$, We can define for all

$$\psi \in B_{p',p'}^{\frac{1}{p}-s}(\mathbb{R}^{n-1}, \Lambda^{k-1}),$$

and $\Psi \in H^{1-s,p'}(\mathbb{R}_+^n, \Lambda^{k-1})$ such that $\Psi|_{\partial\mathbb{R}_+^n} = \psi$, the following functional,

$$\kappa_u(\Psi) := \int_{\mathbb{R}_+^n} \langle u(x), d\Psi(x) \rangle dx - \int_{\mathbb{R}_+^n} \langle \delta u(x), \Psi(x) \rangle dx.$$

First, the map $(u, \Psi) \mapsto \kappa_u(\Psi)$ is well-defined and bilinear on $\dot{D}_p^s(\delta, \mathbb{R}_+^n, \Lambda^k) \times H^{1-s,p'}(\mathbb{R}_+^n, \Lambda^{k-1})$, i.e., in particular only depends on the boundary value ψ of Ψ . It is straightforward from duality that,

$$\begin{aligned} \|\kappa_u(\Psi)\| &\lesssim_{s,p,n} \|u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} \|d\Psi\|_{\dot{H}^{-s,p'}(\mathbb{R}_+^n)} + \|\delta u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} \|\Psi\|_{\dot{H}^{-s,p'}(\mathbb{R}_+^n)} \\ &\lesssim_{s,p,n} \|u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} \|\Psi\|_{\dot{H}^{1-s,p'}(\mathbb{R}_+^n)} + \|\delta u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} \|\Psi\|_{\dot{H}^{-s,p'}(\mathbb{R}_+^n)} \\ &\lesssim_{s,p,n} \|u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} \|\Psi\|_{H^{1-s,p'}(\mathbb{R}_+^n)} + \|\delta u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} \|\Psi\|_{H^{-s,p'}(\mathbb{R}_+^n)} \\ &\lesssim_{s,p,n} \left(\|u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} + \|\delta u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} \right) \|\Psi\|_{H^{1-s,p'}(\mathbb{R}_+^n)}, \end{aligned}$$

where above inequalities follows from

$$H^{-s,p'}(\mathbb{R}_+^n) \hookrightarrow \dot{H}^{-s,p'}(\mathbb{R}_+^n), H^{1-s,p'}(\mathbb{R}_+^n) \hookrightarrow \dot{H}^{1-s,p'}(\mathbb{R}_+^n),$$

since $-1 + \frac{1}{p} < s \leq 0$, then from $H^{1-s,p'}(\mathbb{R}_+^n) \hookrightarrow H^{-s,p'}(\mathbb{R}_+^n)$.

Now, if we have $\Psi_1, \Psi_2 \in H^{1-s,p'}(\mathbb{R}_+^n, \Lambda^{k-1})$ such that $\Psi_1|_{\partial\mathbb{R}_+^n} = \Psi_2|_{\partial\mathbb{R}_+^n} = \psi$, we introduce

$$\Psi_0 = \Psi_1 - \Psi_2 \in H_0^{1-s,p'}(\mathbb{R}_+^n, \Lambda^{k-1}).$$

Therefore, let's consider $(\Phi_k)_{k \in \mathbb{N}} \subset C_c^\infty(\mathbb{R}_+^n, \Lambda^{k-1})$ such that,

$$\Phi_k \xrightarrow[k \rightarrow +\infty]{} \Psi_0 \text{ in } H_0^{1-s,p'}(\mathbb{R}_+^n, \Lambda^{k-1}).$$

We can deduce,

$$\begin{aligned} \kappa_u(\Psi_1) - \kappa_u(\Psi_2) &= \kappa_u(\Psi_0) = \lim_{k \rightarrow +\infty} \left[\int_{\mathbb{R}_+^n} \langle u(x), d\Phi_k(x) \rangle dx - \int_{\mathbb{R}_+^n} \langle \delta u(x), \Phi_k(x) \rangle dx \right] \\ &= 0. \end{aligned}$$

Thus, the following equality occurs, where we also consider the extension operator $\text{Ext}_{\mathbb{R}_+^n}$ from [BL76, Exercises 25, 26, p. 166], see also [Gau23, Theorem 2.45],

$$\tilde{\kappa}_u(\psi) := \kappa_u \left(\text{Ext}_{\mathbb{R}_+^n} \otimes \psi \right) = \kappa_u(\Psi),$$

and with estimate, also obtained from [Gau23, Theorem 2.45],

$$\|\tilde{\kappa}_u(\psi)\| \lesssim_{s,p,n} \left(\|u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} + \|du\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} \right) \|\psi\|_{B_{p',p'}^{\frac{1}{p}-s}(\mathbb{R}^{n-1})}.$$

By duality, there exists a unique function depending linearly on

$$u, \nu \lrcorner u|_{\partial\mathbb{R}_+^n} \in B_{p,p}^{s-\frac{1}{p}}(\mathbb{R}^{n-1}, \Lambda^{k-1})$$

such that (A.1) holds.

To guarantee that the representation formula makes sense, one may use the usual integration by parts formula with $u, \Psi \in \mathcal{S}(\overline{\mathbb{R}_+^n}, \Lambda)$.

Step 1.2: Proof of (i), for $s > 0$. For the same assumption on u , and Ψ as before, everything works similarly except the way we bounded bilinearly the map $(u, \Psi) \mapsto \kappa_u(\Psi)$ on $\dot{D}_p^s(\delta, \mathbb{R}_+^n, \Lambda^k) \times H^{1-s,p'}(\mathbb{R}_+^n, \Lambda^{k-1})$. For $r \in (1, +\infty)$ such that $\frac{1}{r} = \frac{1}{p} - \frac{s}{n}$, we deduce from Sobolev embeddings and duality that

$$\begin{aligned} \|\kappa_u(\Psi)\| &\lesssim_{s,p,n} \|u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} \|d\Psi\|_{\dot{H}^{-s,p'}(\mathbb{R}_+^n)} + \|\delta u\|_{L^r(\mathbb{R}_+^n)} \|\Psi\|_{L^{r'}(\mathbb{R}_+^n)} \\ &\lesssim_{r,s,p,n} \|u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} \|\Psi\|_{H^{1,r'}(\mathbb{R}_+^n)} + \|\delta u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} \|\Psi\|_{H^{1,r'}(\mathbb{R}_+^n)} \\ &\lesssim_{r,s,p,n} \left(\|u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} + \|\delta u\|_{\dot{H}^{s,p}(\mathbb{R}_+^n)} \right) \|\Psi\|_{H^{1,r'}(\mathbb{R}_+^n)}. \end{aligned}$$

Thus everything goes similarly.

Step 2.1: Proof of (ii) for $s < 0$, is very similar to the one of above Step 1.1.

Step 2.2: Proof of 2.(ii), for $s > 0$, is somewhat similar to the one of Step 1.2 but needs further explanations. We use Sobolev embeddings, and generalized Hölder inequalities using Lorentz spaces,

$$\dot{B}_{p,q}^s(\mathbb{R}_+^n) \hookrightarrow L^{r,q}(\mathbb{R}_+^n), B_{\tilde{r},q}^\varepsilon(\mathbb{R}_+^n) \hookrightarrow L^{r',q'}(\mathbb{R}_+^n) \hookrightarrow \dot{B}_{p',q'}^{-s}(\mathbb{R}_+^n), \text{ for } \varepsilon > 0,$$

$$\begin{aligned} \|\kappa_u(\Psi)\| &\lesssim_{s,p,n} \|u\|_{\dot{B}_{p,q}^s(\mathbb{R}_+^n)} \|d\Psi\|_{\dot{B}_{p',q'}^{-s}(\mathbb{R}_+^n)} + \|\delta u\|_{L^{r,q}(\mathbb{R}_+^n)} \|\Psi\|_{L^{r',q'}(\mathbb{R}_+^n)} \\ &\lesssim_{r,s,p,n} \|u\|_{\dot{B}_{p,q}^s(\mathbb{R}_+^n)} \|d\Psi\|_{L^{r',q'}(\mathbb{R}_+^n)} + \|\delta u\|_{\dot{B}_{p,q}^s(\mathbb{R}_+^n)} \|\Psi\|_{L^{r',q'}(\mathbb{R}_+^n)} \\ &\lesssim_{r,s,p,n} \left(\|u\|_{\dot{B}_{p,q}^s(\mathbb{R}_+^n)} + \|\delta u\|_{\dot{B}_{p,q}^s(\mathbb{R}_+^n)} \right) \|\Psi\|_{B_{\tilde{r},q'}^{1+\varepsilon}(\mathbb{R}_+^n)}. \end{aligned}$$

Step 2.3: Proof of (ii), for $s = 0$ is shown via similar Sobolev embeddings arguments and is left to the reader.

Step 3: Proof of (iii), follows from [Gau24b, Proposition 4.4], with explicit representation formula for any suitable k -differential forms u :

$$\begin{aligned} \nu \lrcorner u|_{\partial\mathbb{R}_+^n} &= -\mathbf{e}_n \lrcorner u|_{\partial\mathbb{R}_+^n} = (-1)^k \sum_{1 \leq \ell_1 < \dots < \ell_{k-1} < n} u_{\ell_1 \ell_2 \dots \ell_{k-1} n}(\cdot, 0) dx_{\ell_1} \wedge \dots \wedge dx_{\ell_{k-1}} \\ &= (-1)^k \sum_{I' \in \mathcal{I}_{n-1}^{k-1}} u_{I', n}(\cdot, 0) dx_{I'}. \end{aligned}$$

A similar treatment yields the same conclusion for the boundary term $\nu \wedge u|_{\partial\mathbb{R}_+^n}$, so that one ends the proof here. \square

Remark A.4. — Let's make further comment about estimates used in the proof of Theorem A.2 above, in particular the ones used in Step 2.2.

We recall that from Sobolev embeddings, see [BCD11, Proposition 2.39], for $0 < s_0 < s < s_1 < 1/p$, $r_0, r_1, p \in (1, +\infty)$, $1/r_j = 1/p - s_j/n$, we have by the definition of function spaces by restriction

$$\dot{B}_{p,1}^{s_j}(\mathbb{R}_+^n) \hookrightarrow L^{r_j}(\mathbb{R}_+^n), \quad j \in \{0, 1\}.$$

If $(s, 1/r) = (1 - \theta)(s_0, 1/r_0) + \theta(s_1, 1/r_1)$, by real interpolation, for $q \in [1, +\infty]$ we obtain,

$$\dot{B}_{p,q}^s(\mathbb{R}_+^n) = \left(\dot{B}_{p,1}^{s_0}(\mathbb{R}_+^n), \dot{B}_{p,1}^{s_1}(\mathbb{R}_+^n) \right)_{\theta,q} \hookrightarrow \left(L^{r_0}(\mathbb{R}_+^n), L^{r_1}(\mathbb{R}_+^n) \right)_{\theta,q} = L^{r,q}(\mathbb{R}_+^n).$$

And one may proceed similarly for the reverse embedding,

$$L^{r',q'}(\mathbb{R}_+^n) \hookrightarrow \dot{B}_{p',q'}^{-s}(\mathbb{R}_+^n).$$

For more details about Lorentz spaces and their interpolation, one could consult [Lun18, Section 1, Examples 1.10, 1.11 & 1.27] and [BL76, Chapter 5, Section 5.3].

BIBLIOGRAPHY

- [ABHN11] W. Arendt, C. J. K. Batty, M. Hieber, and F. Neubrander, *Vector-valued Laplace transforms and Cauchy problems*, 2nd ed., Monographs in Mathematics, vol. 96, Birkhäuser, 2011. ↑1469, 1484, 1505
- [ACDH04] P. Auscher, T. Coulhon, X. T. Duong, and S. Hofmann, *Riesz transform on manifolds and heat kernel regularity*, Ann. Sci. Éc. Norm. Supér. **37** (2004), no. 6, 911–957. ↑1461
- [AM04] A. Axelsson and A. McIntosh, *Hodge decompositions on weakly Lipschitz domains*, Advances in analysis and geometry. New developments using Clifford algebras, Trends in Mathematics, Birkhäuser, 2004, pp. 3–29. ↑1477
- [Ama95] H. Amann, *Linear and quasilinear parabolic problems, Vol. I: Abstract linear theory*, Monographs in Mathematics, vol. 89, Birkhäuser, 1995. ↑1463, 1464
- [BCD11] H. Bahouri, J.-Y. Chemin, and R. Danchin, *Fourier analysis and nonlinear partial differential equations*, Grundlehren der Mathematischen Wissenschaften, vol. 343, Springer, 2011. ↑1465, 1471, 1472, 1482, 1483, 1484, 1530
- [BL76] J. Bergh and J. Löfström, *Interpolation spaces. An introduction*, Grundlehren der Mathematischen Wissenschaften, vol. 223, Springer, 1976. ↑1465, 1470, 1472, 1473, 1514, 1515, 1516, 1517, 1518, 1519, 1520, 1529, 1530

- [Bog86] M. E. Bogovskii, *Decomposition of $L_p(\Omega, \mathbf{R}^n)$ into the direct sum of subspaces of solenoidal and potential vector fields*, Sov. Math., Dokl. **33** (1986), no. 1, 161–165. ↑1460
- [Che99] J.-Y. Chemin, *Théorèmes d'unicité pour le système de Navier–Stokes tridimensionnel*, J. Anal. Math. **77** (1999), no. 1, 27–50. ↑1464
- [CM10] M. Costabel and A. McIntosh, *On Bogovskii and regularized Poincaré integral operators for de Rham complexes on Lipschitz domains*, Math. Z. **265** (2010), 297–320. ↑1477
- [DC94] M. Do Carmo, *Differential Forms and Applications*, Universitext, Springer, 1994, original Portuguese edition published by IMPA, 1971. ↑1477
- [Den22] C. Denis, *Existence and uniqueness in critical spaces for the magnetohydrodynamical system in \mathbb{R}^n* , preprint, HAL-03660738v2, 2022. ↑1462
- [DHMT21] R. Danchin, M. Hieber, P. B. Mucha, and P. Tolksdorf, *Free Boundary Problems via Da Prato–Grisvard Theory*, 2021, <https://arxiv.org/abs/2011.07918v2>. ↑1464, 1465, 1466, 1471, 1472, 1483, 1505, 1506, 1513, 1514, 1519, 1520, 1521
- [DHP03] R. Denk, M. Hieber, and J. Prüss, *\mathcal{R} -boundedness, Fourier multipliers, and problems of elliptic and parabolic type*, Memoirs of the American Mathematical Society, vol. 788, American Mathematical Society, 2003. ↑1463, 1464
- [DK07] R. Denk and T. Krainer, *\mathcal{R} -boundedness, pseudodifferential operators, and maximal regularity for some classes of partial differential operators*, Manuscr. Math. **124** (2007), no. 3, 319–342. ↑1464
- [DM09] R. Danchin and P. B. Mucha, *A critical functional framework for the inhomogeneous Navier–Stokes equations in the half-space*, J. Funct. Anal. **256** (2009), no. 3, 881–927. ↑1464
- [DM15] ———, *Critical functional framework and maximal regularity in action on systems of incompressible flows*, Mémoires de la Société Mathématique de France. Nouvelle Série, vol. 143, Société Mathématique de France, 2015. ↑1464, 1483
- [DPG75] G. Da Prato and P. Grisvard, *Sommes d'opérateurs linéaires et équations différentielles opérationnelles*, J. Math. Pures Appl. **54** (1975), no. 3, 305–387. ↑1464, 1465
- [DW05] M. Duelli and L. Weis, *Spectral projections, Riesz transforms and H^∞ -calculus for bisectorial operator*, Nonlinear elliptic and parabolic problems. A special tribute to the work of Herbert Amann, Zürich, Switzerland, June 28–30, 2004, Progress in Nonlinear Differential Equations and their Applications, vol. 64, Birkhäuser, 2005, pp. 99–111. ↑1470
- [Ege15] M. Egert, *On Kato's conjecture and mixed boundary conditions*, Ph.D. thesis, Technischen Universität Darmstadt, 2015. ↑1469, 1470, 1491
- [FKS05] R. Farwig, H. Kozono, and H. Sohr, *An L^q approach to Stokes and Navier–Stokes in general domains*, Acta Math. **195** (2005), 21–53. ↑1460
- [FKS07] ———, *On the Helmholtz decomposition in general unbounded domains*, Arch. Math. **88** (2007), 239–248. ↑1460
- [FMM98] E. Fabes, O. Mendez, and M. Mitrea, *Boundary layers on Sobolev–Besov spaces and Poisson's equation for the Laplacian in Lipschitz domains*, J. Funct. Anal. **159** (1998), no. 2, 323–368. ↑1460
- [FY06] H. Fujiwara and M. Yamazaki, *The Helmholtz decomposition in Sobolev and Besov spaces*, Asymptotic analysis and singularities. Hyperbolic and dispersive PDEs and fluid mechanics, Advanced Studies in Pure Mathematics, vol. 47, Mathematical Society of Japan, 2006, pp. 99–116. ↑1460
- [Gal11] G. P. Galdi, *An Introduction to the Mathematical Theory of the Navier–Stokes Equations*, 2nd ed., Springer Monographs in Mathematics, Springer, 2011. ↑1460

- [Gau23] A. Gaudin, *Homogeneous Sobolev and Besov spaces on half-spaces*, Ph.D. thesis, Aix-Marseille Université, Marseille, France, 2023, <https://hal.science/tel-04169055>. ↑1529
- [Gau24a] ———, *Homogeneous Sobolev global-in-time maximal regularity and related trace estimates*, *J. Evol. Equ.* **24** (2024), no. 1, article no. 15. ↑1464, 1466, 1505, 1507
- [Gau24b] ———, *On homogeneous Sobolev and Besov spaces on the whole and the half space*, *Tunis. J. Math.* **6** (2024), no. 2, 343–404. ↑1465, 1466, 1472, 1473, 1474, 1475, 1476, 1482, 1483, 1490, 1491, 1492, 1508, 1509, 1511, 1512, 1513, 1514, 1515, 1516, 1517, 1518, 1530
- [GHT13] M. Geissert, H. Heck, and C. Trunk, *\mathcal{H}^∞ -calculus for a system of Laplace operators with mixed order boundary conditions*, *Discrete Contin. Dyn. Syst., Ser. S* **6** (2013), no. 5, 1259–1275. ↑1460, 1463
- [GS91] Y. Giga and H. Sohr, *Abstract L^p estimates for the Cauchy problem with applications to the Navier–Stokes equations in exterior domains*, *J. Funct. Anal.* **102** (1991), no. 1, 72–94. ↑1464
- [Gui91a] D. Guidetti, *On elliptic problems in Besov spaces*, *Math. Nachr.* **152** (1991), 247–275. ↑1509
- [Gui91b] ———, *On interpolation with boundary conditions*, *Math. Z.* **207** (1991), no. 3, 439–460. ↑1509, 1511
- [Haa06] M. Haase, *The functional calculus for sectorial operators*, *Operator Theory: Advances and Applications*, vol. 169, Birkhäuser, 2006. ↑1465, 1468, 1469, 1470, 1485, 1489, 1494, 1505
- [HHK06] B. H. Haak, M. Haase, and P. C. Kunstmann, *Perturbation, interpolation, and maximal regularity*, *Adv. Differ. Equ.* **11** (2006), no. 2, 201–240. ↑1465
- [Hie20] M. Hieber, *Analysis of Viscous Fluid Flows: An Approach by Evolution Equations*, *Mathematical Analysis of the Navier–Stokes Equations: Cetraro, Italy 2017* (G. P. Galdi and Y. Shibata, eds.), *Lecture Notes in Mathematics*, vol. 2254, Springer, 2020, pp. 1–146. ↑1464
- [HM13] M. Hieber and S. Monniaux, *Well-posedness results for the Navier–Stokes equations in the rotational framework*, *Discrete Contin. Dyn. Syst.* **33** (2013), no. 11–12, 5143–5151. ↑1464
- [HMM11] S. Hofmann, M. Mitrea, and S. Monniaux, *Riesz transforms associated with the Hodge Laplacian in Lipschitz subdomains of Riemannian manifolds*, *Ann. Inst. Fourier* **61** (2011), no. 4, 1323–1349. ↑1462
- [HNPS16] M. Hieber, M. Nesensohn, J. Prüss, and K. Schade, *Dynamics of nematic liquid crystal flows: The quasilinear approach*, *Ann. Inst. Henri Poincaré, Anal. Non Linéaire* **33** (2016), no. 2, 397–408. ↑1464
- [Jos11] J. Jost, *Riemannian Geometry and Geometric Analysis*, 6th ed., Universitext, Springer, 2011. ↑1477
- [KW04] P. C. Kunstmann and L. Weis, *Maximal L^p -regularity for Parabolic Equations, Fourier Multiplier Theorems and H^∞ -functional Calculus*, *Functional Analytic Methods for Evolution Equations*, *Lecture Notes in Mathematics*, vol. 1855, Springer, 2004, pp. 65–311. ↑1463, 1464
- [Lun18] A. Lunardi, *Interpolation theory*, 3rd ed., Appunti. Scuola Normale Superiore di Pisa (Nuova Serie), Edizioni della Normale, 2018. ↑1470, 1530
- [McI86] A. McIntosh, *Operators which have an H_∞ functional calculus*, *Miniconference on operator theory and partial differential equations* (North Ryde, 1986), *Proceedings of the Centre for Mathematical Analysis, Australian National University*, vol. 14, Australian National University, Canberra, 1986, pp. 210–231. ↑1489, 1494

- [MM08] M. Mitrea and S. Monniaux, *The regularity of the Stokes operator and the Fujita–Kato approach to the Navier–Stokes initial value problem in Lipschitz domains*, J. Funct. Anal. **254** (2008), no. 6, 1522–1574. ↑1460
- [MM09a] ———, *The nonlinear Hodge–Navier–Stokes equations in Lipschitz domains*, Differ. Integral Equ. **22** (2009), no. 3-4, 339–356. ↑1464, 1524
- [MM09b] ———, *On the analyticity of the semigroup generated by the Stokes operator with Neumann-type boundary conditions on Lipschitz subdomains of Riemannian manifolds*, Trans. Am. Math. Soc. **361** (2009), no. 6, 3125–3157. ↑1462
- [MM18] A. McIntosh and S. Monniaux, *Hodge–Dirac, Hodge–Laplacian and Hodge–Stokes operators in L^p spaces on Lipschitz domains*, Rev. Mat. Iberoam. **34** (2018), no. 4, 1711–1753. ↑1461, 1476, 1489
- [MMS08] D. Mitrea, M. Mitrea, and M.-C. Shaw, *Traces of Differential Forms on Lipschitz Domains, the Boundary De Rham Complex, and Hodge Decompositions*, Indiana Univ. Math. J. **57** (2008), no. 5, 2061–2095. ↑1525, 1528
- [Mon13] S. Monniaux, *Various boundary conditions for Navier–Stokes equations in bounded Lipschitz domains*, Discrete Contin. Dyn. Syst., Ser. S **6** (2013), no. 5, 1355–1369. ↑1464
- [Mon21] ———, *Existence in critical spaces for the magnetohydrodynamical system in 3D bounded Lipschitz domains*, J. Elliptic Parabol. Equ. **7** (2021), 311–322. ↑1462, 1464, 1476
- [MS18] S. Monniaux and Z. Shen, *Stokes problems in irregular domains with various boundary conditions*, Handbook of mathematical analysis in mechanics of viscous fluids (Y. Giga and A. Novotný, eds.), Springer, 2018, pp. 207–248. ↑1460
- [OS16] T. Ogawa and S. Shimizu, *End-point maximal L^1 -regularity for the Cauchy problem to a parabolic equation with variable coefficients*, Math. Ann. **365** (2016), no. 1-2, 661–705. ↑1464
- [OS21] ———, *Global well-posedness for the incompressible Navier–Stokes equations in the critical Besov space under the Lagrangian coordinates*, J. Differ. Equations **274** (2021), 613–651. ↑1464
- [OS22] ———, *Maximal L^1 -regularity for parabolic initial-boundary value problems with inhomogeneous data*, J. Evol. Equ. **22** (2022), no. 2, article no. 30. ↑1464
- [Ouh05] E. M. Ouhabaz, *Analysis of heat equations on domains*, London Mathematical Society Monographs, vol. 31, Princeton University Press, 2005. ↑1489
- [Sch95] G. Schwarz, *Hodge Decomposition A Method for Solving Boundary Value Problems*, Lecture Notes in Mathematics, Springer, 1995. ↑1461
- [Soh01] H. Sohr, *The Navier–Stokes Equations: An Elementary Functional Analytic Approach*, Birkhäuser Advanced Texts. Basler Lehrbücher, Birkhäuser, 2001. ↑1459, 1480
- [SS92] C. G. Simader and H. Sohr, *A new approach to the Helmholtz decomposition and the Neumann problem in L^q -spaces for bounded and exterior domains*, Mathematical Problems Relating to the Navier–Stokes Equations (G. P. Galdi, ed.), Series on Advances in Mathematics for Applied Sciences, vol. 11, World Scientific, 1992, pp. 1–35. ↑1460
- [Ste70] E. M. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton Mathematical Series, vol. 30, Princeton University Press, 1970. ↑1485
- [Tol17] P. Tolksdorf, *On the L^p -theory of the Navier–Stokes equation on Lipschitz domains*, Ph.D. thesis, Technische Universität Darmstadt, 2017. ↑1460
- [Tol18] ———, *On the L^p -theory of the Navier–Stokes equations on three-dimensional bounded Lipschitz domains*, Math. Ann. **371** (2018), no. 1-2, 445–460. ↑1464
- [Tri78] H. Triebel, *Interpolation Theory, Function Spaces, Differential Operators*, North-Holland Mathematical Library, vol. 18, Elsevier, 1978. ↑1470

- [Tri83] ———, *Theory of Function Spaces. I*, Monographs in Mathematics, vol. 78, Birkhäuser, 1983. ↑1465
- [TW20] P. Tolksdorf and K. Watanabe, *The Navier–Stokes equations in exterior Lipschitz domains: L^p -theory*, J. Differ. Equations **269** (2020), no. 7, 5765–5801. ↑1464
- [Wei01] L. Weis, *Operator–Valued Multiplier Theorems and Maximal L_p –Regularity*, Math. Ann. **319** (2001), no. 4, 735–758. ↑1464

Manuscript received on 8th March 2023,
revised on 3rd May 2024,
accepted on 22nd August 2024.

Recommended by Editors S. Vu Ngoc and N. Raymond.
Published under license CC BY 4.0.



eISSN: 2644-9463

This journal is a member of Centre Mersenne.



Anatole GAUDIN
Technische Universität Clausthal,
Institut für Mathematik,
Erzstraße 1,
38678 Clausthal-Zellerfeld (Germany)
anatole.gaudin@tu-clausthal.de