Annales Henri Lebesgue **8** (2025) 1-34



ZAKHAR KABLUCHKO

LEE-YANG ZEROES OF THE CURIE-WEISS FERROMAGNET, UNITARY HERMITE POLYNOMIALS, AND THE BACKWARD HEAT FLOW ZÉROS DE LEE-YANG POUR LE MODÈL

ZÉROS DE LEE-YANG POUR LE MODÈLE DE CURIE-WEISS FERROMAGNÉTIQUE, POLYNÔMES D'HERMITE UNITAIRES ET FLOT DE LA CHALEUR RÉTROGRADE

ABSTRACT. — The backward heat flow on the real line started from the initial condition z^n results in the classical n^{th} Hermite polynomial whose zeroes are distributed according to the Wigner semicircle law in the large n limit. Similarly, the backward heat flow with the periodic initial condition $(\sin \frac{\theta}{2})^n$ leads to trigonometric or unitary analogues of the Hermite polynomials. These polynomials are closely related to the partition function of the Curie–Weiss model and appeared in the work of Mirabelli on finite free probability. We relate the n^{th} unitary Hermite polynomial to the expected characteristic polynomial of a unitary random matrix obtained by running a Brownian motion on the unitary group U(n). We identify the

Keywords: Curie–Weiss model, unitary Hermite polynomials, Lee–Yang zeroes, finite free probability, free multiplicative convolution, free unitary normal distribution, saddle-point method. 2020 Mathematics Subject Classification: 82B20, 30C15, 30C10, 60B10, 46L54, 30F99, 30E15. DOI: https://doi.org/10.5802/ahl.227

^(*) The author is supported by the German Research Foundation under Germany's Excellence Strategy EXC 2044 – 390685587, *Mathematics Münster: Dynamics – Geometry – Structure*.

global distribution of zeroes of the unitary Hermite polynomials as the free unitary normal distribution. We also compute the asymptotics of these polynomials or, equivalently, the free energy of the Curie–Weiss model in a complex external field. We identify the global distribution of the Lee–Yang zeroes of this model. Finally, we show that the backward heat flow applied to a high-degree real-rooted polynomial (respectively, trigonometric polynomial) induces, on the level of the asymptotic distribution of its roots, a free Brownian motion (respectively, free unitary Brownian motion).

RÉSUMÉ. — Le flot de la chaleur rétrograde sur la droite réelle, partant de la condition initiale z^n , donne le $n^{\text{ème}}$ polynôme d'Hermite classique, dont les zéros sont distribués selon la loi du demi-cercle de Wigner pour n grand. De la même manière, le flot de la chaleur rétrograde avec la condition initiale périodique $(\sin \frac{\theta}{2})^n$ conduit à des analogues trigonométriques ou unitaires des polynômes d'Hermite. Ces polynômes sont étroitement liés à la fonction de partition du modèle de Curie-Weiss et sont apparus dans les travaux de Mirabelli sur les probabilités libres finies. Nous relions le $n^{\text{ème}}$ polynôme d'Hermite unitaire à l'espérance du polynôme caractéristique d'une matrice aléatoire unitaire obtenue en exécutant un mouvement brownien sur le groupe unitaire U(n). Nous identifions la distribution globale des zéros des polynômes d'Hermite unitaires comme la distribution normale unitaire libre. Nous calculons également l'asymptotique de ces polynômes ou, de manière équivalente, l'énergie libre du modèle de Curie-Weiss dans un champ extérieur complexe. Nous identifions la distribution globale des zéros de Lee-Yang de ce modèle. Enfin, nous montrons que le flot de la chaleur rétrograde appliqué à un polynôme (respectivement, polynôme trigonométrique) de haut degré à racines réelles induit, au niveau de la distribution asymptotique de ses racines, un mouvement brownien libre (respectivement, mouvement brownien unitaire libre).

1. Introduction

1.1. Hermite polynomials and their trigonometric analogues

One possible way to define the classical (probabilist) Hermite polynomials $\text{He}_0(z) = 1$, $\text{He}_1(z) = z$, $\text{He}_2(z) = z^2 - 1$,... is the formula

(1.1)
$$\operatorname{He}_{n}(z) = \exp\left\{-\frac{1}{2}\partial_{z}^{2}\right\} z^{n} = n! \sum_{m=0}^{[n/2]} \frac{(-1)^{m}}{m!2^{m}} \cdot \frac{z^{n-2m}}{(n-2m)!}, \quad n \in \mathbb{N}_{0}.$$

where ∂_z denotes differentiation in z and the exponential can be understood as an infinite series which terminates after finitely many non-zero summands. More generally, we have

(1.2)
$$\exp\left\{-\frac{1}{2}\sigma^2\partial_z^2\right\}z^n = \sigma^n\operatorname{He}_n\left(\frac{z}{\sigma}\right), \quad \sigma > 0.$$

This can be expressed by saying that the n^{th} Hermite polynomial arises when solving the *backward* heat equation $\partial_t f(z;t) = -\frac{1}{2} \partial_z^2 f(z;t)$ on the real line with the initial condition $f(z;0) = z^n$.

We shall be interested in the trigonometric (or unitary) analogues of the Hermite polynomials. To introduce them, it will be convenient to adopt the following (somewhat unconventional) terminology. A trigonometric polynomial $T_n(\theta)$ of degree $n \in \mathbb{N}_0$ is an expression of the form

(1.3)
$$T_n(\theta) = \frac{P_n(e^{i\theta})}{e^{in\theta/2}},$$

where $P_n(z) \in \mathbb{C}[z]$ is an algebraic polynomial of degree *n* such that $P_n(0) \neq 0$. For example, if $P_n(z) = (z-1)^n$, then $T_n(\theta) = (2 \operatorname{i} \sin \frac{\theta}{2})^n$.

In general, $T_n(\theta)$ is a linear combination of the functions $\theta \mapsto e^{i(k-\frac{n}{2})\theta}$, $k = 0, \ldots, n$, with complex coefficients (such that the first and the last coefficient do not vanish). If n = 2d is even, then $T_n(\theta)$ can be represented as a linear combination of the functions $1, \sin \theta, \cos \theta, \ldots, \sin(d\theta), \cos(d\theta)$. This case corresponds to the usual definition of trigonometric polynomials. If n = 2d + 1 is odd, then $T_n(\theta)$ can be written as a linear combination of the functions $\sin(\ell\theta/2)$ and $\cos(\ell\theta/2)$ with $\ell = 1, 3, 5, \ldots, n$. This case is somewhat unconventional. From the representation $P_n(z) = C \prod_{j=1}^n (z - z_j)$, where $z_1, \ldots, z_n \in \mathbb{C} \setminus \{0\}$ are the complex zeroes of P_n , one deduces the existence of a representation

$$T_n(\theta) = C' \prod_{j=1}^n \sin \frac{\theta - \theta_j}{2}$$

where $\theta_1, \ldots, \theta_n \in \mathbb{C}$ are chosen to satisfy $e^{i\theta_j} = z_j \neq 0$.

We shall be interested in trigonometric polynomials with real roots only or, equivalently, in algebraic polynomials having all roots on the unit circle. If we consider $(z-1)^n$ as the "simplest" algebraic polynomial of degree n with this property, then the "simplest" real-rooted trigonometric polynomial of degree n is $(\sin \frac{\theta}{2})^n$ (up to a constant factor). To derive the trigonometric analogues of the Hermite polynomials He_n , we look at the backward heat flow on \mathbb{R} with the initial condition $(\sin \frac{\theta}{2})^n$. More precisely, we take some parameter $\sigma^2 > 0$, let ∂_{θ} be the differentiation operator in θ and consider, similarly to (1.2), the expression

(1.4)

$$T_{n}(\theta; \sigma^{2}) := \exp\left\{-\frac{1}{2}\sigma^{2}\partial_{\theta}^{2}\right\} \left(\sin\frac{\theta}{2}\right)^{n}$$

$$= (2 i)^{-n} \exp\left\{-\frac{1}{2}\sigma^{2}\partial_{\theta}^{2}\right\} \frac{(e^{i\theta}-1)^{n}}{e^{in\theta/2}}$$

$$= (2 i)^{-n} \exp\left\{-\frac{1}{2}\sigma^{2}\partial_{\theta}^{2}\right\} \sum_{j=0}^{n} (-1)^{n-j} {n \choose j} e^{i\theta(j-\frac{n}{2})}$$

$$= (2 i)^{-n} \sum_{j=0}^{n} (-1)^{n-j} {n \choose j} e^{\frac{1}{2}\sigma^{2}(j-\frac{n}{2})^{2}} e^{i\theta(j-\frac{n}{2})}.$$

In the last line we used the identity $e^{-\frac{1}{2}\sigma^2\partial_{\theta}^2} e^{ic\theta} = e^{\frac{1}{2}\sigma^2c^2} e^{ic\theta}$. The algebraic polynomials corresponding to these trigonometric polynomials via (1.3) are given by

(1.5)
$$H_n(z;\sigma^2) = \sum_{j=0}^n (-1)^{n-j} \binom{n}{j} \exp\left\{-\frac{\sigma^2 j(n-j)}{2}\right\} z^j, \quad n \in \mathbb{N},$$

up to a multiplicative constant which was chosen to make $H_n(z; \sigma^2)$ monic.

If we view $t := \sigma^2$ as the time, then the trigonometric polynomial $T_n(\theta; t)$ satisfies the backward heat equation $\partial_t T_n(\theta; t) = -\frac{1}{2} \partial_{\theta}^2 T_n(\theta; t)$, while $H_n(z; t)$ satisfies the PDE

$$\partial_t H_n(z;t) = -\frac{1}{2}(z\partial_z)(n-z\partial_z)H_n(z;t), \quad H_n(z;0) = (z-1)^n,$$

TOME 8 (2025)

very similar to the heat-type PDE's that appeared in [Tao18] and [HH22, Section 2.3]. Indeed, using the fact that $(z\partial_z)(n-z\partial_z)z^j = j(n-j) \cdot z^j$, we deduce

$$\exp\left\{-\frac{t}{2}(z\partial_z)(n-z\partial_z)\right\}(z-1)^n$$
$$=\sum_{j=0}^n(-1)^{n-j}\binom{n}{j}\exp\left\{-\frac{t}{2}(z\partial_z)(n-z\partial_z)\right\}z^j$$
$$=\sum_{j=0}^n(-1)^{n-j}\binom{n}{j}\exp\left\{-\frac{t}{2}j(n-j)\right\}z^j=H_n(z;t).$$

1.2. Connection to finite free probability

In the following, we shall refer to the polynomials $H_n(z; \sigma^2)$ defined by (1.5) as unitary Hermite polynomials with parameter $\sigma^2 > 0$. These polynomials appeared in the work of Mirabelli [Mir21] on finite free probability, a theory developed by Marcus, Spielman, Srivastava [MSS22] and Marcus [Mar18]. This theory studies the finite free additive convolution \boxplus_n and the finite free multiplicative convolution \boxtimes_n which are bilinear operations on the space of algebraic polynomials of degree at most n defined [Mar18, MSS22] as follows:

(1.6)
$$\left(\sum_{i=0}^{n} \frac{\alpha_i}{i!} z^i\right) \boxplus_n \left(\sum_{j=0}^{n} \frac{\beta_j}{j!} z^j\right) = \frac{1}{n!} \sum_{\ell=0}^{n} \frac{z^\ell}{\ell!} \sum_{\substack{i,j \in \{0,\dots,n\}\\i+j=n+\ell}} \alpha_i \beta_j,$$

(1.7)
$$\left(\sum_{j=0}^{n} \alpha_j z^j\right) \boxtimes_n \left(\sum_{j=0}^{n} \beta_j z^j\right) = \sum_{j=0}^{n} (-1)^{n-j} \frac{\alpha_j \beta_j}{\binom{n}{j}} z^j$$

It has been shown in [Mar18, Mir21, MSS22] that there is an analogue of the central limit theorem for these convolutions (for every fixed $n \in \mathbb{N}$). The classical Hermite polynomials play the role of the normal distribution for \boxplus_n ; see [Mar18, Theorem 6.7] and [Mir21, Theorems 3.2, 3.5]. Similarly, the unitary Hermite polynomials $H_n(z; \sigma^2/(n-1))$ play the role of the normal distribution for \boxtimes_n ; see [Mir21, Theorems 3.16, 3.23, 3.32]. For example, the analogue of the de Moivre–Laplace theorem for \boxtimes_n is as follows. Fix some even number n = 2d and consider degree 2d polynomials

$$Q_N(z) := \left(z^2 - 2z\cos(\sigma/\sqrt{N}) + 1\right)^d, \qquad N \in \mathbb{N},$$

having two zeroes at $e^{i\sigma/\sqrt{N}}$ and $e^{-i\sigma/\sqrt{N}}$, both of multiplicity d. Then, it can be shown that

$$\lim_{N \to \infty} \underbrace{Q_N(z) \boxtimes_{2d} \dots \boxtimes_{2d} Q_N(z)}_{N \text{ times}} = H_{2d} \left(z; \frac{\sigma^2}{2d-1} \right).$$

The operations \boxplus_n (respectively, \boxtimes_n) are known (in a suitable sense) to converge, as $n \to \infty$, to the classical free additive (respectively, multiplicative) convolutions \boxplus , respectively, \boxtimes . For information on (infinite) free probability we refer to [NS06, VDN92] for its finite counterpart to [AP18, AGVP23, Mar18, Mir21, MSS22].

1.3. Connection to the Curie–Weiss model

It has been observed by Mirabelli [Mir21, Section 3.2.5] that the unitary Hermite polynomials are closely related to the *Curie–Weiss model* (or the Ising model on the complete graph), which is one of the simplest models of statistical mechanics. The partition function of the Curie–Weiss model at inverse temperature $\beta > 0$ and with external magnetic field $h \in \mathbb{R}$ is given by

(1.8)
$$Z_n(\beta, h) = \sum_{(\sigma_1, \dots, \sigma_n) \in \{\pm 1\}^n} e^{\frac{\beta}{2n} \left(\sum_{k=1}^n \sigma_k\right)^2 + h \sum_{k=1}^n \sigma_k}$$

For every $j \in \{0, ..., n\}$ there exist $\binom{n}{j}$ configurations $(\sigma_1, ..., \sigma_n)$ in which the number of +1's is j. Since for every such configuration we have $\sum_{k=1}^{n} \sigma_k = 2j - n$, the above partition function can be written as

(1.9)
$$Z_n(\beta,h) = \sum_{j=0}^n \binom{n}{j} e^{\frac{\beta}{2n}(2j-n)^2 + h(2j-n)} = H_n\left(-e^{2h};\frac{4\beta}{n}\right) \cdot e^{n(\frac{\beta}{2}-h)} \cdot (-1)^n.$$

The fact that various quantities related to $Z_n(\beta, h)$ satisfy PDE's (the heat equation, the Burgers equation and a Hamilton–Jacobi equation) is known since Newman [New86]; see also [DM24, DMTW19].

The behavior of the Curie–Weiss model at *real* parameters β and *h* is very well understood; see [EN78a, EN78b, ENR80] as well as the books by Ellis [Ell06, Sections IV.4, V.9] and Friedli and Velenik [FV18, Chapter 2]. For a recent approach using the theory of mod- ϕ -convergence we refer to [MN15].

The behavior at *complex* parameters β and h, and in particular the location of the complex zeroes of $Z_n(\beta, h)$ is also of interest and has attracted attention in the theoretical physics literature [DBF20, DF20, GPS86, KBHK16]; see also the recent paper [HJN23]. These authors were motivated by the Lee–Yang program [Fis65, LY52, YL52] which relates phase transitions to the complex zeroes of the partition function. Shamis and Zeitouni [SZ18] analyzed the partition function and its zeroes at complex β (with h = 0) in a small neighborhood of the critical value $\beta = 1$, while the behavior outside this neighborhood remains largely unknown. The results of the present paper clarify the asymptotic behavior of $Z_n(\beta, h)$ at complex h (with fixed real $\beta > 0$) and, in particular, identify the global limiting distribution of the complex zeroes of $Z_n(\beta, h)$. Thus, we analyse the so-called Lee–Yang zeroes in contrast to the Fisher zeroes analyzed in [SZ18]. Partition function zeroes of the Ising model on Cayley trees and hierarchical lattices have been studied in [BLR17, BLR20, CHJR19], where pointers to earlier literature and a discussion of conjectures on the Ising model on \mathbb{Z}^d can be found.

1.4. Summary of results

The main results of the present paper, to be stated in Section 2, can be summarized as follows:

- (a) We prove that the empirical distribution of zeroes of $H_n(z; \sigma^2/n)$ converges weakly on the unit circle to the free unitary normal distribution \mathfrak{N}_{σ^2} , thereby identifying the limiting distribution of the Lee–Yang zeroes of the Curie–Weiss model; see Theorem 2.2 and Corollary 2.7.
- (b) We compute the asymptotics of $H_n(z; \sigma^2/n)$ for complex z with $|z| \neq 1$, thereby identifying the free energy of the Curie–Weiss model with complex external field; see Theorem 2.4 and Corollary 2.9.
- (c) It is well known [PB21, Section 6.1.2] that the expected characteristic polynomial of a Wigner random matrix of size n coincides with the n^{th} classical Hermite polynomial. We prove a unitary analogue of this result. More precisely, let $(U_n(t))_{t\geq 0}$ be a Brownian motion on the unitary group U(n) such that $U_n(0) = I_n$ is the identity matrix. In Theorem 2.10 we show that

$$\mathbb{E}\det(xI_n - U_n(t)) = e^{-nt/2} H_n\left(e^{t/2}x; t\right), \qquad t \ge 0.$$

An extension to Brownian motion starting at an arbitrary unitary matrix is given in Corollary 2.11.

- (d) Consider a high-degree polynomial (respectively, trigonometric polynomials) whose roots are real and have certain asymptotic distribution. We show that applying the backward heat flow to the polynomial is equivalent, on the level of the asymptotic distribution of the roots, to starting a free Brownian motion (respectively, a free unitary Brownian motion) from the initial distribution of roots; see Theorems 2.13 and 2.16.
- (e) We review the properties of the free unitary normal distribution \mathfrak{N}_{σ^2} and derive some new ones; see Theorem 2.18 and Proposition 2.21.

Notation

Throughout the paper, $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$ denotes the open unit disk, $\mathbb{T} = \{z \in \mathbb{C} : |z| = 1\}$ the unit circle, and $\mathbb{H} = \{\theta \in \mathbb{C} : \operatorname{Im} \theta > 0\}$ the upper half-plane. The closures of \mathbb{D} and \mathbb{H} are denoted by $\overline{\mathbb{D}}$ and $\overline{\mathbb{H}}$, respectively. We write $a_n \sim b_n$ if $a_n/b_n \to 1$ as $n \to \infty$. Weak and vague convergence of measures are denoted by \xrightarrow{w} and \xrightarrow{v} , respectively.

2. Main results

2.1. Empirical distribution of zeroes

The empirical distribution of zeroes of an algebraic polynomial $P_n(z)$ of degree n, i.e. the probability measure assigning to each zero the same weight 1/n, will be denoted by

(2.1)
$$\mu\llbracket P_n \rrbracket := \frac{1}{n} \sum_{z \in \mathbb{C} : P_n(z) = 0} \delta_z.$$



Figure 2.1. Zeroes of the unitary Hermite polynomials $H_n(z; \sigma^2/n)$ with $\sigma^2 = 1, 2, 3, 4$. The degree is n = 200.

We agree that the roots are always counted with multiplicities. It is well known, see, e.g, [Ull80, Gaw87, KM16], that the empirical distribution of zeroes of the classical Hermite polynomial $\text{He}_n(z\sqrt{n})$ converges weakly to the Wigner distribution $\gamma_{0,2}$ with the density $x \mapsto \frac{1}{2\pi}\sqrt{4-x^2}$ on the interval [-2, 2], namely

$$\left[\!\left[\operatorname{He}_{n}(\cdot\sqrt{n})\right]\!\right] = \frac{1}{n} \sum_{\substack{z \in \mathbb{C}:\\\operatorname{He}_{n}(z)=0}} \delta_{z/\sqrt{n}} \xrightarrow{w}_{n \to \infty} \gamma_{0,2}.$$

Given that the Wigner law is the analogue of the normal distribution w.r.t. the free additive convolution \boxplus , one may conjecture that the limiting empirical distribution of zeroes of the *unitary* Hermite polynomials should be related to the analogue of the normal distribution w.r.t. the free *multiplicative* convolution \boxtimes . We shall confirm this intuition. We begin by recording the following important property.

LEMMA 2.1. — All zeroes of the polynomial $H_n(z; \sigma^2)$ are located on the unit circle $\mathbb{T} = \{|z| = 1\}$.

Proof. — The claim is a special case of the Lee–Yang theorem; see [LY52, Appendix II] (where one takes $x_{\alpha\beta} := e^{-\sigma^2/2}$ for all $\alpha, \beta = 1, \ldots, n$) or [Rue69, Section 5.1]. Alternatively, the claim can be deduced from the Pólya–Benz theorem [ABH04, Theorem 1.2] applied to the periodic function $f(\theta) = (\sin \frac{\theta}{2})^n$ and the differential operator $\exp\{-\frac{1}{2}\sigma^2\partial_{\theta}^2\}$ (see the remarks preceding Corollary 1.3 in [ABH04] regarding applicability to non-polynomials). The Pólya–Benz theorem implies that all zeroes of $\exp\{-\frac{1}{2}\sigma^2\partial_{\theta}^2\}(\sin \frac{\theta}{2})^n$ are real. Recalling (1.4) completes the proof.

The empirical distribution of zeroes of $H_n(z; \sigma^2/n)$ will be denoted by

(2.2)
$$\mu_{n;\sigma^2} := \mu \left[\left[H_n(\cdot; \sigma^2/n) \right] \right] = \frac{1}{n} \sum_{z \in \mathbb{T}: H_n(z; \sigma^2/n) = 0} \delta_z.$$

THEOREM 2.2. — Fix some $\sigma^2 > 0$. Then, as $n \to \infty$, the probability measures $\mu_{n;\sigma^2}$ converge weakly on \mathbb{T} to the free unitary normal distribution \mathfrak{N}_{σ^2} with parameter σ^2 ; see Section 2.6 for its definition and properties.

Attempting to prove Theorem 2.2 by the method of moments leads to non-trivial combinatorics. After the preprint version of this paper appeared, such proof has been given in [AFU24].

2.2. Asymptotics of unitary Hermite polynomials

Our asymptotic results on the polynomials $H_n(z; \sigma^2/n)$ will be stated in terms of certain analytic function $\zeta_t(\theta)$ that satisfies

$$\zeta_t(\theta) - t \tan \zeta_t(\theta) = \theta_t$$

where t > 0 is a parameter and θ is a complex variable satisfying Im $\theta > 0$. This function is related to the free unitary Poisson distribution, as has been shown in [Kab21], and to the free unitary normal distribution, which will be demonstrated in Section 2.6 below. The next theorem summarizes the main properties of this function; see [Kab21, Section 4] for proofs and further properties.

THEOREM 2.3. — Fix t > 0. Let $\mathbb{H} := \{\theta \in \mathbb{C} : \operatorname{Im} \theta > 0\}$ be the upper halfplane. For every $\theta \in \mathbb{H}$, the equation $\zeta - t \tan \zeta = \theta$ has a unique, simple solution $\zeta = \zeta_t(\theta)$ in \mathbb{H} . The function $\zeta_t : \mathbb{H} \to \mathbb{H}$ is analytic on \mathbb{H} , admits a continuous extension to the closed upper half-plane $\overline{\mathbb{H}}$, and satisfies

(2.3)
$$\zeta_t(\theta + \pi) = \zeta_t(\theta) + \pi, \qquad \zeta_t(-\bar{\theta}) = -\overline{\zeta_t(\theta)}, \qquad \text{Im } \zeta_t(\theta) > \text{Im } \theta,$$

for all $\theta \in \mathbb{H}$. Locally uniformly in $x \in \mathbb{H}$ we have

$$\zeta_t(\theta) = \lim_{n \to \infty} \underbrace{\left(\theta + t \tan(\theta + t \tan(\dots (\theta + t \tan x) \dots))\right)}_{n \text{ iterations}}, \qquad \theta \in \overline{\mathbb{H}}.$$

Finally, we have

(2.4) $\zeta_t(\theta) - \theta \to it \quad as \quad \text{Im}\,\theta \to +\infty \quad \text{uniformly in } \text{Re}\,\theta \in \mathbb{R}.$

The next theorem is our second main result.

THEOREM 2.4. — Locally uniformly in $\theta \in \mathbb{H}$ we have

(2.5)
$$\lim_{n \to \infty} \frac{1}{n} \log \frac{H_n\left(-\mathrm{e}^{\mathrm{i}\,\theta}; \sigma^2/n\right)}{(-1)^n} = \log\left(1 + \mathrm{e}^{2\,\mathrm{i}\,\zeta_{\sigma^2/4}(\theta/2)}\right) - \frac{\sigma^2}{2\left(1 + \mathrm{e}^{-2\,\mathrm{i}\,\zeta_{\sigma^2/4}(\theta/2)}\right)^2},$$

$$\lim_{n \to \infty} \frac{1}{n} \frac{H'_n\left(-e^{i\theta}; \sigma^2/n\right)}{H_n\left(-e^{i\theta}; \sigma^2/n\right)} = \frac{-e^{-i\theta}}{1 + e^{-2i\zeta_{\sigma^2/4}(\theta/2)}} = -e^{-i\theta}\left(\frac{i}{2}\tan\zeta_{\sigma^2/4}(\theta/2) + \frac{1}{2}\right).$$

The logarithms in (2.5) are chosen such that $\log 1 = 0$ and all functions of the form $\log(\ldots)$ are continuous (and analytic) in $\theta \in \mathbb{H}$.

Remark 2.5. — We can consider all functions appearing in (2.5) and (2.6) as analytic functions of the variable $z := -e^{i\theta} \in \mathbb{D}$ including the value z = 0. Firstly, $z \to 0$ is equivalent to $\operatorname{Im} \theta \to +\infty$. In this regime, (2.4) implies that $\zeta_{\sigma^2/4}(\theta/2) =$ $(\theta/2) + i(\sigma^2/4) + o(1)$ and, consequently, $\operatorname{Im} \zeta_{\sigma^2/4}(\theta/2) \to +\infty$. It follows that the right-hand side of (2.5) converges to 0, while the right-hand side of (2.6) converges to $-e^{-\sigma^2/2}$. Secondly, note that θ corresponding to a given $z \in \mathbb{D} \setminus \{0\}$ is defined only up to a summand of the form $2\pi n$ with $n \in \mathbb{Z}$. Still, the right-hand sides of (2.5) and (2.6) stay invariant under the substitution $\theta \mapsto \theta + 2\pi n$ (since $\zeta_{\sigma^2/4}(\theta/2 + \pi n) =$ $\zeta_{\sigma^2/4}(\theta/2) + \pi n$ by (2.3)) and hence define analytic functions of $z \in \mathbb{D}$. Analogous observations apply to many similar functions below. Note that convergence in (2.5) and (2.6) stays locally uniform in $z \in \mathbb{D}$. For z outside any small disk around 0, this is stated in Theorem 2.4, while the rest follows from Cauchy's integration formula.

Remark 2.6. — Theorem 2.4 describes the asymptotics of $H_n(z; \sigma^2/n)$ for |z| < 1. The asymptotics for |z| > 1 can be derived from the identity $z^n H_n(1/z; \sigma^2/n) = (-1)^n H_n(z; \sigma^2/n)$ following from (1.5). On the circle $\{|z| = 1\}$ one may expect an asymptotic result of Plancherel–Rotach type; see [Sze75, Theorem 8.22.9] for the case of the classical Hermite polynomials.

2.3. Applications to the Curie–Weiss model

We are now going to describe the global limiting distribution of zeroes of $Z_n(\beta, h)$, the partition function of the Curie–Weiss model defined in (1.8). We consider the so-called Lee–Yang zeroes, that is we fix real $\beta > 0$ and allow h to be complex. By the Lee–Yang theorem, all zeroes are purely imaginary; see (1.9) and Lemma 2.1. Observe also that $Z_n(\beta, h + \pi i) = e^{\pi i n} Z_n(\beta, h)$ by (1.8) implying that the zeros are periodic with period πi .

COROLLARY 2.7. — Fix $\beta > 0$. For the partition function of the Curie–Weiss model, the following convergence holds vaguely on \mathbb{R} :

$$\frac{1}{n} \sum_{y \in \mathbb{R} : Z_n(\beta, \mathbf{i} y) = 0} \delta_y \xrightarrow[n \to \infty]{v} \nu_\beta$$

Here, ν_{β} is a measure on \mathbb{R} which is invariant under the shifts $h \mapsto h + \pi \ell$, $\ell \in \mathbb{Z}$, and is characterized by $\nu_{\beta}(A) = \mathfrak{N}_{4\beta}(-e^{2iA})$ for every Borel set $A \subset (-\frac{\pi}{2}, \frac{\pi}{2}]$, where $\mathfrak{N}_{4\beta}$ is the free unitary normal distribution on the unit circle with parameter $\sigma^2 = 4\beta$; see Section 2.6.

Proof. — Let $f : \mathbb{R} \to \mathbb{R}$ be a continuous function with compact support. Define $f^*(y) := \sum_{\ell \in \mathbb{Z}} f(y + \pi \ell)$ for $y \in (-\frac{\pi}{2}, \frac{\pi}{2}]$. Let also $\psi : \mathbb{T} \to (-\frac{\pi}{2}, \frac{\pi}{2}]$ be the inverse map of $y \mapsto -e^{2iy}$. Then, by (1.9),

$$\frac{1}{n} \sum_{y \in \mathbb{R} : Z_n(\beta, i y) = 0} f(y) = \frac{1}{n} \sum_{y \in (-\frac{\pi}{2}, \frac{\pi}{2}] : H_n\left(-e^{2 i y}, \frac{4\beta}{n}\right) = 0} f^*(y) = \frac{1}{n} \sum_{z \in \mathbb{T} : H_n\left(z, \frac{4\beta}{n}\right) = 0} f^*(\psi(z)).$$

By Theorem 2.2, the latter sum converges to $\int_{\mathbb{T}} f^*(\psi(z)) \mathfrak{N}_{4\beta}(dz) = \int_{\mathbb{R}} f(y) \nu_{\beta}(dy)$, and the claim follows.

Remark 2.8. — The Lebesgue density of ν_{β} on \mathbb{R} is given by $y \mapsto 2f_{4\beta}(-e^{2iy}) = \frac{1}{\pi\beta} \operatorname{Im} \zeta_{\beta}(y)$, where $f_{4\beta}$ is the function which will be discussed in Theorem 2.18. It follows from this theorem that the support of ν_{β} is \mathbb{R} for $\beta \ge 1$, while for $0 < \beta < 1$ the support is the union of the intervals

$$\left[\frac{\pi}{2} - \arcsin\sqrt{\beta} - \sqrt{\beta - \beta^2} + \pi\ell, \frac{\pi}{2} + \arcsin\sqrt{\beta} + \sqrt{\beta - \beta^2} + \pi\ell\right], \quad \ell \in \mathbb{Z}.$$

If β increases from 0 to ∞ , then the support of ν_{β} hits the real axis at $\beta = 1$, which is well known to be the point of phase transition for the Curie–Weiss model.

In the next result we compute the free energy of the Curie–Weiss model in the complex h-plane excluding the imaginary axis.

COROLLARY 2.9. — Let $\beta > 0$ and $h \in \mathbb{C}$ with $\operatorname{Re} h > 0$. For the partition function of the Curie–Weiss model defined in (1.8) we have

$$\lim_{n \to \infty} \frac{1}{n} \log Z_n(\beta, h) = \lim_{n \to \infty} \frac{1}{n} \log Z_n(\beta, -h)$$
$$= \frac{\beta}{2} + h + \log \left(1 + e^{2i\zeta_\beta(ih)}\right) - \frac{2\beta}{\left(1 + e^{-2i\zeta_\beta(ih)}\right)^2}$$

Proof. — Recall that $Z_n(\beta, h) = Z_n(\beta, -h)$ is given by (1.8) and (1.9) and apply Theorem 2.4 with $\sigma^2 = 4\beta$ and $\theta = 2ih$.

2.4. Expected characteristic polynomial of the Brownian motion on unitary matrices

It is well known [PB21, Section 6.1.2] that the expected characteristic polynomial of an $n \times n$ Wigner random matrix coincides with the n^{th} Hermite polynomial. Formulas of this type go back to Heine [Sze75, (2.2.11) p. 27]. For this and analogous results on several other types of random matrices including the Wishart matrices whose expected characteristic polynomials are the Laguerre polynomials we refer to [PB21, Sections 6.2, 6.3], [Aom87], [Ede89, Chapter 9], [DE02, Theorem 4.1], [BH00, (15)], [DG04, Proposition 12], [FG06, Proposition 11], [AGN21, Theorem 1.1].

In this section we prove a similar result on unitary Hermite polynomials by relating them to the expected characteristic polynomials of the random matrices obtained by running a Brownian motion on the unitary group U(n). More precisely, we consider the unitary group U(n) as a compact Riemannian manifold endowed with the Riemannian metric induced by its natural embedding into $\mathbb{C}^{n \times n} \equiv \mathbb{R}^{2n^2}$. On the Lie algebra $\mathfrak{u}(n) = \{A \in \operatorname{Mat}_{n \times n}(\mathbb{C}) : A^* = -A\}$ (which can be identified with the tangent space of U(n) at the identity matrix I_n) the scalar product takes the form $\langle A, B \rangle = \operatorname{Tr}(AB^*) = -\operatorname{Tr}(AB)$.

Let now $(U_n(t))_{t\geq 0}$ be the Brownian motion on the unitary group U(n) starting at the identity matrix I_n at time t = 0. The eigenvalues $\lambda_1(t), \ldots, \lambda_n(t)$ of the unitary random matrix $U_n(t)$ represent a special case of Dyson's Brownian motions [Dys62, Section III] on the circle; see also [Bia09, Bui24, CL01, FKLZ24, Hal18, HW96] for further information on this process.

To define Dyson's Brownian motions on the circle, fix parameters $n \in \mathbb{N}$ and $\lambda > 0$. Let $(B_1(t))_{t \ge 0}, \ldots, (B_n(t))_{t \ge 0}$ be *n* independent standard Brownian motions on \mathbb{R} . We are interested in real-valued stochastic processes $X_1(t) \le \ldots \le X_n(t)$, defined for $t \ge 0$ and solving stochastic differential equations

(2.7)
$$dX_j = dB_j + \lambda \cdot \left(\sum_{\substack{k \in \{1,\dots,n\}\\k \neq j}} \cot \frac{X_j - X_k}{2}\right) dt, \quad j = 1,\dots,n,$$

with the initial condition $X_1(0) = \ldots = X_n(0) = 0$. If $\lambda = 1/2$, then we can identify $e^{i X_1(t)}, \ldots, e^{i X_n(t)}$ with the eigenvalues $\lambda_1(t), \ldots, \lambda_n(t)$ of the unitary random matrix $U_n(t)$. More precisely, it is known that the measure-valued process

$$\left(\sum_{\ell=1}^n \delta_{\mathrm{e}^{\mathrm{i}\,X_\ell(t)}}\right)_{t\,\geqslant\,0}$$

has the same distribution as the process

$$\left(\sum_{\ell=1}^n \delta_{\lambda_\ell(t)}\right)_{t \ge 0}.$$

We are interested in the following polynomial in x which, for $\lambda = 1/2$, reduces to the characteristic polynomial of $U_n(t)$:

$$P_{n,\lambda}(x;t) := \prod_{j=1}^{n} \left(x - \mathrm{e}^{\mathrm{i} X_j(t)} \right).$$

THEOREM 2.10. — For every $\lambda > 0, n \in \mathbb{N}, t \ge 0$ and $x \in \mathbb{C}$ we have

$$\mathbb{E}P_{n,\lambda}(x;t) = \sum_{j=0}^{n} (-1)^{n-j} \binom{n}{j} e^{-\frac{1}{2}(n-j)t - \lambda j(n-j)t} x^{j} = e^{-\frac{1}{2}nt} H_n\left(e^{t/2}x; 2\lambda t\right)$$

Proof. — To simplify the notation, we shall usually suppress the dependence of quantities under consideration on n and λ . Let $e_{\ell}(t)$ be the ℓ^{th} elementary symmetric polynomial of $e^{i X_1(t)}, \ldots, e^{i X_n(t)}$, that is

$$e_{\ell}(t) = \sum_{1 \leq j_1 < \dots < j_{\ell} \leq n} e^{i X_{j_1}(t) + \dots + i X_{j_{\ell}}(t)}, \qquad \ell = 1, \dots, n,$$

Put also $e_0(t) = 1$. Since $P_{n,\lambda}(x;t) = \sum_{\ell=0}^n (-1)^\ell e_\ell(t) x^{n-\ell}$ by Vieta's formula, it suffices to show that for all $\ell \in \{1, \ldots, n\}$ we have

$$\mathbb{E}e_{\ell}(t) = \binom{n}{\ell} e^{-\frac{1}{2}\ell t - \lambda\ell(n-\ell)t}$$

To this end, we shall derive stochastic differential equations satisfied by $e_{\ell}(t)$. Using the Itô formula, see, e.g., [RY99, Chapter IV, Theorem (3.3)], we have

$$de_{\ell} = \sum_{1 \leq j_1 < \dots < j_{\ell} \leq n} d\left(e^{iX_{j_1} + \dots + iX_{j_{\ell}}}\right)$$
$$= \sum_{1 \leq j_1 < \dots < j_{\ell} \leq n} \left(\sum_{s=1}^{\ell} i e^{iX_{j_1} + \dots + iX_{j_{\ell}}} dX_{j_s} - \frac{\ell}{2} e^{iX_{j_1} + \dots + iX_{j_{\ell}}} dt\right).$$

Write $V(x) := \lambda \cot \frac{x}{2}$. Recalling (2.7), we obtain

(2.8)
$$de_{\ell} = \sum_{1 \leq j_1 < \dots < j_{\ell} \leq n} i e^{i X_{j_1} + \dots + i X_{j_{\ell}}} (dB_{j_1} + \dots + dB_{j_{\ell}}) - \frac{\ell}{2} e_{\ell} dt + R dt$$

TOME 8 (2025)

with

$$R := \sum_{\substack{1 \leq j_1 < \dots < j_\ell \leq n}} i e^{i X_{j_1} + \dots + i X_{j_\ell}} \\ \begin{pmatrix} \sum_{\substack{m \in \{1,\dots,n\} \\ m \neq j_1}} V(X_{j_1} - X_m) + \dots + \sum_{\substack{m \in \{1,\dots,n\} \\ m \neq j_\ell}} V(X_{j_\ell} - X_m) \end{pmatrix} \\ = \sum_{\substack{1 \leq j_1 < \dots < j_\ell \leq n}} i e^{i X_{j_1} + \dots + i X_{j_\ell}} \\ \begin{pmatrix} \sum_{\substack{m \in \{1,\dots,n\} \setminus \{j_1,\dots,j_\ell\}}} \left(V(X_{j_1} - X_m) + \dots + V(X_{j_\ell} - X_m) \right) \end{pmatrix} \end{cases}$$

where in the second line we used that V(-x) = -V(x). After some re-indexing, we can write

,

$$R = \frac{i}{(\ell+1)!} \sum_{\substack{k_0, \dots, k_{\ell} \in \{1, \dots, n\} \\ \text{pairwise distinct}}} e^{iX_{k_0} + iX_{k_1} + \dots + iX_{k_{\ell}}}$$
$$\sum_{\substack{s, p \in \{0, \dots, \ell\} \\ s \neq p}} \frac{V(X_{k_p} - X_{k_s})}{e^{iX_{k_s}}}$$
$$= \frac{i}{2(\ell+1)!} \sum_{\substack{k_0, \dots, k_{\ell} \in \{1, \dots, n\} \\ \text{pairwise distinct}}} e^{iX_{k_0} + iX_{k_0} + \dots + iX_{k_{\ell}}}$$
$$\sum_{\substack{s, p \in \{0, \dots, \ell\} \\ s \neq p}} \left(\frac{V(X_{k_p} - X_{k_s})}{e^{iX_{k_s}}} + \frac{V(X_{k_s} - X_{k_p})}{e^{iX_{k_p}}} \right).$$

To simplify the expression in the brackets, note that

$$\frac{V(x-y)}{e^{iy}} + \frac{V(y-x)}{e^{ix}} = \lambda \cot\left(\frac{x-y}{2}\right) \left(e^{i(x-y)} - 1\right) e^{-ix} = i\lambda \left(e^{-ix} + e^{-iy}\right).$$

It follows that

$$R = \frac{-\lambda}{2(\ell+1)!} \sum_{\substack{k_0,\dots,k_\ell \in \{1,\dots,n\}\\\text{pairwise distinct}}} e^{i X_{k_0} + i X_{k_1} + \dots + i X_{k_\ell}} \sum_{\substack{s,p \in \{0,\dots,\ell\}\\s \neq p}} \left(e^{-i X_{k_s}} + e^{-i X_{k_p}} \right)$$
$$= -\lambda \ell(n-\ell) e_\ell(t).$$

To justify the last identity, observe that the double sum in the first line must be a multiple of e_{ℓ} for symmetry reasons and that it contains $(n)_{\ell+1} \cdot (\ell+1)\ell \cdot 2$ summands, while e_{ℓ} contains $\binom{n}{\ell}$ summands. Taking the quotient of these two numbers, it follows that the double sum in the first line equals $2(\ell+1)!\ell(n-\ell)e_{\ell}$.

Finally, recalling (2.8), we arrive at the stochastic differential equation

(2.9)
$$de_{\ell} = -\left(\frac{\ell}{2} + \lambda \ell (n-\ell)\right) e_{\ell} dt + \sum_{j=1}^{n} i e^{i X_j} e_{\ell-1}^{(j)} dB_j,$$

where $e_{\ell-1}^{(j)}$ is the $(\ell-1)^{\text{st}}$ elementary symmetric polynomial of $e^{iX_1}, \ldots, e^{iX_{j-1}}, e^{iX_{j+1}}, \ldots, e^{iX_n}$ (excluding e^{iX_j}). From the Itô formula it follows that $e^{\frac{1}{2}\ell t + \lambda\ell(n-\ell)t} e_{\ell}(t)$ is a martingale. Recalling that $e_{\ell}(0) = \binom{n}{\ell}$ we conclude that

$$\mathbb{E}e_{\ell}(t) = \binom{n}{\ell} e^{-\frac{1}{2}\ell t - \lambda\ell(n-\ell)t},$$

and the proof of Theorem 2.10 is complete.

Following a suggestion of an anonymous referee let us extend the above result to the Brownian motion $(VU_n(t))_{t\geq 0}$ on the unitary group U(n) starting from an arbitrary unitary matrix $V \in U(n)$.

COROLLARY 2.11. — For every unitary matrix $V \in U(n)$ and all $t \ge 0$ and $z \in \mathbb{C}$ we have

(2.10)
$$Q_n(z;t) = \mathbb{E} \det(zI_n - VU_n(t)) = \det(zI_n - V) \boxtimes_n e^{-\frac{1}{2}nt} H_n\left(e^{t/2}z;t\right).$$

Also, the following PDE holds:

(2.11)
$$\partial_t Q_n(z;t) = -\frac{1}{2}(z\partial_z + 1)(n - z\partial_z)Q_n(z;t), \qquad Q_n(z;0) = \det(zI_n - V).$$

Proof. — It is possible to prove (2.10) by changing the initial condition for $e_{\ell}(t)$ in the proof of Theorem 2.10, however, we find it more instructive to give a proof that uses finite free probability. Let W be a Haar-distributed random matrix in U(n) which is independent of everything else. Then, $VU_n(t)$ has the same law as $VW^{-1}U_n(t)W$. Hence,

$$\mathbb{E}\det(zI_n - VU_n(t)) = \mathbb{E}\left[\det\left(zI_n - VW^{-1}U_n(t)W\right)\right]$$
$$= \mathbb{E}\left[\mathbb{E}\left[\det\left(zI_n - VW^{-1}U_n(t)W\right) \middle| U_n(t)\right]\right].$$

In the conditional expectation, $U_n(t)$ is fixed and the integration is over the distribution of W, i.e. the Haar distribution on U(n). By [MSS22, Theorem 1.5], the conditional expectation can be expressed using \boxtimes_n , which leads to

$$\mathbb{E} \det(zI_n - VU_n(t)) = \mathbb{E} \Big[\det(zI_n - V) \boxtimes_n \det(zI_n - U_n(t)) \Big]$$
$$= \det(zI_n - V) \boxtimes_n \mathbb{E} \det(zI_n - U_n(t))$$
$$= \det(zI_n - V) \boxtimes_n e^{-\frac{1}{2}nt} H_n \left(e^{t/2} z; t\right).$$

We used the bilinearity of \boxtimes_n and Theorem 2.10 with $\lambda = 1/2$. The proof of (2.10) is complete.

Since \boxtimes_n commutes with $z\partial_z$, see [MSS22, p. 810], it suffices to prove the PDE (2.11) for $Q_n(z;t) = e^{-\frac{1}{2}nt} H_n(e^{t/2}z;t)$ with the initial condition $Q_n(z;0) = (z-1)^n$, which corresponds to the case when $V = I_n$. Using that $(z\partial_z + 1)(n - z\partial_z)z^j = (j+1)(n-j) \cdot z^j$, we deduce

$$\exp\left\{-\frac{t}{2}(z\partial_{z}+1)(n-z\partial_{z})\right\}(z-1)^{n}$$

$$=\sum_{j=0}^{n}(-1)^{n-j}\binom{n}{j}\exp\left\{-\frac{t}{2}(z\partial_{z}+1)(n-z\partial_{z})\right\}z^{j}$$

$$=\sum_{j=0}^{n}(-1)^{n-j}\binom{n}{j}\exp\left\{-\frac{t}{2}(j+1)(n-j)\right\}z^{j}$$

$$=e^{-\frac{1}{2}nt}H_{n}\left(e^{t/2}z;t\right),$$

where we used (1.5) in the last step.

Remark 2.12. — The operator $(z\partial_z + 1)(n - z\partial_z) = -z^2\partial_z^2 + (n - 2)z\partial_z + n$ is (up to sign) the same as the one appearing in Conjecture 2.14 and Proposition 2.15 of Hall and Ho [HH22].

Biane [Bia97a, Bia97b] proved that, as $n \to \infty$, the process $(U_n(t/n))_{t\geq 0}$ converges (in a suitable sense) to the free unitary Brownian motion. In particular, by [Bia97a, Theorem 1], the spectral distribution of $U_n(t/n)$ converges weakly to the free unitary normal distribution \mathfrak{N}_t (making the appearance of this distribution in Theorem 2.2 quite natural); see also [CDG01, Section 3.3] for related large deviation results. Exact combinatorial formulas for moments of the form $\mathbb{E}[\operatorname{Tr}(U_n^{m_1}(t)) \dots \operatorname{Tr}(U_n^{m_r}(t))]$ have been derived in [Lév08].

2.5. The action of the backward heat flow on the roots

Consider a sequence of polynomials (or trigonometric polynomials) of increasing degrees whose empirical distributions of roots approach some probability measure. One may ask what happens to the asymptotic distribution of roots if we apply to these polynomials certain operator. One special case, in which the operator is the repeated differentiation, has been studied in [AGVP23, BHS24, Gal22, HK23, Kab21, KT20, OS21, Ste19, Ste20, Ste23]. For trigonometric polynomials, it has been shown in [Kab21] that, on the level of roots, the repeated differentiation induces the free unitary Poisson process. In his blog, Tao [Tao17, Tao18] discusses the evolution of zeroes of a polynomial which undergoes a (backward) heat flow. As this paper was almost complete, Jonas Jalowy brought to our attention the recent preprint by Hall and Ho [HH22] who studied the action of the backward heat flow on the characteristic polynomials of the Ginibre matrices (whose eigenvalues obey the circular law). We shall consider two settings: algebraic polynomials and trigonometric polynomials, both with real roots, and show that the backward heat flow induces free (additive or unitary) Brownian motion on the level of roots.

2.5.1. Heat flow acting on algebraic polynomials

Let $(P_n(z))_{n \in \mathbb{N}}$ be a sequence of algebraic polynomials from $\mathbb{R}[z]$. We suppose that $P_n(z) = \sum_{j=0}^n a_{j:n} z^j$ is real-rooted (that is, it has only real roots) and that all roots are contained in some fixed interval [-C, C] with C not depending on n. Moreover,

we suppose that the empirical distribution of roots of P_n converges weakly to some probability measure μ on [-C, C], that is

(2.12)
$$\mu\llbracket P_n \rrbracket = \frac{1}{n} \sum_{z \in \mathbb{R} : P_n(z) = 0} \delta_z \xrightarrow[n \to \infty]{w} \mu.$$

The roots are counted with multiplicities, as always. We are interested in the action which the backward heat flow induces on the roots of P_n , in the large *n* limit. More precisely, we consider the heat equation on the real line with initial condition given by $P_n(z)$:

(2.13)
$$\partial_t g_n(z;t) = \frac{1}{2} \partial_z^2 g_n(z;t), \quad g_n(z;0) = P_n(z), \quad z \in \mathbb{R}, \ t \in \mathbb{R}.$$

The solution is explicit and can be written as

(2.14)
$$g_n(z; -s) = e^{-\frac{s}{2}\partial_z^2} P_n(z) = \sum_{j=0}^n a_{j:n} e^{-\frac{s}{2}\partial_z^2} z^j = \sum_{j=0}^n a_{j:n} \operatorname{He}_j\left(\frac{z}{\sqrt{s}}\right) s^{j/2},$$

 $z \in \mathbb{R}, \ s \in \mathbb{R},$

where $\operatorname{He}_{j}(z)$ is the j^{th} probabilist Hermite polynomial defined by (1.1) or (1.2). Note that the solution exists both for positive and negative times since the term $\operatorname{He}_{j}(z/\sqrt{s})s^{j/2}$ does not contain fractional powers of s, see (1.1), and makes sense irrespective of the sign of s. Moreover, for every $s \in \mathbb{R}$, the function $z \mapsto g_{n}(z; -s)$ is a polynomial. In the sequel, we shall focus on the case s > 0, which corresponds to the backward heat equation. It is known to be ill-posed for initial conditions more general than polynomials. Since we assume that the initial condition $P_{n}(z)$ is real-rooted, the polynomials $g_{n}(z; -s)$ remain real-rooted for all $s \ge 0$ by the Pólya–Benz theorem; see [Ben34] or [ABH04, Theorem 1.2]. Another proof of this fact can be found [Tao17].

THEOREM 2.13. — Fix r > 0. Under the above assumptions, the empirical distribution of zeroes of the polynomials $g_n(z; -r^2/n)$ converges weakly (as $n \to \infty$) to the free additive convolution $\mu \boxplus \gamma_{0,2r}$ of μ and the Wigner semicircle distribution $\gamma_{0,2r}$ with density $x \mapsto \frac{1}{2\pi r^2}\sqrt{4r^2 - x^2}$ on the interval [-2r, 2r].

To prove this theorem, we shall use a result from finite free probability [Mar18, MSS22]. Recall from (1.6) the definition of the finite free additive convolution \boxplus_n . It is known from [AP18, Corollary 5.5 and Theorem 5.4], see also [Mar18, Theorem 4.3], that, as $n \to \infty$, the finite free additive convolution \boxplus_n approaches the free additive convolution \boxplus in the following sense.

PROPOSITION 2.14. — Let $(p_n)_{n \in \mathbb{N}}$ and $(q_n)_{n \in \mathbb{N}}$ be sequences of polynomials in $\mathbb{R}[z]$ whose roots are contained in some fixed interval [-C, C]. Suppose that deg $p_n = \deg q_n = n$ and the empirical distributions of zeroes of p_n and q_n converge weakly to certain probability measures ν and ρ on [-C, C]. Then, the empirical distribution of zeroes of $p_n \boxplus_n q_n$ (which is also real-rooted by [MSS22, Theorem 1.3]) converges weakly to $\nu \boxplus \rho$.

Indeed, $\mu[\![p_n]\!] \to \nu$ and $\mu[\![q_n]\!] \to \rho$ (weakly) implies that the moments of $\mu[\![p_n]\!]$ and $\mu[\![q_n]\!]$ converge to the corresponding moments of ν and ρ (since all measures are

concentrated on a fixed interval). By the results cited above, this implies that the moments of the empirical distribution of zeroes of $p_n \boxplus_n q_n$ converge to those of $\nu \boxplus \rho$. This implies that the probability measures $\mu \llbracket p_n \boxplus_n q_n \rrbracket$ (which are concentrated on [-2C, 2C] by [MSS22, Theorem 1.3]) converge weakly to $\nu \boxplus \rho$.

Proof of Theorem 2.13. It is known [MSS22] that \boxplus_n commutes with differentiation in the sense that $\partial_z(p(z) \boxplus_n q(z)) = (\partial_z p(z)) \boxplus_n q(z)$ for arbitrary polynomials p(z)and q(z) of degree at most n. By bilinearity of \boxplus_n it follows that it also commutes with the operator $\exp\{-\frac{s}{2}\partial_z^2\}$, for all $s \in \mathbb{R}$. Hence,

$$g_n(z; -s) = e^{-\frac{s}{2}\partial_z^2} P_n(z) = e^{-\frac{s}{2}\partial_z^2} \left(P_n(z) \boxplus_n z^n \right)$$
$$= P_n(z) \boxplus_n \left(e^{-\frac{s}{2}\partial_z^2} z^n \right) = P_n(z) \boxplus_n \operatorname{He}_n \left(\frac{z}{\sqrt{s}} \right) s^{n/2},$$

where we used (1.2) in the last equality. We take $s = r^2/n$ and let $n \to \infty$. By a classical result, the empirical distribution of zeroes of the Hermite polynomial $\operatorname{He}_n(z\sqrt{n}/r)$ converges weakly to the Wigner distribution with the density $x \mapsto \frac{1}{2\pi r^2}\sqrt{4r^2 - x^2}$ on the interval [-2r, 2r]; see, e.g., [Gaw87, KM16, Ull80]. Applying Proposition 2.14 completes the proof.

Remark 2.15. — Tao [Tao17] derived the following system of differential equations satisfied by the roots $z_{1:n}(s), \ldots, z_{n:n}(s)$ of the polynomial $g_n(z; -s)$:

(2.15)
$$\partial_s z_{i:n}(s) = \sum_{\substack{k \in \{1,\dots,n\}\\k \neq i}} \frac{1}{z_{i:n}(s) - z_{k:n}(s)}, \qquad i = 1,\dots,n,$$

and proved that the solutions are well-defined for $s \ge 0$ but may run into a singularity for $s \le 0$. These equations are Dyson's Brownian motions with vanishing variance (or with $\beta = \infty$); see [AGZ10, Theorem 4.3.2]. Therefore, one can view Theorem 2.13 as the $\beta = \infty$ case of [AGZ10, Proposition 4.3.10]. In this form, it has been established in [VW22, Theorem 1.1]. For other results relating Dyson's Brownian motions to free convolutions we refer to [AKM12, DE05, GK24, VW20, VW22].

2.5.2. Heat flow acting on trigonometric polynomials

Let us now state analogous results for trigonometric polynomials. For every even number n = 2d, $d \in \mathbb{N}$, let $T_n(\theta) := \sum_{\ell=-d}^d c_{\ell:n} e^{i\ell\theta}$ be a trigonometric polynomial with complex coefficients $c_{\ell:n}$ satisfying $c_{-\ell:n} = \overline{c_{\ell:n}}$ for all $\ell \in \{-d, \ldots, d\}$ (meaning that $T_n(\theta)$ takes real values for real θ). Moreover, assume that $T_n(\theta)$ is real-rooted meaning that it has n real roots $\theta_{1;n}, \ldots, \theta_{n;n}$ (counting multiplicities) and that its empirical distribution of zeroes converges weakly as $n \to \infty$. The latter assumption will be written in the following form:

(2.16)
$$\nu \llbracket T_n \rrbracket := \frac{1}{n} \sum_{j=1}^n \delta_{\mathrm{e}^{\mathrm{i}\,\theta_{j;\,n}}} \xrightarrow[n \to \infty]{w} \nu$$

weakly on the unit circle \mathbb{T} , for some probability measure ν on \mathbb{T} . Consider the heat equation on the real line with periodic initial condition given by $T_n(\theta)$:

(2.17)
$$\partial_t f_n(\theta; t) = \frac{1}{2} \partial_\theta^2 f_n(\theta; t), \quad f_n(\theta; 0) = T_n(\theta), \quad \theta \in \mathbb{R}, \ t \in \mathbb{R}.$$

Its solution is explicit and can be written as

(2.18)
$$f_n(\theta;t) = e^{\frac{t}{2}\partial_{\theta}^2} T_n(\theta) = \sum_{\ell=-d}^d c_{\ell:n} e^{\frac{t}{2}\partial_{\theta}^2} e^{i\ell\theta} = \sum_{\ell=-d}^d c_{\ell:n} e^{-\frac{t}{2}\ell^2} e^{i\ell\theta}$$

For every $t \in \mathbb{R}$, the function $\theta \mapsto f_n(\theta; t)$ is a trigonometric polynomial. In particular, the solution makes sense both for $t \ge 0$ and for t < 0. In the sequel we shall focus on the latter case, which corresponds to the backward heat equation. For s > 0, the trigonometric polynomial $f_n(\theta; -s)$ remains real-rooted. This claim can be deduced from the Pólya–Benz theorem [ABH04, Corollary 1.3]. Another proof can be found in [Tao18] (where differential equations analogous to (2.15) in the circular setting are derived; see also [CL01, HW96] for the corresponding stochastic differential equations).

THEOREM 2.16. — Take some $\sigma^2 > 0$. In the setting described above, including Assumption (2.16), the empirical distribution $\nu \llbracket f_n(\cdot; -\sigma^2/n) \rrbracket$ of zeroes of the solution $f_n(\theta; -\sigma^2/n)$ of the heat equation at time $t_n := -\sigma^2/n$, converges weakly (as $n = 2d \to \infty$) to the free unitary convolution $\nu \boxtimes \mathfrak{N}_{\sigma^2}$ of ν and the free unitary normal distribution \mathfrak{N}_{σ^2} .

We again rely on a result from finite free probability [MSS22, Mar18]. Recall from (1.7) the definition of the finite free multiplicative convolution \boxtimes_n . The following fact proved in [AGVP23, Proposition 3.4] (see also [Kab21, Proposition 2.9] for the version stated here) states that, in a suitable sense, \boxtimes_n converges to the free multiplicative convolution \boxtimes as $n \to \infty$.

PROPOSITION 2.17. — Let $(p_n(z))_{n \in \mathbb{N}}$ and $(q_n(z))_{n \in \mathbb{N}}$ be sequences of polynomials in $\mathbb{C}[z]$ with deg $p_n = \deg q_n = n$. Suppose that all roots of p_n and q_n are located on the unit circle and that, as $n \to \infty$, the empirical distributions of zeroes $\mu[\![p_n]\!]$ and $\mu[\![q_n]\!]$ converge weakly to two probability measures ν and ρ on \mathbb{T} . Then, all roots of the polynomial $p_n \boxtimes_n q_n$ are also located on \mathbb{T} and $\mu[\![p_n \boxtimes_n q_n]\!]$ converges weakly to $\nu \boxtimes \rho$.

Proof of Theorem 2.16. For $t \in \mathbb{R}$ we consider the algebraic polynomial $P_{2d}(z;t)$ in the complex variable z defined by

(2.19)
$$\frac{P_{2d}(e^{i\theta};t)}{e^{id\theta}} = f_{2d}(\theta;t),$$

More concretely, it follows from (2.18) that

$$P_{2d}(z;t) = \sum_{\ell=-d}^{d} c_{\ell:2d} e^{-\frac{t}{2}\ell^2} z^{\ell+d} = \sum_{j=0}^{2d} c_{j-d:2d} e^{-\frac{t}{2}(j-d)^2} z^j$$
$$= \sum_{j=0}^{2d} c_{j-d:s2d} e^{-\frac{t}{2}(j^2-2jd+d^2)} z^j.$$

TOME 8 (2025)



Figure 2.2. Densities of the free unitary normal distributions \mathfrak{N}_{σ^2} . The plots show the functions $\theta \mapsto f_{\sigma^2}(e^{i\theta})$ for $\theta \in [-\pi,\pi]$. Left: $\sigma^2 = 0.5, 1, 1.5, \ldots, 4$. Right: $\sigma^2 = 4, 4.5, 5, \ldots, 7$.

It follows from (1.7) and (1.5) that for $t = t_{2d} = -\sigma^2/(2d) < 0$ we can write

$$P_{2d}(z;t) = e^{-\frac{td^2}{2}} \left(\sum_{j=0}^{2d} (-1)^{2d-j} {2d \choose j} e^{\frac{t}{2}j(2d-j)} z^j \right) \boxtimes_{2d} \left(\sum_{j=0}^{2d} c_{j-d:2d} z^j \right)$$
$$= e^{-\frac{td^2}{2}} \left(H_{2d} \left(z; \frac{\sigma^2}{2d} \right) \boxtimes_{2d} P_{2d}(z;0) \right).$$

Recall from Theorem 2.2 that the empirical distribution of zeroes of $H_{2d}(z; \sigma^2/(2d))$ converges weakly to \mathfrak{N}_{σ^2} , while the empirical distribution of zeroes of $P_{2d}(z; 0)$ converges to ν by (2.19) and (2.16):

$$\frac{1}{2d} \sum_{z \in \mathbb{C} : H_{2d}\left(z; \sigma^2/(2d)\right) = 0} \delta_z \xrightarrow[d \to \infty]{w} \mathfrak{N}_{\sigma^2}, \qquad \frac{1}{2d} \sum_{z \in \mathbb{C} : P_{2d}(z; 0) = 0} \delta_z = \frac{1}{2d} \sum_{j=1}^{2d} \delta_{e^{i\theta_{j;2d}}} \xrightarrow[d \to \infty]{w} \nu.$$

To complete the proof, apply Proposition 2.17.

2.6. Free unitary normal distribution and its properties

In this section we recall the notion of free multiplicative convolution \boxtimes of probability measures on the unit circle which was introduced by Voiculescu in [Voi85] and [Voi87]; see also [VDN92, §3.6]. Also, we recall the definition of the free unitary normal distribution, which was introduced by Bercovici and Voiculescu [BV92] and studied in [Bia97a, Bia97b, Céb16, Zho14, Zho15]. Finally, we shall state some properties of this distribution.

Given two probability measures μ_1 and μ_2 on the unit circle \mathbb{T} , it is possible to construct a C^* -probability space and two mutually free unitaries u_1 and u_2 with spectral distributions μ_1 and μ_2 , respectively. Then, the free multiplicative convolution of μ_1 and μ_2 , denoted by $\mu_1 \boxtimes \mu_2$, is the spectral distribution of $u_1 u_2$. It does not depend on the choice of u_1 and u_2 ; see [VDN92] for more details. Free multiplicative convolution is linearized by the S-transform which is defined as follows. The ψ -transform of a probability measure μ on the unit circle \mathbb{T} is defined by

(2.20)
$$\psi_{\mu}(z) = \int_{\mathbb{T}} \frac{uz}{1 - uz} \mu(\mathrm{d}u) = \sum_{\ell=1}^{\infty} z^{\ell} \int_{\mathbb{T}} u^{\ell} \mu(\mathrm{d}u), \quad z \in \mathbb{D}$$

Note that $\psi_{\mu}(z)$ is an analytic function on \mathbb{D} satisfying $\psi_{\mu}(0) = 0$ and $\operatorname{Re} \psi_{\mu}(z) > -\frac{1}{2}$. These properties characterize the class of ψ -transforms; see [FHS20, Lemma 2.9].

If $\psi'_{\mu}(0) = \int_{\mathbb{T}} u\mu(\mathrm{d}u) \neq 0$, then the analytic function ψ_{μ} has an inverse on some sufficiently small disk around the origin and the *S*-transform of μ is defined by

(2.21)
$$S_{\mu}(z) = \frac{1+z}{z} \psi_{\mu}^{-1}(z).$$

The free unitary normal distribution \mathfrak{N}_{σ^2} with parameter $\sigma^2 > 0$ introduced in [BV92, Lemmas 6.3], is a probability measure on \mathbb{T} with

(2.22)
$$S_{\mathfrak{N}_{\sigma^2}}(z) = e^{\sigma^2 \left(z + \frac{1}{2}\right)}.$$

The next theorem summarizes some properties of the free unitary normal distributions; see Figure 2.2 for the plots of their densities. Almost all of these properties are known from the work of Biane [Bia97a], [Bia97b, Proposition 10]; see also the discussion in [Kem17, Proposition 2.24] and [Lév08, Remark 6.8] for further pointers to the literature. For similar properties of the free unitary Poisson distribution we refer to [Kab21, Section 5].

THEOREM 2.18. — The density of the free unitary normal distribution \mathfrak{N}_{σ^2} w.r.t. the length measure on \mathbb{T} is given by

(2.23)
$$f_{\sigma^2}(e^{i\theta}) = \frac{2}{\pi\sigma^2} \operatorname{Im} \zeta_{\sigma^2/4} \left(\frac{\theta - \pi}{2}\right), \quad \theta \in [-\pi, \pi),$$

where $\zeta_{\sigma^2/4}(\cdot)$ is the function appearing in Theorem 2.3. The distribution \mathfrak{N}_{σ^2} is invariant under complex conjugation meaning that $f_{\sigma^2}(e^{i\theta}) = f_{\sigma^2}(e^{-i\theta})$.

- (a) For $\sigma^2 > 4$, the function $\theta \mapsto f_{\sigma^2}(e^{i\theta})$ is strictly positive and real analytic on \mathbb{R} .
- (b) For $0 < \sigma^2 < 4$, the function $\theta \mapsto f_{\sigma^2}(e^{i\theta})$ is continuous on its period $[-\pi, \pi]$. It is strictly positive and real-analytic on the interval $(-m_{\sigma^2}, +m_{\sigma^2})$, and vanishes on its complement $[-\pi, \pi] \setminus (-m_{\sigma^2}, +m_{\sigma^2})$, where

(2.24)
$$m_{\sigma^2} := 2 \arcsin \frac{\sigma}{2} + \frac{\sigma}{2} \sqrt{4 - \sigma^2} = \int_0^\sigma \sqrt{4 - t^2} \, \mathrm{d}t.$$

At the points $\pm m_{\sigma^2}$, the function $\theta \mapsto f_{\sigma^2}(e^{i\theta})$ vanishes and has square-root singularities with

(2.25)
$$f_{\sigma^2}\left(e^{i\left(m_{\sigma^2}-\varepsilon\right)}\right) = f_{\sigma^2}\left(e^{i\left(\varepsilon-m_{\sigma^2}\right)}\right) \sim \frac{1}{\pi\sigma^2}\sqrt[4]{\frac{4\sigma^2}{4-\sigma^2}} \cdot \sqrt{\varepsilon}, \quad \varepsilon \downarrow 0.$$

(c) For $\sigma^2 = 4$, the function $\theta \mapsto f_{\sigma^2}(e^{i\theta})$ is strictly positive and real-analytic on the interval $(-\pi, \pi)$. At $\theta = \pm \pi$, it vanishes and has cubic-root singularities

with

(2.26)
$$f_{\sigma^2}\left(e^{i(\pi-\varepsilon)}\right) = f_{\sigma^2}\left(e^{i(\varepsilon-\pi)}\right) \sim \frac{\sqrt{3}}{4\pi} \left(\frac{3\varepsilon}{2}\right)^{1/3}, \quad \varepsilon \downarrow 0$$

Proof of Theorem 2.18. — We shall show in Lemma 3.3 below that

$$\psi_{\mathfrak{N}_{\sigma^2}}(-\mathrm{e}^{\mathrm{i}\,\theta}) = \frac{\mathrm{i}\left(\theta - 2\zeta_{\sigma^2/4}(\theta/2)\right)}{\sigma^2} - \frac{1}{2}, \quad \theta \in \mathbb{H}.$$

Consider the Poisson integral of \mathfrak{N}_{σ^2} given for $z \in \mathbb{D}$ by

$$F_{\mathfrak{N}_{\sigma^2}}(z) := \frac{1}{2\pi} \operatorname{Re} \int_{\mathbb{T}} \frac{u+z}{u-z} \,\mathfrak{N}_{\sigma^2}(\mathrm{d}\, u) = \frac{1}{2\pi} \operatorname{Re} \int_{\mathbb{T}} \frac{1+z\bar{u}}{1-z\bar{u}} \,\mathfrak{N}_{\sigma^2}(\mathrm{d}\, u) \\ = \frac{1}{2\pi} \operatorname{Re} \left(1+2\psi_{\mathfrak{N}_{\sigma^2}}(\bar{z})\right).$$

Since the function $\zeta_{\sigma^2/4}$ admits a continuous extension to $\overline{\mathbb{H}}$, the above formula defines $F_{\mathfrak{N}_{\sigma^2}}(z)$ as a continuous function on $\overline{\mathbb{D}}$. Under these circumstances, it is well known (see, e.g., [Kab21, Lemma 5.2]) that the density of \mathfrak{N}_{σ^2} with respect to the length measure on \mathbb{T} is given by

$$f_{\sigma^2}(\mathbf{e}^{\mathbf{i}\,\theta}) = F_{\mathfrak{N}_{\sigma^2}}(\mathbf{e}^{\mathbf{i}\,\theta}) = \frac{1}{2\pi} \operatorname{Re}\left(1 + 2\psi_{\mathfrak{N}_{\sigma^2}}\left(\mathbf{e}^{-\mathbf{i}\,\theta}\right)\right) = \frac{1}{2\pi} \operatorname{Re}\left(1 + 2\psi_{\mathfrak{N}_{\sigma^2}}\left(-\mathbf{e}^{\mathbf{i}(\pi-\theta)}\right)\right)$$
$$= \frac{2}{\pi\sigma^2} \operatorname{Im}\zeta_{\sigma^2/4}\left(\frac{\theta-\pi}{2}\right)$$

for all $\theta \in \mathbb{R}$. All other claims follow from the properties of the function $\zeta_{\sigma^2/4}$ derived in [Kab21, Section 4]. In particular, Equation (2.25) follows from (2.23) and [Kab21, Remark 4.3], while Equation (2.26) is a consequence of [Kab21, Lemma 4.5]. \Box

Remark 2.19. — It follows from (2.23) and the series expansion of $\zeta_{\sigma^2/4}$ given in [Kab21, Theorem 4.11] that

(2.27)
$$f_{\sigma^2}(e^{i\theta}) = \frac{1}{2\pi} + \frac{1}{\pi} \sum_{\ell=1}^{\infty} \frac{1}{\ell} e^{-\ell\sigma^2/2} q_{\ell-1}(-\ell\sigma^2) \cos(\ell\theta),$$

where $q_m(x) = \sum_{j=0}^m \frac{x^j}{j!} \binom{m+1}{j+1}$, $m \in \mathbb{N}_0$, and the series converges uniformly in $\theta \in \mathbb{R}$. Note that (2.27) characterizes the moments of \mathfrak{N}_{σ^2} . In this form, the result be found in [Bia97a, Lemma 1].

Remark 2.20. — The function ζ_t is closely related to the function $\chi(t, z)$ appearing in [Bia97b, p. 273]. A function very similar to m_{σ^2} appears in [RS15, Theorem 5.1].

It follows from (2.27) that, as $\sigma^2 \to \infty$, the density $f_{\sigma^2}(e^{i\theta})$ converges to $1/(2\pi)$ uniformly, which implies that the free unitary normal distribution converges to the uniform distribution weakly on \mathbb{T} . On the other hand, the next result states that, as $\sigma^2 \to 0$, the free unitary normal distribution \mathfrak{N}_{σ^2} can be approximated by a Wigner semicircle distribution supported on a small arc of length $\sim 4\sigma$ centered at 1.

PROPOSITION 2.21. — For all $x \in (-2, 2)$ we have $\lim_{\sigma^2 \downarrow 0} \sigma f_{\sigma^2}(e^{i\sigma x}) = \frac{1}{2\pi}\sqrt{4-x^2}$.

Proof. — We shall only sketch the idea without giving full details. First of all, from (2.24) we have $m_{\sigma^2} \sim 2\sigma$ as $\sigma^2 \downarrow 0$. Let us take x = 0. It has been argued in [Kab21, Section 4] that for all $\tau \ge 0$, we have $\zeta_{\sigma^2/4}(-\frac{\pi}{2} + i\tau) = -\frac{\pi}{2} + i\tilde{y}_{\sigma^2/4}(\tau)$, where $\tilde{y} = \tilde{y}_{\sigma^2/4}(\tau) > 0$ is the unique positive solution of $\tilde{y} - \frac{1}{4}\sigma^2 \operatorname{cotanh} \tilde{y} = \tau$. For $\tau = 0$ the equation takes the form $\tilde{y} = \frac{1}{4}\sigma^2 \operatorname{cotanh} \tilde{y}$ and its unique positive solution satisfies $\tilde{y}_{\sigma^2/4}(0) \sim \frac{\sigma}{2}$ as $\sigma^2 \downarrow 0$. It follows that $\zeta_{\sigma^2/4}(-\frac{\pi}{2}) = -\frac{\pi}{2} + i\frac{\sigma}{2} + o(\sigma)$, and it follows from (2.23) that $\sigma f_{\sigma^2}(1) \rightarrow \frac{1}{\pi}$ as $\sigma^2 \downarrow 0$. For general $x \in (-2, 2)$, Equation (2.23) expresses $f_{\sigma^2}(e^{i\sigma x})$ through the value $\zeta = \zeta_{\sigma^2/4}(\frac{\sigma x - \pi}{2})$ which solves the equation

$$\zeta - \frac{\sigma^2}{4} \tan \zeta = \frac{\sigma x - \pi}{2}.$$

As $\sigma^2 \downarrow 0$, we can search for a solution in the form $\zeta = -\frac{\pi}{2} + c(x)\sigma + o(\sigma)$ with some unknown c(x) satisfying $c(0) = \frac{i}{2}$ by the above analysis of the case x = 0. Inserting this into the equation for ζ and taking $\sigma^2 \downarrow 0$, we obtain $4c(x) + \frac{1}{c(x)} = 2x$. The solution is $c(x) = \frac{1}{4}x + \frac{1}{4}\sqrt{x^2 - 4}$. It follows that $\operatorname{Im} \zeta_{\sigma^2/4}(\frac{\sigma x - \pi}{2}) \sim \frac{\sigma}{4}\sqrt{4 - x^2}$ for $x \in (-2, 2)$, as $\sigma^2 \downarrow 0$. Inserting this into (2.23) completes the argument. \Box

3. Proofs of Theorems 2.2 and 2.4

In this section we prove Theorems 2.2 and 2.4. Throughout the proof, we use the following notational convention: θ denotes a variable ranging in the upper half-plane \mathbb{H} , while $z := -e^{i\theta}$ is a variable ranging in the unit disk \mathbb{D} . Most of the time we shall be occupied with the proof of the following result.

PROPOSITION 3.1. — Fix $\sigma^2 > 0$. If $r = r(\sigma^2) > 0$ is sufficiently small, then locally uniformly in $|z| \leq r$ we have

(3.1)
$$\lim_{n \to \infty} \frac{1}{n} \cdot \frac{H'_n(z; \sigma^2/n)}{H_n(z; \sigma^2/n)} = \frac{-\mathrm{e}^{-\mathrm{i}\,\theta}}{1 + \mathrm{e}^{-2\mathrm{i}\,\zeta_{\sigma^2/4}(\theta/2)}} = -\mathrm{e}^{-\mathrm{i}\,\theta}\left(\frac{\mathrm{i}}{2}\tan\zeta_{\sigma^2/4}(\theta/2) + \frac{1}{2}\right).$$

As we argued in Remark 2.5, the right-hand side of (3.1) can be considered as an analytic function of $z \in \mathbb{D}$ including the value z = 0 (corresponding to $\operatorname{Im} \theta \to +\infty$) where it equals $-e^{-\sigma^2/2}$.

3.1. Proof of Theorems 2.2 and 2.4 assuming Proposition 3.1

First we need to compute the ψ -transforms of the empirical distribution of zeroes of $H_n(z; \sigma^2/n)$ and of the free unitary normal distribution \mathfrak{N}_{σ^2} . This is done in the next two lemmas.

LEMMA 3.2. — The ψ -transform of the probability measure

$$\mu_{n;\,\sigma^2} := \frac{1}{n} \sum_{z \,\in\, \mathbb{T} \,:\, H_n(z;\,\sigma^2/n) = 0} \delta_z$$

is given by

$$\psi_{\mu_{n;\,\sigma^2}}(z) = -\frac{z}{n} \cdot \frac{H'_n\left(z;\sigma^2/n\right)}{H_n\left(z;\sigma^2/n\right)}, \qquad z \in \mathbb{D}.$$

TOME 8 (2025)

Proof. — Using the definition of the ψ -transform given in (2.20) and the identity $\bar{u} = 1/u$ for $u \in \mathbb{T}$, we can write

$$\overline{\psi_{\mu_{n;\sigma^2}}(\bar{z})} = \frac{1}{n} \sum_{\ell=1}^{\infty} z^{\ell} \sum_{\substack{u \in \mathbb{T}: \\ H_n(u;\sigma^2/n) = 0}} \frac{1}{u^{\ell}} = \frac{1}{n} \sum_{\substack{u \in \mathbb{T}: \\ H_n(u;\sigma^2/n) = 0}} \sum_{\ell=1}^{\infty} \frac{z^{\ell}}{u^{\ell}} = \frac{1}{n} \sum_{\substack{u \in \mathbb{T}: \\ H_n(u;\sigma^2/n) = 0}} \frac{z}{u-z}$$
$$= -\frac{z}{n} \cdot \frac{H'_n(z;\sigma^2/n)}{H_n(z;\sigma^2/n)}.$$

To complete the proof, note that the polynomials $H_n(z; \sigma^2/n)$ have real coefficients, implying that its non-real zeroes come in complex-conjugated pairs and hence $\overline{\psi_{\mu_{n;\sigma^2}}(\bar{z})} = \psi_{\mu_{n;\sigma^2}}(z)$.

LEMMA 3.3. — The ψ -transform of the free unitary normal distribution \mathfrak{N}_{σ^2} with parameter $\sigma^2 > 0$ is given by

$$\psi_{\mathfrak{N}_{\sigma^2}}(-\mathrm{e}^{\mathrm{i}\,\theta}) = -\frac{\mathrm{i}}{2}\tan\zeta_{\sigma^2/4}(\theta/2) - \frac{1}{2} = \frac{\mathrm{i}\left(\theta - 2\zeta_{\sigma^2/4}(\theta/2)\right)}{\sigma^2} - \frac{1}{2}, \qquad \theta \in \mathbb{H}.$$

Proof. — By the definition of \mathfrak{N}_{σ^2} given in (2.22), for all complex w with sufficiently small |w| we have

(3.2)
$$w = \psi_{\mathfrak{N}_{\sigma^2}} \left(\frac{w}{1+w} e^{\sigma^2 \left(w + \frac{1}{2}\right)} \right)$$

Let us use the shorthand $\zeta := \zeta_{\sigma^2/4}(\theta/2)$ with some $\theta \in \mathbb{H}$ and put

$$w := -\frac{i}{2}\tan\zeta - \frac{1}{2} = \frac{i(\theta - 2\zeta)}{\sigma^2} - \frac{1}{2}$$

If Im θ is sufficiently large, then |w| is sufficiently small; see (2.4). Also, the definition of w implies that

$$\frac{w}{1+w} e^{\sigma^2 \left(w+\frac{1}{2}\right)} = \frac{-\frac{i}{2} \tan \zeta - \frac{1}{2}}{-\frac{i}{2} \tan \zeta + \frac{1}{2}} e^{i(\theta-2\zeta)} = \frac{-\frac{e^{i\zeta} - e^{-i\zeta}}{e^{i\zeta} + e^{-i\zeta}} - 1}{-\frac{e^{i\zeta} - e^{-i\zeta}}{e^{i\zeta} + e^{-i\zeta}} + 1} e^{i(\theta-2\zeta)}$$
$$= -e^{i\theta}.$$

Inserting this into (3.2) yields $\psi_{\mathfrak{N}_{\sigma^2}}(-e^{i\theta}) = w = -\frac{i}{2} \tan \zeta_{\sigma^2/4}(\theta/2) - \frac{1}{2}$ for all $\theta \in \mathbb{H}$ with sufficiently large Im θ . The latter restriction can be dropped by uniqueness of analytic continuation since both sides are analytic functions of $\theta \in \mathbb{H}$. \Box

Proof of Theorems 2.2 and 2.4 assuming Proposition 3.1. From Proposition 3.1 combined with Lemmas 3.2 and 3.3 we conclude that

(3.3)
$$\psi_{\mu_{n;\sigma^{2}}}(z) = -\frac{z}{n} \cdot \frac{H'_{n}(z;\sigma^{2}/n)}{H_{n}(z;\sigma^{2}/n)} \xrightarrow[n \to \infty]{} -\frac{1}{1 + e^{-2i\zeta_{\sigma^{2}/4}(\theta/2)}} = -\frac{i}{2} \tan \zeta_{\sigma^{2}/4}(\theta/2) - \frac{1}{2} = \psi_{\mathfrak{N}_{\sigma^{2}}}(z),$$

locally uniformly on $\{|z| \leq r\}$. By standard results, see [FHS20, Lemmas 2.6 and 2.11], this already implies uniform convergence on $\{|z| \leq R\}$ for every $R \in (0, 1)$ and the weak convergence $\mu_{n;\sigma^2} \to \mathfrak{N}_{\sigma^2}$. For convenience of the reader, we provide a full proof.

By Cauchy's integral formula, (3.3) implies the convergence of the derivatives of any order at 0 of the above ψ -transforms. By (2.20) this means that the Fourier coefficients of $\mu_{n;\sigma^2}$ converge to those of \mathfrak{N}_{σ^2} , namely

$$\int_{\mathbb{T}} u^{\ell} \mu_{n;\sigma^2}(\mathrm{d}u) = \frac{\psi_{\mu_{n;\sigma^2}}^{(\ell)}(0)}{\ell!} \xrightarrow[n \to \infty]{} \frac{\psi_{\mathfrak{N}_{\sigma^2}}^{(\ell)}(0)}{\ell!} = \int_{\mathbb{T}} u^{\ell} \mathfrak{N}_{\sigma^2}(\mathrm{d}u),$$

for all $\ell \in \mathbb{N}$. Since we are dealing with measures invariant under complex conjugation, this convergence continues to hold for all $\ell \in \mathbb{Z}$. By [Bil68, p. 50], it follows that $\mu_{n;\sigma^2} \to \mathfrak{N}_{\sigma^2}$ weakly on \mathbb{T} , thus proving Theorem 2.2.

To prove Theorem 2.4, we need the following classical consequence of Montel's fundamental normality test; see [Bur79, p. 252] and also [Bur79, p. 219] for a slightly weaker Vitali–Porter theorem.

THEOREM 3.4 (Carathéodory–Landau theorem). — Let f_1, f_2, \ldots and f be analytic functions defined on the unit disk \mathbb{D} and taking values in $\mathbb{C}\setminus\{a_1, a_2\}$, where $a_1, a_2 \in \mathbb{C}$ are distinct. If $\lim_{n\to\infty} f_n(z_i) = f(z_i)$ for all $i \in \mathbb{N}$, where $z_1, z_2, \ldots \in \mathbb{D}$ is a sequence of distinct points having an accumulation point in \mathbb{D} , then $f_n \to f$ locally uniformly on \mathbb{D} .

We are now in position to prove (2.6) of Theorem 2.4. Since $\operatorname{Re} \psi_{\mu_{n;\sigma^2}}(z) > -\frac{1}{2}$ for all $z \in \mathbb{D}$, Theorem 3.4 lifts (3.3) to locally uniform convergence on \mathbb{D} . That is, for every $R \in (0,1)$, we have $\lim_{n\to\infty} \psi_{\mu_{n;\sigma^2}}(z) = \psi_{\mathfrak{N}_{\sigma^2}}(z)$ uniformly on $\{|z| \leq R\}$. In view of Lemmas 3.2 and 3.3 we conclude that

(3.4)
$$\lim_{n \to \infty} \frac{1}{n} \cdot \frac{H'_n(z; \sigma^2/n)}{H_n(z; \sigma^2/n)} = \frac{-e^{-i\theta}}{1 + e^{-2i\zeta_{\sigma^2/4}(\theta/2)}} = -e^{-i\theta} \left(\frac{i}{2} \tan \zeta_{\sigma^2/4}(\theta/2) + \frac{1}{2}\right)$$

uniformly on $\{|z| \leq R\}$. This proves (2.6).

Let us prove (2.5) of Theorem 2.4. Denote the right-hand side of (3.4) by $h_{\sigma^2/4}(z)$ with the usual convention $z = -e^{i\theta}$. Integrating the uniform convergence in (3.4) along the segment joining 0 and z we obtain

(3.5)
$$\frac{\frac{1}{n}\log\frac{H_n(z;\sigma^2/n)}{(-1)^n} = \int_0^z \frac{1}{n} \cdot \frac{H'_n(y;\sigma^2/n)}{H_n(y;\sigma^2/n)} dy}{\underset{n \to \infty}{\longrightarrow} \int_0^z h_{\sigma^2/4}(y) dy = \log\left(1 + e^{2i\zeta_{\sigma^2/4}(\theta/2)}\right) - \frac{\sigma^2}{2\left(1 + e^{-2i\zeta_{\sigma^2/4}(\theta/2)}\right)^2}.$$

To prove the last equality, one observes that the right-hand side of (3.5) vanishes at z = 0 and that its derivative in z is $h_{\sigma^2/4}(z)$; see (3.12) for the details of the calculation.

3.2. Proof of Proposition 3.1

Proof. — We start with an integral representation of the unitary Hermite polynomials. A related result can be found in [SZ18, Proposition 2.1].

LEMMA 3.5. — For all $n \in \mathbb{N}$, $\sigma^2 > 0$ and $z \in \mathbb{C}$ we have

(3.6)
$$H_n(z;\sigma^2/n) = (-1)^n \frac{\sqrt{n}}{\sqrt{2\pi\sigma^2}} \int_{-\infty}^{+\infty} \left(1 - z \,\mathrm{e}^t\right)^n \mathrm{e}^{-\frac{n}{2}\left(\frac{t}{\sigma} + \frac{\sigma}{2}\right)^2} \,\mathrm{d}t$$

Proof. — As in [SZ18], we use the Hubbard–Stratonovich trick, that is the identity $e^{\frac{1}{2}a^2} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} e^{-\frac{1}{2}s^2} e^{as} ds$, for all $a \in \mathbb{C}$. With $a = \sigma(j - \frac{n}{2})/\sqrt{n}$ it follows that

$$e^{\frac{1}{2n}\sigma^2(j-\frac{n}{2})^2} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} e^{-\frac{1}{2}s^2} e^{s\sigma(j-\frac{n}{2})/\sqrt{n}} ds.$$

With this in mind, we can represent the unitary Hermite polynomials as follows:

$$\begin{aligned} H_n\left(z;\sigma^2/n\right) &= \sum_{j=0}^n \binom{n}{j} (-1)^{n-j} z^j \exp\left\{\frac{\sigma^2 \left(j^2 - jn\right)}{2n}\right\} \\ &= \mathrm{e}^{-\frac{1}{8}\sigma^2 n} \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} \sum_{j=0}^n \binom{n}{j} (-1)^{n-j} z^j \, \mathrm{e}^{-\frac{1}{2}s^2} \, \mathrm{e}^{s\sigma\left(j-\frac{n}{2}\right)/\sqrt{n}} \, \mathrm{d}s \\ &= (-1)^n \, \mathrm{e}^{-\frac{1}{8}\sigma^2 n} \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} \left(1 - z \, \mathrm{e}^{s\sigma/\sqrt{n}}\right)^n \, \mathrm{e}^{-\frac{1}{2}s^2} \, \mathrm{e}^{-\frac{1}{2}s\sigma\sqrt{n}} \, \mathrm{d}s \\ &= (-1)^n \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} \left(1 - z \, \mathrm{e}^{s\sigma/\sqrt{n}}\right)^n \, \mathrm{e}^{-\frac{1}{2}\left(s+\frac{1}{2}\sigma\sqrt{n}\right)^2} \, \mathrm{d}s. \end{aligned}$$

After the substitution $t = s\sigma/\sqrt{n}$ we arrive at (3.6).

Our next aim is to prove the following result which will imply Proposition 3.1 by differentiation.

PROPOSITION 3.6. — Fix some $\sigma^2 > 0$. If $r = r(\sigma^2) > 0$ is sufficiently small, then uniformly over the disk $\{z \in \mathbb{C} : |z| \leq r\}$ we have

$$\lim_{n \to \infty} \frac{1}{n} \log \frac{H_n(z; \sigma^2/n)}{(-1)^n} = \log \left(1 + e^{2i\zeta_{\sigma^2/4}(\theta/2)} \right) - \frac{\sigma^2}{2\left(1 + e^{-2i\zeta_{\sigma^2/4}(\theta/2)} \right)^2}$$

where $\theta \in \mathbb{H}$ is such that $-e^{i\theta} = z$ if $z \neq 0$. In the case z = 0, which corresponds to $\operatorname{Im} \theta \to +\infty$, the right-hand side is defined to be 0, by continuity. The branch of the logarithm on the left-hand side is chosen such that $\log 1 = 0$ at z = 0 and the function $\log(\ldots)$ is continuous (and analytic).

We shall write the integral in (3.6) as $\int_{-\infty}^{0} + \int_{0}^{+\infty}$ and analyze both summands separately. As we shall show, the main contribution comes from the negative half-axis.

Saddle point analysis

With the help of the saddle-point method [SFS85, §45.4, p. 423] we shall analyze the integral $\int_{-\infty}^{0} e^{nS(t;z)} dt$, where

(3.7)
$$S(t;z) = \log(1-z\,\mathrm{e}^t) - \frac{1}{2}\left(\frac{t}{\sigma} + \frac{\sigma}{2}\right)^2, \quad |z| < 1, \, \mathrm{Re}\,t < 0.$$



Figure 3.1. Level lines of the function $t \mapsto \operatorname{Re} S(t; z)$ in the left half-plane $\{\operatorname{Re} t \leq 0\}$. Left panel: z = 0. Right panel: $z \neq 0$. In both cases, $\sigma = 2$. The level set passing through the saddle point is shown in black. Level lines where the function takes smaller (respectively, larger) values than at the saddle point are shown in blue (respectively, red).

Observe that S(t; z) is an analytic function of its two variables since for |z| < 1 and Re t < 0 we have $|z e^t| < 1$, implying that $\log(1 - z e^t)$ is well-defined and analytic. The saddle-point equation takes the form

(3.8)
$$\frac{\mathrm{d}}{\mathrm{d}t}S(t;z) = -\frac{z\,\mathrm{e}^t}{1-z\,\mathrm{e}^t} - \frac{t}{\sigma^2} - \frac{1}{2} = 0$$

LEMMA 3.7. — For every $z \in \mathbb{D}$, Equation (3.8) has a unique solution $t_0 := t_0(z; \sigma^2)$ in the left half-plane {Re t < 0}. For $z \in \mathbb{D} \setminus \{0\}$, the solution is given by

(3.9)
$$t_0(z;\sigma^2) = \frac{i\sigma^2}{2} \tan \zeta_{\sigma^2/4}(\theta/2) = 2i \left(\zeta_{\sigma^2/4}(\theta/2) - \theta/2 \right),$$

where we recall the notation $z = -e^{i\theta}$ for some $\theta \in \mathbb{H}$. For z = 0, the solution is $t_0(0; \sigma^2) = -\frac{1}{2}\sigma^2$.

Proof. — Let $z = -e^{i\theta} \neq 0$ with $\theta \in \mathbb{H}$. Every complex number t with $\operatorname{Re} t < 0$ can be represented as $t = \frac{i\sigma^2}{2} \tan \zeta$ with some $\zeta \in \mathbb{H}$ which is unique up to an additive term of the form $\pi n, n \in \mathbb{Z}$. With this notation, Equation (3.8) takes the form

$$0 = \frac{1}{2} + \frac{t}{\sigma^2} + \frac{z \,\mathrm{e}^t}{1 - z \,\mathrm{e}^t} = \frac{1}{2} + \frac{i}{2} \tan \zeta - \frac{1}{1 - z^{-1} \,\mathrm{e}^{-t}} = \frac{1}{1 + \mathrm{e}^{-2\,\mathrm{i}\,\zeta}} - \frac{1}{1 + \mathrm{e}^{-\,\mathrm{i}\,\theta - \frac{\mathrm{i}\,\sigma^2}{2}\,\mathrm{tan}\,\zeta}}.$$

It follows that $2i\zeta = i\theta + \frac{i\sigma^2}{2}\tan\zeta + 2\pi in$ for some $n \in \mathbb{Z}$. Hence, for $\zeta^* := \zeta - \pi n$ we obtain the equation $\zeta^* = \frac{\theta}{2} + \frac{\sigma^2}{4}\tan\zeta^*$. Since $\zeta^* \in \mathbb{H}$, it follows that $\zeta^* = \zeta_{\sigma^2/4}(\theta/2)$, see Theorem 2.3, and the proof is complete.

We are now going to apply the saddle-point method [SFS85, §45.4, p. 423] to the integral $\int_{-\infty}^{0} e^{nS(t;z)} dt$. To this end, we shall replace the initial contour of integration

by a contour γ (depending on z and σ^2) with the following properties. The contour γ starts at $-\infty$, ends at 0, and stays in the left half-plane Re $t \leq 0$. Furthermore, γ passes through the saddle point $t_0 = t_0(z) = t_0(z; \sigma^2)$, and satisfies Re $S(t; z) < \text{Re } S(t_0(z); z)$ for all $t \in \gamma \setminus \{t_0(z)\}$. Finally, in the neighborhood of the point $t_0(z)$, the contour is required to pass through both sectors in which Re $S(t; z) < \text{Re } S(t_0(z); z)$.

We shall prove the existence of such a contour provided |z| is sufficiently small (even though we conjecture that it exists for all $z \in \mathbb{D}$). Let $\partial_1 S$ and $\partial_2 S$ denote the partial derivatives of $S(\cdot, \cdot)$ in the first/second argument. First of all, for z = 0, the function $S(t; 0) = -\frac{1}{2}(\frac{t}{\sigma} + \frac{\sigma}{2})^2$ attains its strict maximum on $(-\infty, 0]$ at $t_0(0) = -\frac{1}{2}\sigma^2$. This critical point is simple meaning that $(\partial_1^2 S)(-\frac{1}{2}\sigma^2; 0) < 0$. It follows that the negative half-line $(-\infty, 0]$ satisfies the conditions listed above; see the left panel of Figure 3.1. Observe also that in a small disk D centered at $-\frac{1}{2}\sigma^2$, the set where Re $S(t; 0) < \text{Re } S(-\frac{1}{2}\sigma^2; 0)$ consists of two sectors whose boundaries are straight lines crossing D at four points A_1, A_2, A_3, A_4 . After a small perturbation of z, we still have a critical point at $t_0(z)$ which is close to $t_0(0)$ and simple, by analyticity of $t_0(z)$ and S; see Lemma 3.7. It follows that, in the disk D, the set where Re $S(t; z) < \text{Re } S(t_0(z); z)$ still consists of two sectors; see the right panel of Figure 3.1. The four points where the boundaries of these sectors cross D are close to the points A_1, A_2, A_3, A_4 , again by analyticity. It follows that the saddle-point contour γ satisfying the above condition exists (and can be obtained by perturbing the segment of the real line contained in D).

Applying the saddle-point asymptotics [SFS85, §45.4, p. 423], we have

(3.10)
$$\int_{-\infty}^{0} e^{nS(t;z)} dt \sim \sqrt{-\frac{2\pi}{n(\partial_1^2 S)(t_0(z);z)}} e^{nS(t_0(z);z)}, \quad \text{as } n \to \infty$$

This holds pointwise in z provided |z| is sufficiently small (the uniformity in z will be addressed later). A formula for $(\partial_1^2 S)(t_0(z); z)$ will be given in (3.17). The choice of the square root branch, irrelevant for our purposes, is explained in [SFS85, p. 425]. It follows from (3.7) and (3.9) that for $z = -e^{i\theta} \neq 0$ we have

(3.11)
$$S(t_0(z); z) = \log\left(1 - z e^{t_0(z)}\right) - \frac{1}{2} \left(\frac{t_0(z)}{\sigma} + \frac{\sigma}{2}\right)^2 = \log\left(1 + e^{2i\zeta_{\sigma^2/4}(\theta/2)}\right) - \frac{\sigma^2}{2\left(1 + e^{-2i\zeta_{\sigma^2/4}(\theta/2)}\right)^2}.$$

Note that for z = 0 we have $S(t_0(0); 0) = 0$. For future use let us note the identity

(3.12)

$$\frac{\mathrm{d}}{\mathrm{d}z}S(t_0(z);z) = (\partial_1 S)(t_0(z);z)\partial_z t_0(z) + (\partial_2 S)(t_0(z);z) = (\partial_2 S)(t_0(z);z) \\
= -\frac{\mathrm{e}^{t_0(z)}}{1-z\,\mathrm{e}^{t_0(z)}} = \frac{1}{z}\left(\frac{1}{2} + \frac{t_0(z)}{\sigma^2}\right) \\
= -\mathrm{e}^{-\mathrm{i}\theta}\left(\frac{\mathrm{i}}{2}\tan\zeta_{\sigma^2/4}(\theta/2) + \frac{1}{2}\right) = \frac{-\mathrm{e}^{-\mathrm{i}\theta}}{1+\mathrm{e}^{-2\mathrm{i}\zeta_{\sigma^2/4}(\theta/2)}},$$

where we used that $(\partial_1 S)(t_0(z); z) = 0$ since $t = t_0(z)$ solves the saddle-point equation (3.8).

Contribution of the positive half-axis is negligible

We claim that if $r = r(\sigma^2) > 0$ is sufficiently small, then uniformly over the disk $|z| \leq r$ it holds that

(3.13)
$$\int_0^\infty \left(1 - z \operatorname{e}^t\right)^n \operatorname{e}^{-\frac{n}{2}\left(\frac{t}{\sigma} + \frac{\sigma}{2}\right)^2} \mathrm{d}t = O\left(\operatorname{e}^{-\frac{1}{16}\sigma^2 n}\right), \quad \text{as } n \to \infty$$

If $0 \leq t \leq \sigma^2$, then by choosing a sufficiently small r > 0 we can achieve that $|1 - z e^t| \leq 1 + |z| e^t < e^{\frac{1}{16}\sigma^2}$ for all $|z| \leq r$ and hence, for all $n \in \mathbb{N}$,

$$\left| \left(1 - z \, \mathrm{e}^t \right)^n \mathrm{e}^{-\frac{n}{2} \left(\frac{t}{\sigma} + \frac{\sigma}{2} \right)^2} \right| \leq \mathrm{e}^{\frac{1}{16} \sigma^2 n} \, \mathrm{e}^{-\frac{1}{8} \sigma^2 n} = \mathrm{e}^{-\frac{1}{16} \sigma^2 n} \, .$$

It follows that

(3.14)
$$\int_{0}^{\sigma^{2}} \left(1 - z e^{t}\right)^{n} e^{-\frac{n}{2} \left(\frac{t}{\sigma} + \frac{\sigma}{2}\right)^{2}} dt = O\left(-e^{\frac{1}{16}\sigma^{2}n}\right), \quad \text{as } n \to \infty.$$

For $t \ge \sigma^2$ we argue as follows. Since $1 \le e^t$ for all $t \ge 0$, we have $|1 - z e^t| \le 1 + |z| e^t \le e^t (1 + r)$ provided $|z| \le r$. It follows that

$$\left| \int_{\sigma^2}^{\infty} \left(1 - z \, \mathrm{e}^t \right)^n \mathrm{e}^{-\frac{n}{2} \left(\frac{t}{\sigma} + \frac{\sigma}{2} \right)^2} \, \mathrm{d}t \right| \leq (1 + r)^n \int_{\sigma^2}^{\infty} \mathrm{e}^{-\frac{n}{2} \left(\frac{t}{\sigma} - \frac{\sigma}{2} \right)^2} \, \mathrm{d}t$$
$$= (1 + r)^n \, \mathrm{e}^{-\frac{1}{8}\sigma^2(n-1)} \int_{\sigma^2}^{\infty} \mathrm{e}^{-\frac{1}{2} \left(\frac{t}{\sigma} - \frac{\sigma}{2} \right)^2} \, \mathrm{d}t$$

since $\frac{1}{2}(\frac{t}{\sigma} - \frac{\sigma}{2})^2 \ge \frac{1}{8}\sigma^2$ for $t \ge \sigma^2$. Note that the integral on the right-hand side converges. Also, if r > 0 is sufficiently small, then $1 + r \le e^{\frac{1}{16}\sigma^2}$ and it follows that (3.15) $\int_{\sigma^2}^{\infty} (1 - z e^t)^n e^{-\frac{n}{2}(\frac{t}{\sigma} + \frac{\sigma}{2})^2} dt = O\left(e^{-\frac{1}{16}\sigma^2 n}\right), \quad \text{as } n \to \infty.$

Combining (3.14) and (3.15) yields (3.13), thus completing the proof that the contribution of the positive half-line is negligible.

Exact asymptotics of H_n

It follows from (3.10) and (3.13), together with the fact that $S(t_0(z); z)$ converges to 0 as $z \to 0$, that, for sufficiently small |z|, the contribution of the positive half-axis is negligible in the sense that

$$\int_{-\infty}^{\infty} e^{nS(t;z)} dt \sim \sqrt{-\frac{2\pi}{n(\partial_1^2 S)(t_0(z);z)}} e^{nS(t_0(z);z)}, \quad \text{as } n \to \infty$$

In view of Lemma 3.5, we can write this as

(3.16)
$$\frac{H_n(z;\sigma^2/n)}{(-1)^n} \sim \sqrt{-\frac{1}{\sigma^2(\partial_1^2 S)(t_0(z);z)}} e^{nS(t_0(z);z)} \quad \text{as } n \to \infty$$

This holds pointwise for every $|z| \leq r$ provided that r > 0 is sufficiently small. Although we shall not need this fact, let us mention that after some work it is possible to verify that

(3.17)
$$\left(\partial_1^2 S\right) \left(t_0(z); z\right) = \frac{z^{-1} e^{-t_0(z)}}{\left(1 - z^{-1} e^{-t_0(z)}\right)^2} - \frac{1}{\sigma^2} = -\frac{1}{4 \cos^2 \zeta_{\sigma^2/4}(\theta/2)} - \frac{1}{\sigma^2}.$$

TOME 8 (2025)

Proof of the uniform convergence

Now we would like to take the logarithm of both sides of (3.16). Observe that the left-hand side cannot become zero for $z \in \mathbb{D}$ by Lemma 2.1. Trivially, the same conclusion holds for the right-hand side. Therefore, applying the function $w \mapsto \log |w|$ to both sides of (3.16) and dividing by n we get

(3.18)
$$\lim_{n \to \infty} \frac{1}{n} \log \left| \frac{H_n(z; \sigma^2/n)}{(-1)^n} \right| = \operatorname{Re} S(t_0(z); z).$$

This holds pointwise in z provided $|z| \leq r$ with r > 0 sufficiently small. Note that we did not apply the function $w \mapsto \log w$ to avoid difficulties with the choice of the branch. Now, our aim is to prove that

(3.19)
$$\lim_{n \to \infty} \frac{1}{n} \log \frac{H_n(z; \sigma^2/n)}{(-1)^n} = S(t_0(z); z)$$

locally uniformly in |z| < r and with the same convention for the logarithm as in Proposition 3.6. To this end, we shall use the following lemma which strengthens [Kab21, Lemma 3.8].

LEMMA 3.8. — Let $h_1(z), h_2(z), \ldots$ be a sequence of holomorphic functions defined on some domain $\mathcal{D} \subset \mathbb{C}$. If $\operatorname{Re} h_n(z) \to 0$ pointwise on \mathcal{D} and the sequence $(\operatorname{Re} h_n(z))_{n \in \mathbb{N}}$ is locally uniformly bounded from above on \mathcal{D} , then $h'_n(z) \to 0$ locally uniformly on \mathcal{D} .

Remark 3.9. — If, additionally, $h_n(y_0) \to 0$ for some $y_0 \in \mathcal{D}$, then, integrating, we conclude that $h_n(z) \to 0$ locally uniformly on \mathcal{D} .

Proof of Lemma 3.8. — Consider some closed disk B contained in \mathcal{D} and centered at $z_0 \in \mathcal{D}$. We know that $\operatorname{Re} h_n(z) \leq C$, for all $n \in \mathbb{N}$ and $z \in B$. Also, we know that $\operatorname{Re} f_n(z_0)$ converges (and hence is bounded below). By Harnack's inequality applied to the non-negative harmonic functions $C - \operatorname{Re} h_n(z)$ it follows that $C - \operatorname{Re} h_n(z)$ is uniformly bounded from above on B. We conclude that the sequence $(\operatorname{Re} h_n(z))_{n \in \mathbb{N}}$ is locally uniformly bounded, both from above and from below.

Take some $z_0 \in \mathcal{D}$ and let $B_r(z_0) \subset \mathcal{D}$ be a closed disk of radius r centered at z_0 and contained in \mathcal{D} . It is known (see, e.g., the proof of [Kab21, Lemma 3.8]) that for all z in the interior of $B_r(z_0)$, we have

(3.20)
$$h'_{n}(z) = \frac{1}{\pi i} \oint_{|w-z_{0}|=r} \frac{\operatorname{Re} h_{n}(w)}{(w-z)^{2}} \mathrm{d}w,$$

where the integration contour is the boundary of $B_r(z_0)$, oriented counter-clockwise. Having (3.20) at our disposal, we claim that $h'_n(z) \to 0$ uniformly over $z \in B_{r/2}(z_0)$. Indeed, (3.20) implies

$$\sup_{z \in B_{r/2}(z_0)} |h'_n(z)| \leq \frac{1}{\pi} \cdot \left(\sup_{\substack{|w-z_0|=r\\z \in B_{r/2}(z_0)}} \frac{1}{|w-z|^2} \right) \cdot \int_{|w-z_0|=r} |\operatorname{Re} h_n(w)| |\mathrm{d}w|$$

Since the supremum is finite and the integral converges to 0 by the dominated convergence theorem, the desired conclusion follows. $\hfill \Box$

We apply Lemma 3.8 with $h_n(z) := \frac{1}{n} \log \frac{H_n(z; \sigma^2/n)}{(-1)^n} - S(t_0(z); z)$. These functions are analytic on $\{|z| < r\}$ and we have $h_n(0) = 0$ as well as $\operatorname{Re} h_n(z) \to 0$ pointwise, by (3.18). Observe that the functions $(\operatorname{Re} h_n(z))_{n \in \mathbb{N}}$ are locally uniformly bounded from above since by (1.5) and the triangle inequality,

$$\frac{1}{n}\log\left|\frac{H_n\left(z;\sigma^2/n\right)}{(-1)^n}\right| \leqslant \frac{1}{n}\log\left(\sum_{j=0}^n \binom{n}{j}|z|^j\right) \leqslant \log(1+|z|), \quad z \in \mathbb{C}.$$

Lemma 3.8 (together with Remark 3.9) yields (3.19). Taking into account (3.11), this proves Proposition 3.6. Since a locally uniform convergence of analytic functions can be differentiated, we infer that

$$\lim_{n \to \infty} \frac{1}{n} \cdot \frac{H'_n(z; \sigma^2/n)}{H_n(z; \sigma^2/n)} = \frac{\mathrm{d}}{\mathrm{d}z} S(t_0(z); z) = \frac{-\mathrm{e}^{-\mathrm{i}\,\theta}}{1 + \mathrm{e}^{-2\mathrm{i}\,\zeta_{\sigma^2/4}(\theta/2)}} \\ = -\mathrm{e}^{-\mathrm{i}\,\theta} \left(\frac{\mathrm{i}}{2} \tan\zeta_{\sigma^2/4}(\theta/2) + \frac{1}{2}\right);$$

see (3.12). The proof of Proposition 3.1 is complete.

Acknowledgements

The author is grateful to two anonymous referees for several insightful comments and to Octavio Arizmendi, Jorge Garza-Vargas, Jonas Jalowy, Matthias Löwe and Daniel Perales for useful discussions and, in particular, for pointing out the works of Mirabelli [Mir21] and Hall and Ho [HH22].

BIBLIOGRAPHY

- [ABH04] Alexandru Aleman, Ditry Beliaev, and Haakan Hedenmalm, Real zero polynomials and Pólya–Schur type theorems, J. Anal. Math. 94 (2004), 49–60. ↑7, 15, 17
- [AFU24] Octavio Arizmendi, Katsunori Fujie, and Yuki Ueda, New combinatorial identity for the set of partitions and limit theorems in finite free probability theory, Int. Math. Res. Not. 2024 (2024), no. 14, 10524–10558. ↑7
- [AGN21] Gernot Akemann, Friedrich Götze, and Thorsten Neuschel, Characteristic polynomials of products of non-Hermitian Wigner matrices: finite-N results and Lyapunov universality, Electron. Commun. Probab. 26 (2021), article no. 30. ↑10
- [AGVP23] Octavio Arizmendi, Jorge Garza-Vargas, and Daniel Perales, Finite free cumulants: multiplicative convolutions, genus expansion and infinitesimal distributions, Trans. Am. Math. Soc. 376 (2023), no. 6, 4383–4420. ↑4, 14, 17
- [AGZ10] Greg W. Anderson, Alice Guionnet, and Ofer Zeitouni, An introduction to random matrices, Cambridge Studies in Advanced Mathematics, vol. 118, Cambridge University Press, 2010. ↑16
- [AKM12] Sergio Andraus, Makoto Katori, and Seiji Miyashita, Interacting particles on the line and Dunkl intertwining operator of type A: application to the freezing regime, J. Phys. A. Math. Theor. 45 (2012), no. 39, article no. 395201. [↑]16
- [Aom87] Kazuhiko Aomoto, Jacobi polynomials associated with Selberg integrals, SIAM J. Math. Anal. 18 (1987), no. 2, 545–549. ↑10

- [AP18] Octavio Arizmendi and Daniel Perales, Cumulants for finite free convolution, J. Comb. Theory, Ser. A 155 (2018), 244–266. ↑4, 15
- [Ben34] Eduard Benz, Über lineare, verschiebungstreue Funktionaloperationen und die Nullstellen ganzer Funktionen, Comment. Math. Helv. 7 (1934), no. 1, 243–289. ↑15
- [BH00] Edouard Brézin and Shinobu Hikami, Characteristic polynomials of random matrices, Commun. Math. Phys. 214 (2000), no. 1, 111–135. ↑10
- [BHS24] Rikard Bøgvad, Christian Hägg, and Boris Shapiro, Rodrigues' descendants of a polynomial and Boutroux curves, Constr. Approx. 59 (2024), no. 3, 737–798. ↑14
- [Bia97a] Philippe Biane, Free Brownian motion, free stochastic calculus and random matrices, Free probability theory (Waterloo, ON, 1995), Fields Institute Communications, vol. 12, American Mathematical Society, 1997, pp. 1–19. ↑14, 18, 19, 20
- [Bia97b] _____, Segal-Bargmann transform, functional calculus on matrix spaces and the theory of semi-circular and circular systems, J. Funct. Anal. 144 (1997), no. 1, 232– 286. ↑14, 18, 19, 20
- [Bia09] _____, Matrix valued Brownian motion and a paper by Pólya, Séminaire de Probabilités XLII, Lecture Notes in Mathematics, vol. 1979, Springer, 2009, pp. 171–185. ↑10
- [Bil68] Patrick P. Billingsley, Convergence of probability measures, John Wiley & Sons, 1968. ↑23
- [BLR17] Pavel Bleher, Mikhail Lyubich, and Roland K. W. Roeder, Lee-Yang zeros for the DHL and 2D rational dynamics. I. Foliation of the physical cylinder, J. Math. Pures Appl. (9) 107 (2017), no. 5, 491–590. ^{↑5}
- [BLR20] _____, Lee-Yang-Fisher zeros for the DHL and 2D rational dynamic. II. Global pluripotential interpretation, J. Geom. Anal. **30** (2020), no. 1, 777–833. ↑5
- [Bui24] Wouter Buijsman, Efficient circular Dyson Brownian motion algorithm, Phys. Rev. Res. 6 (2024), no. 2, article no. 023264. ↑10
- [Bur79] Robert B. Burckel, An introduction to classical complex analysis. Vol. 1, Pure and Applied Mathematics, vol. 82, Academic Press Inc., 1979. ↑23
- [BV92] Hari Bercovici and Dan-Virgil Voiculescu, Lévy-Hinčin type theorems for multiplicative and additive free convolution, Pac. J. Math. 153 (1992), no. 2, 217–248. ↑18, 19
- [CDG01] Thierry Cabanal Duvillard and Alice Guionnet, Large deviations upper bounds for the laws of matrix-valued processes and non-communicative entropies, Ann. Probab. 29 (2001), no. 3, 1205–1261. ↑14
- [Céb16] Guillaume Cébron, Matricial model for the free multiplicative convolution, Ann. Probab.
 44 (2016), no. 4, 2427–2478. ↑18
- [CHJR19] Ivan Chio, Caleb He, Anthony L. Ji, and Roland K. W. Roeder, Limiting measure of Lee-Yang zeros for the Cayley tree, Commun. Math. Phys. 370 (2019), no. 3, 925–957. ↑5
- [CL01] Emmanuel Cépa and Dominique Lépingle, Brownian particles with electrostatic repulsion on the circle: Dyson's model for unitary random matrices revisited, ESAIM, Probab. Stat. 5 (2001), 203–224. [↑]10, 17
- [DBF20] Aydin Deger, Fredrik Brange, and Christian Flindt, Lee-Yang theory, high cumulants, and large-deviation statistics of the magnetization in the Ising model, Phys. Rev. B 102 (2020), no. 17, article no. 174418. ↑5
- [DE02] Ioana Dumitriu and Alan Edelman, Matrix models for beta ensembles, J. Math. Phys. 43 (2002), no. 11, 5830–5847. ↑10
- [DE05] _____, Eigenvalues of Hermite and Laguerre ensembles: large beta asymptotics, Ann. Inst. Henri Poincaré, Probab. Stat. 41 (2005), no. 6, 1083–1099. ↑16

- [DF20] Aydin Deger and Christian Flindt, Lee-Yang theory of the Curie-Weiss model and its rare fluctuations, Phys. Rev. Res. 2 (2020), no. 3, article no. 033009. $\uparrow 5$
- [DG04] Persi Diaconis and Alex Gamburd, Random matrices, magic squares and matching polynomials, Electron. J. Comb. 11 (2004), no. 2, article no. 2. ↑10
- [DM24] Tomas Dominguez and Jean-Christophe Mourrat, Statistical Mechanics of Mean-Field Disordered Systems: A Hamilton-Jacobi Approach, EMS Zurich Lectures in Advanced Mathematics Series, European Mathematical Society, 2024. ↑5
- [DMTW19] Radu Dascaliuc, Nicholas Michalowski, Enrique Thomann, and Edward C. Waymire, Complex Burgers equation: A probabilistic perspective, Sojourns in Probability Theory and Statistical Physics. I. Spin Glasses and Statistical Mechanics, A Festschrift for Charles M. Newman, Springer Proceedings in Mathematics & Statistics, vol. 298, Springer, 2019, pp. 138–170. ↑5
- [Dys62] Freeman J. Dyson, A Brownian-motion model for the eigenvalues of a random matrix,
 J. Math. Phys. 3 (1962), 1191–1198. ↑10
- [Ede89] Alan Edelman, Eigenvalues and condition numbers of random matrices, Ph.D. thesis, Massachusetts Institute of Technology, Cambridge, USA, 1989, https://dspace.mit. edu/handle/1721.1/14322. ↑10
- [Ell06] Richard S. Ellis, *Entropy, large deviations, and statistical mechanics*, Classics in Mathematics, Springer, 2006, reprint of the 1985 original. ↑5
- [EN78a] Richard S. Ellis and Charles M. Newman, Limit theorems for sums of dependent random variables occurring in statistical mechanics, Z. Wahrscheinlichkeitstheor. Verw. Geb. 44 (1978), no. 2, 117–139. ↑5
- [EN78b] _____, The statistics of Curie–Weiss models, J. Stat. Phys. 19 (1978), no. 2, 149–161. $\uparrow 5$
- [ENR80] Richard S. Ellis, Charles M. Newman, and Jay S. Rosen, Limit theorems for sums of dependent random variables occurring in statistical mechanics. II. Conditioning, multiple phases, and metastability, Z. Wahrscheinlichkeitstheor. Verw. Geb. 51 (1980), no. 2, 153–169. ↑5
- [FG06] Peter J. Forrester and Alex Gamburd, Counting formulas associated with some random matrix averages, J. Comb. Theory, Ser. A 113 (2006), no. 6, 934–951. ↑10
- [FHS20] Uwe Franz, Takahiro Hasebe, and Sebastian Schleißinger, Monotone increment processes, classical Markov processes, and Loewner chains, Diss. Math. 552 (2020), 1–119. ↑19, 22
- [Fis65] Michael E. Fisher, The nature of critical points, Statistical Physics, Weak interactions, Field Theory (W. E. Brittin, ed.), Lecture Notes in Theoretical Physics, vol. 7C, University of Colorado Press, 1965, pp. 1–159. ↑5
- [FKLZ24] Peter J. Forrester, Mario Kieburg, Shi-Hao Li, and Jiyuan Zhang, Dip-ramp-plateau for Dyson Brownian motion from the identity on U(N), Probab. Math. Phys. 5 (2024), no. 2, 321–355. ↑10
- [FV18] Sacha Friedli and Yvan Velenik, Statistical mechanics of lattice systems. A concrete mathematical introduction, Cambridge University Press, 2018. ↑5
- [Gal22] Andre Galligo, Modeling complex root motion of real random polynomials under differentiation, ISSAC'22 — Proceedings of the 2022 International Symposium on Symbolic and Algebraic Computation, ACM Press, 2022, pp. 245–253. ↑14
- [Gaw87] Wolfgang Gawronski, On the asymptotic distribution of the zeros of Hermite, Laguerre, and Jonquière polynomials, J. Approx. Theory 50 (1987), no. 3, 214–231. ↑7, 16
- [GK24] Vadim Gorin and Victor Kleptsyn, Universal objects of the infinite beta random matrix theory, J. Eur. Math. Soc. 26 (2024), no. 9, 3429–3496. ↑16

[GPS86]	M. Lawrence Glasser, Vladimir Privman, and Lawrence S. Schulman, Complex tem-
	perature plane zeros in the mean-field approximation, J. Stat. Phys. 45 (1986), no. 3-4,
	451–457. ↑5

- [Hal18] Brian C. Hall, The eigenvalue process for Brownian motion in U(N), Preprint at https://www3.nd.edu/~bhall/publications, 2018. $\uparrow 10$
- [HH22] Brian C. Hall and Ching-Wei Ho, *The heat flow conjecture for random matrices*, 2022, https://arxiv.org/abs/2202.09660v1. ↑4, 14, 29
- [HJN23] Qi Hou, Jianping Jiang, and Charles M. Newman, Motion of Lee-Yang Zeros, J. Stat. Phys. 190 (2023), no. 3, article no. 56. ↑5
- [HK23] Jeremy Hoskins and Zakhar Kabluchko, Dynamics of zeroes under repeated differentiation, Exp. Math. 32 (2023), no. 4, 573–599. [↑]14
- [HW96] David G. Hobson and Wendelin Werner, Non-colliding Brownian motions on the circle, Bull. Lond. Math. Soc. 28 (1996), no. 6, 643–650. ↑10, 17
- [Kab21] Zakhar Kabluchko, Repeated differentiation and free unitary Poisson process, 2021, https://arxiv.org/abs/2112.14729v1. ↑8, 14, 17, 19, 20, 21, 28
- [KBHK16] Mariana Krasnytska, Bertrand Berche, Yurij Holovatch, and Ralph Kenna, Partition function zeros for the Ising model on complete graphs and on annealed scale-free networks, J. Phys. A. Math. Theor. 49 (2016), no. 13, article no. 135001. ↑5
- [Kem17] Todd Kemp, *Heat kernel empirical laws on* \mathbb{U}_N and \mathbb{GL}_N , J. Theor. Probab. **30** (2017), no. 2, 397–451. $\uparrow 19$
- [KM16] Miklós Kornyik and György Michaletzky, Wigner matrices, the moments of roots of Hermite polynomials and the semicircle law, J. Approx. Theory 211 (2016), 29–41. ↑7, 16
- [KT20] Alexander Kiselev and Changhui Tan, The Flow of Polynomial Roots Under Differentiation, 2020, https://arxiv.org/abs/2012.09080. ↑14
- [Lév08] Thierry Lévy, Schur-Weyl duality and the heat kernel measure on the unitary group, Adv. Math. 218 (2008), no. 2, 537–575. ↑14, 19
- [LY52] Tsung-Dao Lee and Chen-Ning Yang, Statistical theory of equations of state and phase transitions. II. Lattice gas and Ising model, Phys. Rev., II. Ser. 87 (1952), 410–419. ↑5, 7
- [Mar18] Adam W. Marcus, Polynomial convolutions and (finite) free probability, 2018, https: //arxiv.org/abs/2108.07054. ⁴, 15, 17
- [Mir21] Benjamin Mirabelli, Hermitian, Non-Hermitian and Multivariate Finite Free Probability, Ph.D. thesis, Princeton University, Princeton, USA, 2021, https://dataspace. princeton.edu/handle/88435/dsp019593tz21r. ↑4, 5, 29
- [MN15] Pierre-Loïc Méliot and Ashkan Nikeghbali, Mod-Gaussian convergence and its applications for models of statistical mechanics, In memoriam Marc Yor — Séminaire de Probabilités XLVII, Lecture Notes in Mathematics, vol. 2137, Springer, 2015, pp. 369– 425. ↑5
- [MSS22] Adam W. Marcus, Daniel A. Spielman, and Nikhil Srivastava, Finite free convolutions of polynomials, Probab. Theory Relat. Fields 182 (2022), no. 3-4, 807–848. ⁴, 13, 15, 16, 17
- [New86] Charles M. Newman, Shock waves and mean field bounds: Concavity and analyticity of the magnetization at low temperature. Appendix to Percolation theory: A selective survey of rigorous results, Advances in multiphase flow and related problems (George C. Papanicolauo, ed.), Society for Industrial and Applied Mathematics, 1986, pp. 147–167. ↑5

- [NS06] Alexandru Nica and Roland Speicher, Lectures on the combinatorics of free probability, London Mathematical Society Lecture Note Series, vol. 335, Cambridge University Press, 2006. ↑4
- [OS21] Sean O'Rourke and Stefan Steinerberger, A Nonlocal Transport Equation Modeling Complex Roots of Polynomials under Differentiation, Proc. Am. Math. Soc. 149 (2021), 1581–1592. ↑14
- [PB21] Marc Potters and Jean-Philippe Bouchaud, A first course in random matrix theory: for physicists, engineers and data scientists, Cambridge University Press, 2021. ↑6, 10
- [RŚ15] Dan Romik and Piotr Śniady, Jeu de taquin dynamics on infinite Young tableaux and second class particles, Ann. Probab. 43 (2015), no. 2, 682–737. ↑20
- [Rue69] David Ruelle, Statistical mechanics. Rigorous results, The Mathematical Physics Monographs Series, W. A. Benjamin, Inc., 1969. ↑7
- [RY99] Daniel Revuz and Marc Yor, Continuous martingales and Brownian motion, 3rd ed., Grundlehren der Mathematischen Wissenschaften, vol. 293, Springer, 1999. ↑11
- [SFS85] Yu. V. Sidorov, Mikhai V. Fedoryuk, and M. I. Shabunin, Lectures on the theory of functions of a complex variable, Mir Publishers, 1985. ↑24, 25, 26
- [Ste19] Stefan Steinerberger, A nonlocal transport equation describing roots of polynomials under differentiation, Proc. Am. Math. Soc. **147** (2019), no. 11, 4733–4744. ↑14
- [Ste20] _____, Conservation Laws for the Density of Roots of Polynomials under Differentiation, 2020, https://arxiv.org/abs/2001.09967. ↑14
- [Ste23] _____, Free convolution powers via roots of polynomials, Exp. Math. **32** (2023), no. 4, 567-572. $\uparrow 14$
- [SZ18] Mira Shamis and Ofer Zeitouni, The Curie–Weiss model with complex temperature: phase transitions, J. Stat. Phys. 172 (2018), no. 2, 569–591. ↑5, 23, 24
- [Sze75] Gabor Szegő, Orthogonal polynomials, 4th ed., Colloquium Publications, vol. 23, American Mathematical Society, 1975. ↑9, 10
- [Tao17] Terence Tao, Heat flow and zeroes of polynomials, 2017, https://terrytao. wordpress.com/2017/10/17/heat-flow-and-zeroes-of-polynomials/. ¹⁴, 15, 16
- [Tao18] _____, Heat flow and zeroes of polynomials II: zeroes on a circle, 2018, heat-flow-and-zeroes-of-polynomials-ii-zeroes-on-a-circle/. ↑4, 14, 17
- [Ull80] Joseph L. Ullman, Orthogonal polynomials associated with an infinite interval, Mich. Math. J. 27 (1980), no. 3, 353–363. ↑7, 16
- [VDN92] Dan-Virgil Voiculescu, Kenneth J. Dykema, and Alexandru Nica, Free random variables, CRM Monograph Series, vol. 1, American Mathematical Society, 1992. ↑4, 18
- [Voi85] Dan-Virgil Voiculescu, Symmetries of some reduced free product C^{*}-algebras, Operator algebras and their connections with topology and ergodic theory (Buşteni, 1983), Lecture Notes in Mathematics, vol. 1132, Springer, 1985, pp. 556–588. ↑18
- [Voi87] _____, Multiplication of certain noncommuting random variables, J. Oper. Theory 18 (1987), no. 2, 223–235. ↑18
- [VW20] Michael Voit and Jeannette H. C. Woerner, Functional central limit theorems for multivariate Bessel processes in the freezing regime, Stochastic Anal. Appl. 39 (2020), no. 1, 136–156. ↑16
- [VW22] _____, Limit theorems for Bessel and Dunkl processes of large dimensions and free convolutions, Stochastic Processes Appl. 143 (2022), 207–253. ↑16
- [YL52] Chen-Ning Yang and Tsung-Dao Lee, Statistical theory of equations of state and phase transitions. I. Theory of condensation, Phys. Rev., II. Ser. 87 (1952), 404–409. ↑5

- [Zho14] Ping Zhong, Free Brownian motion and free convolution semigroups: multiplicative case, Pac. J. Math. **269** (2014), no. 1, 219–256. ↑18
- [Zho15] _____, On the free convolution with a free multiplicative analogue of the normal distribution, J. Theor. Probab. **28** (2015), no. 4, 1354–1379. ↑18

Manuscript received on 2nd April 2023, revised on 9th September 2024, accepted on 18th October 2024.

Recommended by Editors S. Gouëzel and M. Bousquet-Mélou. Published under license CC BY 4.0.



eISSN: 2644-9463 This journal is a member of Centre Mersenne.



Zakhar KABLUCHKO Institut für Mathematische Stochastik, Universität Münster, Orléans-Ring 10, 48149 Münster (Germany) zakhar.kabluchko@uni-muenster.de