Annales Henri Lebesgue 8 (2025) 67-100



## SIMON ALLAIS PIERRE-ALEXANDRE ARLOVE SHEILA SANDON

# SPECTRAL SELECTORS ON LENS SPACES AND APPLICATIONS TO THE GEOMETRY OF THE GROUP OF CONTACTOMORPHISMS

INVARIANTS SPECTRAUX SUR LES ESPACES LENTICULAIRES ET APPLICATIONS À LA GÉOMÉTRIE DU GROUPE DES CONTACTOMORPHISMES

*Keywords:* Contactomorphism group, spectral selectors, conjugation invariant norms, non-linear Maslov index, generating functions.

<sup>2020</sup> Mathematics Subject Classification: 53D10, 53D35, 57R17.

DOI: https://doi.org/10.5802/ahl.229

<sup>(\*)</sup> The first author was funded by the postdoctoral fellowships of the Fondation Sciences Mathématiques de Paris and of the Interdisciplinary Thematic Institute IRMIA++, as part of the ITI 2021-2028 program of the University of Strasbourg, CNRS and Inserm, supported by IdEx Unistra (ANR-10-IDEX-0002) and by SFRI-STRAT'US project (ANR-20-SFRI-0012) under the framework of the French Investments for the Future Program. The second author is partially supported by the Deutsche Forschungsgemeinschaft under the Collaborative Research Center SFB/TRR 191 – 281071066 (Symplectic Structures in Geometry, Algebra and Dynamics). The third author is partially supported by the ANR project COSY (ANR-21-CE40-0002).

ABSTRACT. — Using Givental's non-linear Maslov index we define a sequence of spectral selectors on the universal cover of the identity component of the contactomorphism group of any lens space. As applications, we prove for lens spaces with equal weights that the standard Reeb flow is a geodesic for the discriminant and oscillation norms, and we define for general lens spaces a stably unbounded conjugation invariant spectral pseudonorm.

RÉSUMÉ. — En utilisant l'indice de Maslov non-linéaire de Givental nous définissons une suite d'invariants spectraux sur le recouvrement universel de la composante de l'identité du groupe des contactomorphismes de n'importe quel espace lenticulaire. Comme applications nous démontrons pour les espaces lenticulaires de poids égaux que le flot de Reeb standard est une géodésique pour les normes discriminante et d'oscillation, et nous définissons en général une pseudonorme invariante par conjugaison qui n'est pas bornée au sens stable.

#### 1. Introduction

For any integer  $k \ge 2$  and *n*-tuple  $\underline{w} = (w_1, \dots, w_n)$  of positive integers relatively prime to k, the lens space  $L_k^{2n-1}(\underline{w})$  is the quotient of the unit sphere  $\mathbb{S}^{2n-1}$  in  $\mathbb{R}^{2n} \equiv \mathbb{C}^n$  by the free  $\mathbb{Z}_k$ -action generated by the map

$$(z_1, \cdots, z_n) \mapsto \left(e^{\frac{2\pi i}{k} \cdot w_1} z_1, \cdots, e^{\frac{2\pi i}{k} \cdot w_n} z_n\right).$$

We endow  $L_k^{2n-1}(\underline{w})$  with its canonical contact structure  $\xi_0$ , the kernel of the contact form  $\alpha_0$  whose pullback  $\bar{\alpha}_0$  by the projection  $\mathbb{S}^{2n-1} \to L_k^{2n-1}(\underline{w})$  is equal to the pullback of  $\sum_{j=1}^n x_j dy_j - y_j dx_j$  by the inclusion  $\mathbb{S}^{2n-1} \hookrightarrow \mathbb{R}^{2n}$ . We denote by  $\operatorname{Cont}_0(L_k^{2n-1}(\underline{w}), \xi_0)$  the identity component of the contactomorphism group, and by  $\widetilde{\operatorname{Cont}}_0(L_k^{2n-1}(\underline{w}), \xi_0)$  its universal cover. The non-linear Maslov index is a quasimorphism

$$\mu: \widetilde{\operatorname{Cont}}_0(L_k^{2n-1}(\underline{w}), \xi_0) \to \mathbb{Z},$$

defined by Givental [Giv90] for real projective spaces and extended to general lens spaces in [GKPS21]. Roughly speaking, it counts with multiplicity the number of intersections of contact isotopies with (a certain subspace of) the space of contactomorphisms that have at least one discriminant point.

Recall that a point p of a contact manifold  $(M,\xi)$  is said to be a discriminant point of a contactomorphism  $\phi$  if  $\phi(p) = p$  and  $(\phi^* \alpha)_p = \alpha_p$  for some (hence any) contact form  $\alpha$  for  $\xi$ , and is said to be a translated point of  $\phi$  with respect to a contact form  $\alpha$  if there exists a real number T (in general not unique) such that p is a discriminant point of  $r_{-T}^{\alpha} \circ \phi$ , where  $\{r_t^{\alpha}\}$  denotes the Reeb flow; such T is then said to be a translation of the translated point p. Discriminant and translated points play a key role in certain proofs of several global rigidity results in contact topology, related in particular to contact non-squeezing [AM18, FSZ23, San11a], orderability [AM18, Bhu01, EP00, GKPS21, San11a, San11b], and bi-invariant metrics on the contactomorphism group [Arl23, CS15, San10]. In particular, Givental's nonlinear Maslov index for projective spaces has been used in [EP00], [San13] and [CS15] respectively to prove that real projective spaces are orderable, satisfy a contact analogue of the Arnold conjecture and have unbounded discriminant and oscillation norms. All these results have then been generalized to lens spaces in [GKPS21] (recovering for orderability a result also obtained in [Mil08, San11b]). In the original article of Givental [Giv90], the non-linear Maslov index on projective spaces and a Legendrian version of it have been applied in particular to prove the Weinstein and chord conjectures, and a result on existence of Reeb chords between Legendrian submanifolds Legendrian isotopic to each other. Moreover, an analogue of the nonlinear Maslov index for complex projective spaces has been used by Givental [Giv90] and Théret [Thé98] to prove the Arnold conjectures on fixed points of Hamiltonian symplectomorphisms and Lagrangian intersections.

In the present article we use the non-linear Maslov index to define spectral selectors on the universal cover of the identity component of the contactomorphism group of lens spaces, i.e. maps

$$c_j: \widetilde{\operatorname{Cont}_0}\left(L_k^{2n-1}(\underline{w}), \xi_0\right) \to \mathbb{R}$$

that associate to every element of  $\widetilde{\operatorname{Cont}}_0(L_k^{2n-1}(\underline{w}),\xi_0)$  a real number belonging to its action spectrum. Recall that the action spectrum of a contactomorphism  $\phi$  of a contact manifold  $(M,\xi)$  with respect to a contact form  $\alpha$  is the set  $\mathcal{A}_{\alpha}(\phi)$  of real numbers that are translations of translated points of  $\phi$  with respect to  $\alpha$ . We denote by

$$\Pi: \operatorname{Cont}_0(M,\xi) \to \operatorname{Cont}_0(M,\xi)$$

the standard projection, which sends an element  $\tilde{\phi} = [\{\phi_t\}_{t \in [0,1]}]$  of  $\operatorname{Cont}_0(M,\xi)$  to  $\phi_1$ , and define the action spectrum of an element  $\tilde{\phi}$  of  $\operatorname{Cont}_0(M,\xi)$  by  $\mathcal{A}_{\alpha}(\tilde{\phi}) = \mathcal{A}_{\alpha}(\Pi(\tilde{\phi}))$ . Let

$$\mathcal{L}: \widetilde{\mathrm{Cont}}_0\left(L_k^{2n-1}(\underline{w}), \xi_0\right) \to \widetilde{\mathrm{Cont}}_0\left(\mathbb{S}^{2n-1}, \bar{\xi}_0 = \ker(\bar{\alpha}_0)\right)$$

be the map that sends  $\tilde{\phi} = [\{\phi_t\}_{t \in [0,1]}]$  to the element of  $\operatorname{Cont}_0(\mathbb{S}^{2n-1}, \bar{\xi}_0)$  represented by the lift of  $\{\phi_t\}_{t \in [0,1]}$  to  $(\mathbb{S}^{2n-1}, \bar{\xi}_0)$ . For  $\tilde{\phi} \in \operatorname{Cont}_0(L_k^{2n-1}(\underline{w}), \xi_0)$  we denote

$$\mathcal{A}(\phi) = \mathcal{A}_{\alpha_0}(\phi)$$

and

$$\bar{\mathcal{A}}(\widetilde{\phi}) = \mathcal{A}_{\bar{lpha}_0} \Big( \mathcal{L}(\widetilde{\phi}) \Big) \subset \mathcal{A}(\widetilde{\phi})$$

The sets  $\mathcal{A}(\tilde{\phi})$  and  $\bar{\mathcal{A}}(\tilde{\phi})$  are invariant by translation by  $T_{\underline{w}}$  and  $2\pi$  respectively, where  $T_{\underline{w}}$  denotes the period of the Reeb flow of  $\alpha_0$  on  $L_k^{2n-1}(\underline{w})$ . For a real number T we denote

$$\begin{bmatrix} T \end{bmatrix}_{T_{\underline{w}}} = T_{\underline{w}} \begin{bmatrix} T \\ \overline{T_{\underline{w}}} \end{bmatrix}$$
 and  $\begin{bmatrix} T \end{bmatrix}_{T_{\underline{w}}} = T_{\underline{w}} \begin{bmatrix} T \\ \overline{T_{\underline{w}}} \end{bmatrix}$ 

thus  $[T]_{T_{\underline{w}}}$  and  $[T]_{T_{\underline{w}}}$  are respectively the smallest multiple of  $T_{\underline{w}}$  greater or equal than T and the greatest multiple of  $T_{\underline{w}}$  smaller or equal than T.

Before stating our main result we recall that, since  $(L_k^{2n-1}(\underline{w}), \xi_0)$  is orderable, the relation  $\leq$  on  $\widetilde{\text{Cont}}_0(L_k^{2n-1}(\underline{w}), \xi_0)$  defined by posing  $\widetilde{\phi} \leq \widetilde{\psi}$  if there is a non-negative contact isotopy representing  $\widetilde{\psi} \cdot \widetilde{\phi}^{-1}$  is a bi-invariant partial order; this is the partial order that appears in point (vii) below. Recall also that a translated point p of a contactomorphism  $\phi$  of a contact manifold  $(M, \xi)$  with respect to a contact form  $\alpha$ is said to be non-degenerate for a translation T if there is no vector  $X \in T_p M \setminus \{0\}$  such that  $(r_{-T}^{\alpha} \circ \phi)_*(X) = X$  and dg(X) = 0, where g is the conformal factor of  $\phi$ , i.e. the function defined by the relation  $\phi^* \alpha = e^g \alpha$ . In the case of  $(\mathbb{S}^{2n-1}, \bar{\xi_0})$ , if a translated point of a contactomorphism with respect to  $\bar{\alpha}_0$  is non-degenerate for a certain translation then it is non-degenerate for all the translations; we then just say that it is non-degenerate. For any  $T \in \mathbb{R}$  we denote

$$\widetilde{r_T} = \left[ \{ r_{Tt} \}_{t \in [0,1]} \right],$$

where  $\{r_t\}$  is the Reeb flow on  $L_k^{2n-1}(\underline{w})$  of  $\alpha_0$ . Moreover, we denote by  $\widetilde{id}$  the identity on  $\widetilde{Cont_0}(L_k^{2n-1}(\underline{w}), \xi_0)$ .

Our main result is the following theorem.

THEOREM 1.1 (Spectral selectors). — There exists a non-decreasing sequence of maps

$$\left\{ c_j : \widetilde{\operatorname{Cont}}_0 \left( L_k^{2n-1}(\underline{w}), \xi_0 \right) \to \mathbb{R}, \ j \in \mathbb{Z} \right\}$$

satisfying the following properties:

(i) Spectrality:

$$c_j(\widetilde{\phi}) \in \overline{\mathcal{A}}(\widetilde{\phi})$$
.

(ii) Normalization:

$$c_0(\mathrm{id}) = 0$$

(iii) Relation with translated points: if all the translated points of  $\Pi(\mathcal{L}(\tilde{\phi}))$  with respect to  $\bar{\alpha}_0$  are non-degenerate then the spectral selectors  $\{c_j(\tilde{\phi}), j \in \mathbb{Z}\}$ are all distinct. On the other hand, if

$$c_{j-1}(\widetilde{\phi}) < c_j(\widetilde{\phi}) = c_{j+1}(\widetilde{\phi}) = \dots = c_{j+m}(\widetilde{\phi}) = T < c_{j+m+1}(\widetilde{\phi})$$

for some j and  $1 \leq m \leq 2n-1$  and either k is even or j is odd or m > 1 then  $\Pi(\mathcal{L}(\tilde{\phi}))$  has infinitely many translated points of translation T with respect to  $\bar{\alpha}_0$ .

(iv) Non-degeneracy: if

$$c_{-2n+1}(\widetilde{\phi}) = c_0(\widetilde{\phi}) = 0$$

then  $\Pi(\mathcal{L}(\tilde{\phi}))$  is the identity.

(v) Composition with the Reeb flow: for every  $T \in \mathbb{R}$  we have

$$c_j(\widetilde{r_T}\cdot\widetilde{\phi}) = c_j(\widetilde{\phi}) + T;$$

in particular,  $c_0(\widetilde{r_T}) = T$ .

(vi) Periodicity:

$$c_{j+2n}(\widetilde{\phi}) = c_j(\widetilde{\phi}) + 2\pi$$
.

(vii) Monotonicity: if  $\tilde{\phi} \leq \tilde{\psi}$  then  $c_i(\tilde{\phi}) \leq c_i(\tilde{\psi})$ .

(viii) Continuity: if  $\phi \cdot \tilde{\psi}^{-1}$  is represented by a contact isotopy with Hamiltonian function  $H_t: L_k^{2n-1}(\underline{w}) \to \mathbb{R}$  with respect to  $\alpha_0$  then

$$\int_0^1 \min H_t \, dt \leqslant c_j(\widetilde{\phi}) - c_j(\widetilde{\psi}) \leqslant \int_0^1 \max H_t \, dt$$

Moreover, each  $c_i$  is continuous with respect to the  $\mathcal{C}^1$ -topology.

(ix) Triangle inequality: if either k is even or j is even then

$$c_{j+l}(\widetilde{\phi}\cdot\widetilde{\psi}) \leqslant c_j(\widetilde{\phi}) + \left[c_l(\widetilde{\psi})\right]_{T_{\underline{w}}},$$

in particular

$$\left[c_{j+l}(\widetilde{\phi}\cdot\widetilde{\psi})\right]_{T_{\underline{w}}} \leqslant \left[c_{j}(\widetilde{\phi})\right]_{T_{\underline{w}}} + \left[c_{l}(\widetilde{\psi})\right]_{T_{\underline{w}}} \,.$$

(x) Conjugation invariance:

$$\left[c_j(\tilde{\psi}\cdot\tilde{\phi}\cdot\tilde{\psi}^{-1})\right]_{T_{\underline{w}}} = \left[c_j(\tilde{\phi})\right]_{T_{\underline{w}}}.$$

(xi) Poincaré duality:

$$\left[c_{j}(\widetilde{\phi})\right]_{T_{\underline{w}}} = -\left\lfloor c_{-j-(2n-1)}(\widetilde{\phi}^{-1})\right\rfloor_{T_{\underline{w}}}$$

Using the Hamiltonian version of the non-linear Maslov index for complex projective spaces ([Car13, Giv90, Thé98]) it is possible to define also spectral invariants

$$\left\{ c_j : \widetilde{\operatorname{Ham}}(\mathbb{CP}^n, \omega_0) \to \mathbb{R}, \ j \in \mathbb{Z} \right\}$$

satisfying properties analogue to those of Theorem 1.1, with stronger statements for (ix), (x) and (xi) not involving the  $T_w$ -floors and ceilings. Such spectral invariants coincide with the ones defined by the first author in [All22a], and their projections to  $\mathbb{S}^1$  coincide with the rotation numbers defined by Théret in [Thé98]. Moreover, their properties are analogue to those satisfied by the spectral invariants defined with Floer homology by Entov and Polterovich in [EP03]. The fact that in the contact case the statements of the triangle inequality, conjugation invariance and Poincaré duality properties are weaker than in the symplectic case and involve the  $T_w$ -floors and ceilings is similar to what happens for the spectral selectors of compactly supported contactomorphisms of  $(\mathbb{R}^{2n} \times \mathbb{S}^1, \xi_0)$  defined by the third author in [San11a]. Indeed, these spectral selectors are contact analogues of the spectral selectors of compactly supported Hamiltonian symplectomorphisms of  $(\mathbb{R}^{2n}, \omega_0)$  defined by Viterbo in [Vit92], but they satisfy weaker versions of the triangle inequality, conjugation invariance and Poincaré duality properties involving their (integral) floors and ceilings. Roughly speaking, this difference with respect to the symplectic case can be explained as follows. If we see  $(\mathbb{R}^{2n} \times \mathbb{S}^1, \xi_0)$  and  $(L_k^{2n-1}(1, \cdots, 1), \xi_0)$  as prequantizations of  $(\mathbb{R}^{2n}, \omega_0)$  and  $(\mathbb{CP}^{n-1}, \omega_0)$  respectively then in both cases the contact spectral selectors are generalizations of the symplectic ones, in the sense that the symplectic spectral selectors of Hamiltonian isotopies of  $(\mathbb{R}^{2n}, \omega_0)$  and  $(\mathbb{CP}^{n-1}, \omega_0)$  coincide with the contact spectral selectors of their lifts to  $(\mathbb{R}^{2n} \times \mathbb{S}^1, \xi_0)$  and  $(L_k^{2n-1}(1, \dots, 1), \xi_0)$ respectively. The fact that the contact spectral selectors satisfy weaker versions of the triangle inequality, conjugation invariance and Poincaré duality properties involving their floors and ceilings with respect to the period of the Reeb flow is due to the fact that, while the lifts of Hamiltonian isotopies of  $(\mathbb{R}^{2n}, \omega_0)$  and  $(\mathbb{CP}^{n-1}, \omega_0)$  are exactly the contact isotopies that commute with the standard Reeb flows, general contact isotopies commute with the Reeb flow at time t only when t is a multiple of the period of the Reeb flow. For conjugation invariance, for instance, while in the symplectic case the action spectrum is invariant by conjugation, in the contact

case this is in general not true: the translated points of a contactomorphism are in general not in bijection with those of a conjugation. However, if the Reeb flow is periodic then the translated points of translation equal to the period of the Reeb flow are discriminant points, which are invariant by conjugation, and this fact can be used to prove that the corresponding floor and ceiling of the spectral selectors are invariant by conjugation (see also the discussion in [San11a]).

In [AA23] the first and second authors have defined invariants  $c_{+}$  and  $c_{-}$  for elements of the universal cover of any closed orderable contact manifold and for contactomorphisms of any closed contact manifold with orderable contactomorphism group. In the universal cover case, these invariants satisfy all the properties in Theorem 1.1 (including conjugation invariance if the Reeb flow is periodic, and with stronger versions for the triangle inequality and Poincaré duality properties not involving floors and ceilings) except for periodicity (there are only two invariants  $c_{+}$  and  $c_{-}$ , while we have a sequence  $c_{i}$  related by periodicity), spectrality and (iii). These properties are important for us to obtain the applications discussed below. In particular, spectrality is crucial to obtain Corollary 1.5 and the relation between the pseudonorm  $\nu$  of Corollary 1.6 and the oscillation norm, while periodicity is used in Corollary 1.6 to show that the induced norm  $\nu_*$  is bounded (see also Remarks 1.2 and 1.3 below for two more consequences of these properties). The first and second authors also defined in [AA23] invariants for Legendrian submanifolds and Legendrian isotopies (when the involved spaces are orderable) that do satisfy a spectrality property. Using the Legendrian version of the non-linear Maslov index defined in [Giv90] it should be possible to obtain also a Legendrian version of our spectral selectors, with properties similar to those in Theorem 1.1. However, as far as we can see, the only new application of these spectral selectors with respect to those in [AA23] would be a better lower bound for the number of Reeb chords between Legendrian submanifolds Legendrian isotopic to each other, but (at least in the case of real projective space) such bound is already given by Givental in [Giv90] just using the non-linear Maslov index.

Remark 1.2. — Properties (vi), (ix) and (xi) imply that each  $c_j$  is a quasimorphism.

Remark 1.3. — Properties (i), (iii) and (vi) imply that every contactomorphism of  $(L_k^{2n-1}(\underline{w}), \xi_0)$  contact isotopic to the identity has at least n translated points with respect to  $\alpha_0$ , and at least 2n if either k is even or all the translated points are non-degenerate. We thus recover the corresponding result of [GKPS21], but not the optimal bound obtained by the first author in [All22b], where it is proved that every contactomorphism of  $(L_k^{2n-1}(\underline{w}), \xi_0)$  contact isotopic to the identity has at least 2ntranslated points with respect to  $\alpha_0$ .

Remark 1.4. — Suppose that k is prime. Recall that the cohomological index ind(A) of a subset A of  $L_k^{2n-1}(\underline{w})$  is the dimension over  $\mathbb{Z}_k$  of the image of the map  $\check{H}^*(L_k^{2n-1}(\underline{w});\mathbb{Z}_k) \to \check{H}^*(A;\mathbb{Z}_k)$  on Čech cohomology induced by the inclusion  $A \hookrightarrow L_k^{2n-1}(\underline{w})$ . As we will see, property (iii) can be refined in this case as follows: if

$$c_{j-1}(\widetilde{\phi}) < c_j(\widetilde{\phi}) = c_{j+1}(\widetilde{\phi}) = \dots = c_{j+m}(\widetilde{\phi}) = T < c_{j+m+1}(\widetilde{\phi})$$

for some j and  $1 \leq m \leq 2n - 1$  then the set of translated points of translation T of  $\Pi(\tilde{\phi})$  has cohomological index greater or equal than m, and greater or equal than m+1 if either k=2 or j is odd.

As a first application of Theorem 1.1 we prove that the standard Reeb flow on lens spaces with equal weights is a geodesic for the discriminant and oscillation norms introduced in [CS15]. The definition of the discriminant norm  $\nu_{\rm dis}$  and of the oscillation pseudonorm  $\nu_{\rm osc}$  on the universal cover of the identity component of the contactomorphism group of a closed contact manifold  $(M,\xi)$  are recalled in Section 4 below, as well as the definition of the discriminant and oscillation lengths of contact isotopies. Recall also from [CS15, Proposition 3.2] that the oscillation pseudonorm is non-degenerate if and only if  $(M,\xi)$  is orderable; in particular, it is thus a norm for lens spaces. As in [Arl23], we say that a contact isotopy of a closed orderable contact manifold is a geodesic for the discriminant or for the oscillation norm if its discriminant or oscillation length is equal to the discriminant or oscillation norm of the element of the universal cover it represents. In other words, a contact isotopy is a geodesic for the discriminant or oscillation norm if it minimizes the discriminant or oscillation length in its homotopy class with fixed endpoints. In [Arl23] it is proved that certain contact isotopies of  $(\mathbb{R}^{2n} \times \mathbb{S}^1, \xi_0)$  are geodesics for the discriminant and oscillation norms. We obtain here a similar result for lens spaces with equal weights, answering a question in [CS15]. In [CS15, GKPS21] respectively it is proved that the discriminant and oscillation norms on real projective spaces and on general lens spaces are unbounded, by showing that the classes in the universal cover represented by higher iterations of the Reeb flow have bigger and bigger discriminant and oscillation norms. More precisely, it is proved in [GKPS21] that for every N the discriminant and oscillation norms on  $\widetilde{\text{Cont}}_0(L_k^{2n-1}(\underline{w}), \xi_0)$  of  $\widetilde{r_{6\pi N}}$  and  $\widetilde{r_{20\pi N}}$  respectively are at least equal to N + 1. Since the discriminant length of  $\{r_{6\pi Nt}\}_{t \in [0,1]}$  is 3Nk + 1 and the oscillation length of  $\{r_{20\pi Nt}\}_{t\in[0,1]}$  is 10Nk+1, the results in [GKPS21] (as well as the previous ones in [CS15]) left open the question of whether there exist contact isotopies in the same homotopy class with fixed endpoints as  $\{r_{6\pi Nt}\}_{t \in [0,1]}$ or  $\{r_{20\pi Nt}\}_{t \in [0,1]}$  having shorter discriminant or oscillation lengths. Note that this is what happens for the sphere  $(\mathbb{S}^{2n-1}, \overline{\xi}_0)$ : the N<sup>th</sup> iteration  $\{r_{2\pi Nt}\}_{t \in [0,1]}$  of the Reeb flow  $\{r_{2\pi t}\}_{t \in [0,1]}$  of  $\bar{\alpha}_0$  has discriminant and oscillation length N+1, but by [CS15, Proposition 4.3] the discriminant norm and the oscillation pseudonorm of  $[\{r_{2\pi Nt}\}_{t \in [0,1]}]$  are smaller or equal than 4; in other words, there exist contact isotopies of  $(\mathbb{S}^{2n-1}, \overline{\xi}_0)$  in the same homotopy class with fixed endpoints as certain iterations the Reeb flow having strictly shorter discriminant and oscillation length. As an application of Theorem 1.1, in Section 4 we show that for lens spaces with equal weights this is not possible. More precisely, we show that Theorem 1.1 implies the following result.

COROLLARY 1.5 (Non-shortening of the standard Reeb flow). — For every real number T, the Reeb flow  $\{r_{Tt}\}_{t \in [0,1]}$  of the standard contact form  $\alpha_0$  on a lens space of the form  $L_k^{2n-1}(w, \dots, w)$  is a geodesic for the discriminant and oscillation norms.

In particular,

$$\nu_{\rm dis}(\widetilde{r_T}) = \nu_{\rm osc}(\widetilde{r_T}) = \left\lfloor \frac{k}{2\pi} T \right\rfloor + 1.$$

For general lens spaces  $L_k^{2n-1}(\underline{w})$  we also obtain lower bounds for the discriminant and oscillation norms of the standard Reeb flow that are sharper than those in [CS15, GKPS21], however in general these bounds are not sharp enough to prove that the Reeb flow is a geodesic (see Remark 4.2).

Using the spectral selectors of Theorem 1.1 we also define a stably unbounded conjugation invariant pseudonorm on  $\widetilde{\text{Cont}}_0(L_k^{2n-1}(\underline{w}),\xi_0)$ . More precisely, posing  $c_- = c_{-2n+1}$  and  $c_+ = c_0$  we prove the following result.

COROLLARY 1.6 (Spectral pseudonorm). — The map  $\nu : \widetilde{\text{Cont}_0}(L_k^{2n-1}(\underline{w}), \xi_0) \to T_{\underline{w}} \cdot \mathbb{Z}$  defined by

$$\nu(\widetilde{\phi}) = \max\left\{\left[c_{+}(\widetilde{\phi})\right]_{T_{\underline{w}}}, -\left\lfloor c_{-}(\widetilde{\phi})\right\rfloor_{T_{\underline{w}}}\right\}$$

is a stably unbounded conjugation invariant pseudonorm, which is compatible with the partial order  $\leq$  and satisfies  $\nu(\tilde{\phi}) \leq T_{\underline{w}} \cdot \nu_{\text{osc}}(\tilde{\phi})$  for every  $\tilde{\phi} \in \text{Cont}_0(L_k^{2n-1}(\underline{w}), \xi_0)$ . The induced pseudonorm  $\nu_*$  on  $\text{Cont}_0(L_k^{2n-1}(\underline{w}), \xi_0)$  is non-degenerate and bounded.

Finally, we remark that the spectral selectors of Theorem 1.1 can also be used as in [AA23] to define a time function on  $\widetilde{\text{Cont}}_0(L_k^{2n-1}(\underline{w}), \xi_0)$ , i.e. a function

$$\tau : \widetilde{\mathrm{Cont}}_0 \left( L_k^{2n-1}(\underline{w}), \xi_0 \right) \to \mathbb{R}$$

that is continuous with respect to the  $\mathcal{C}^1$ -topology and satisfies  $\tau(\tilde{\phi}) < \tau(\tilde{\psi})$  whenever  $\tilde{\phi} \leq \tilde{\psi}$  with  $\tilde{\phi} \neq \tilde{\psi}$ . Such function can be defined by

$$\tau(\widetilde{\phi}) = a \sum_{j \in \mathbb{N}} \frac{c_0(\widetilde{\phi} \cdot \widetilde{\psi}_j)}{2^j \max\left(1, \left|\left[c_0(\widetilde{\psi}_j)\right]_{T_{\underline{w}}}\right|, \left|\left[c_0\left(\widetilde{\psi}_j^{-1}\right)\right]_{T_{\underline{w}}}\right|\right)} + b$$

with

$$a = \left(\sum_{j \in \mathbb{N}} \frac{1}{2^j \max\left(1, \left|\left\lceil c_0(\tilde{\psi}_j)\right\rceil_{T_{\underline{w}}}\right|, \left|\left\lceil c_0\left(\tilde{\psi}_j^{-1}\right)\right\rceil_{T_{\underline{w}}}\right|\right)}\right)^{-1}$$

and b chosen so that  $\tau(\widetilde{id}) = 0$ , where  $(\widetilde{\psi}_j)_{j \ge 1}$  is any sequence in  $\widetilde{Cont}_0(L_k^{2n-1}(\underline{w}), \xi_0)$  that is dense with respect to the  $\mathcal{C}^1$ -topology. As in [AA23], the time function  $\tau$  satisfies moreover  $\tau(\widetilde{r_T} \cdot \widetilde{\phi}) = T + \tau(\widetilde{\phi})$  for all T and  $\widetilde{\phi}$ .

The article is organized as follows. In Section 2 we recall the definition of the non-linear Maslov index and discuss the properties that are needed for the construction of the spectral selectors. In Section 3 we define the spectral selectors and prove Theorem 1.1. In Section 4 we prove Corollary 1.5, and in Section 5 we prove Corollary 1.6.

#### Acknowledgments

The third author thanks Alberto Abbondandolo for the invitation to Bochum and kind hospitality in the fall of 2022, when part of this work has been done. We all thank Alberto also for his support and for inspiring discussions. The version of this article that was first posted to arXiv contained a mistake concerning the period of the Reeb flow on lens spaces with general weights and affecting in particular Corollary 1.5, which was stated in more generality than what we can actually prove. We are grateful to Miguel Abreu for noticing this mistake and for very useful discussions.

#### 2. The non-linear Maslov index

In this section we recall the definition of the non-linear Maslov index

$$\mu : \operatorname{Cont}_0(L_k^{2n-1}(\underline{w}), \xi_0) \to \mathbb{Z}$$

following the presentation in [GKPS21], to which we refer for more details. We also discuss the properties of the non-linear Maslov index that are needed for the construction of the spectral selectors. Several of these properties do not appear in [GKPS21], and so we include detailed proofs.

Since the weights  $\underline{w}$  do not play a particular role in the discussion, in this section we denote  $L_k^{2n-1}(\underline{w})$  simply by  $L_k^{2n-1}$ . As in [GKPS21], we first define the nonlinear Maslov index assuming that k is prime and then obtain the general case (Proposition 2.12) by pullback. Assume thus for now that k is prime.

The construction of the non-linear Maslov index is based on generating functions. Recall that a function  $F : E \to \mathbb{R}$  defined on the total space of a fibre bundle  $p : E \to B$  is said to be a generating function if the differential  $dF : E \to T^*E$ is transverse to the fibre conormal bundle  $N_E^*$ , the space of points  $(e, \eta)$  of  $T^*E$ such that  $\eta$  vanishes on the kernel of dp(e). Then the set  $\Sigma_F = (dF)^{-1}(N_E^*)$  of fibre critical points of F is a submanifold of E, and the map

$$i_F: \Sigma_F \to T^*B, \ e \mapsto (p(e), v^*(e))$$

defined by posing  $v^*(e)(X) = dF(\widehat{X})$  for  $X \in T_{p(e)}B$ , where  $\widehat{X}$  is any vector in  $T_e E$  with  $dp(e)(\widehat{X}) = X$ , is a Lagrangian immersion with respect to the canonical symplectic form  $\omega_{\text{can}}$  on  $T^*B$ . If  $i_F$  is an embedding then F is said to be a generating function of the Lagrangian submanifold  $i_F(\Sigma_F)$  of  $(T^*B, \omega_{\text{can}})$ . A function F is said to be a generating function of a symplectomorphism  $\Phi$  of  $(\mathbb{R}^{2n}, \omega_0)$  if it is a generating function of the Lagrangian submanifold of  $(T^*\mathbb{R}^{2n}, \omega_{\text{can}})$  that is the image of the graph of  $\Phi$  by the symplectomorphism  $\tau : \mathbb{R}^{2n} \times \mathbb{R}^{2n} \to T^*\mathbb{R}^{2n}$  defined by

$$\tau(x, y, X, Y) = \left(\frac{x+X}{2}, \frac{y+Y}{2}, Y-y, x-X\right).$$

Any contact isotopy  $\{\phi_t\}_{t \in [0,1]}$  of  $(L_k^{2n-1}, \xi_0)$  starting at the identity can be uniquely lifted to a  $\mathbb{Z}_k$ -equivariant contact isotopy  $\{\bar{\phi}_t\}_{t \in [0,1]}$  of  $(\mathbb{S}^{2n-1}, \bar{\xi}_0)$  starting at the identity, which in turn can be uniquely extended to a conical Hamiltonian isotopy

 $\{\Phi_t\}_{t \in [0,1]}$  of  $(\mathbb{R}^{2n}, \omega_0)$ , i.e. a 1-parameter family of  $(\mathbb{Z}_k \times \mathbb{R}_{>0})$ -equivariant homeomorphisms of  $\mathbb{R}^{2n}$  that is a Hamiltonian isotopy on  $\mathbb{R}^{2n} \setminus \{0\}$ . If M is a multiple of n then we say that a function  $F: \mathbb{R}^{2M} \to \mathbb{R}$  is conical if it is  $\mathcal{C}^1$  with Lipschitz differential, homogeneous of degree 2 with respect to the radial action of  $\mathbb{R}_{>0}$  on  $\mathbb{R}^{2M}$ , and invariant by the diagonal action of  $\mathbb{Z}_k$  on  $\mathbb{R}^{2M}$ . We say that a conical function  $F: E \to \mathbb{R}$  defined on the total space of a trivial vector bundle  $E = \mathbb{R}^{2n} \times \mathbb{R}^{2nN} \to \mathbb{R}^{2n}$  is a conical generating functions of a conical symplectomorphism  $\Phi$  of  $(\mathbb{R}^{2n}, \omega_0)$ , i.e. a  $(\mathbb{Z}_k \times \mathbb{R}_{>0})$ -equivariant homeomorphism of  $\mathbb{R}^{2n}$  that is a symplectomorphism on  $\mathbb{R}^{2n} \setminus \{0\}$ , if it is smooth near its fibre critical points other than the origin,  $dF: E \to T^*E$  is transverse to the fibre conormal bundle  $N_E^*$  except possibly at the origin, and  $i_F$  is a homeomorphism between  $\Sigma_F$  and the image of the graph of  $\Phi$  by  $\tau$ . We say that  $F_t : \mathbb{R}^{2n} \times \mathbb{R}^{2nN} \to \mathbb{R}, t \in [0, 1]$ , is a family of conical generating functions for a contact isotopy  $\{\phi_t\}_{t \in [0,1]}$  of  $(L_k^{2n-1}, \xi_0)$ starting at the identity if for every t the function  $F_t$  is a conical generating function of  $\Phi_t$ , where  $\{\Phi_t\}$  denotes the conical Hamiltonian isotopy of  $(\mathbb{R}^{2n}, \omega_0)$  lifting  $\{\phi_t\}$ , and the map  $(e,t) \mapsto F_t(e)$  is  $\mathcal{C}^1$  with locally Lipschitz differential and smooth near (e, t) whenever e is a fibre critical point of  $F_t$  other than the origin. We say that  $F_t$ ,  $t \in [0, 1]$ , is a based family of conical generating functions for  $\{\phi_t\}_{t \in [0,1]}$  if moreover  $F_0$  is equivalent to the zero function on  $\mathbb{R}^{2n}$ , where we consider on the set of conical generating functions the smallest equivalence relation under which two such functions are equivalent if they differ by a stabilization (i.e. replacing a conical generating function  $F: \mathbb{R}^{2n} \times \mathbb{R}^{2nN} \to \mathbb{R}$  by  $F \oplus Q: \mathbb{R}^{2n} \times \mathbb{R}^{2nN} \times \mathbb{R}^{2nN'} \to \mathbb{R}$  for a non-degenerate  $\mathbb{Z}_k$ -invariant quadratic form Q on  $\mathbb{R}^{2nN'}$ ) or by a fibre preserving conical homeomorphism (i.e. a  $(\mathbb{Z}_k \times \mathbb{R}_{>0})$ -equivariant homeomorphism of  $\mathbb{R}^{2n} \times \mathbb{R}^{2nN}$ that takes each fibre  $\{z\} \times \mathbb{R}^{2nN}$  to itself) that restricts to a diffeomorphism between neighborhoods of fibre critical points other than the origin. It is proved in [GKPS21, Proposition 2.14] that any contact isotopy  $\{\phi_t\}_{t \in [0,1]}$  of  $(L_k^{2n-1}, \xi_0)$  starting at the identity has a based family  $F_t : \mathbb{R}^{2n} \times \mathbb{R}^{2nN} \to \mathbb{R}$  of conical generating functions. A conical function  $F : \mathbb{R}^{2M} \to \mathbb{R}$  induces uniquely a function  $f : L_k^{2M-1} \to \mathbb{R}$ , which

A conical function  $F : \mathbb{R}^{2M} \to \mathbb{R}$  induces uniquely a function  $f : L_k^{2M-1} \to \mathbb{R}$ , which is  $\mathcal{C}^1$  with Lipschitz differential. All the critical points of F have critical value zero and come in  $(\mathbb{Z}_k \times \mathbb{R}_{>0})$ -families; moreover, there is a 1–1 correspondence between the  $(\mathbb{Z}_k \times \mathbb{R}_{>0})$ -families of critical points of F and the critical points of critical value zero of f. If F is a conical generating function of a conical symplectomorphism  $\Phi$  whose restriction  $\overline{\phi}$  to  $\mathbb{S}^{2n-1}$  projects to a contactomorphism  $\phi$  of  $(L_k^{2n-1}, \xi_0)$  then there is a 1–1 correspondence between the critical points of critical value zero of f and the discriminant points of  $\phi$  that lift to discriminant points of  $\overline{\phi}$ . In order to detect discriminant points of contactomorphisms of  $(L_k^{2n-1}, \xi_0)$  we thus study the topology of the sublevel set at zero of the functions on (possibly higher dimensional) lens spaces induced by the corresponding conical generating functions. The topological invariant that we use for this is the cohomological index for subsets of lens spaces: for a subset A of  $L_k^{2M-1}$  such index, which we denote by  $\operatorname{ind}(A)$ , is the dimension over  $\mathbb{Z}_k$  of the image of the map  $\check{H}^*(L_k^{2M-1}; \mathbb{Z}_k) \to \check{H}^*(A; \mathbb{Z}_k)$  on Čech cohomology induced by the inclusion  $A \hookrightarrow L_k^{2M-1}$ . For any conical function  $F : \mathbb{R}^{2M} \to \mathbb{R}$  we thus define

$$\operatorname{ind}(F) = \operatorname{ind}(\{f \leq 0\}).$$

The following property is proved in [GKPS21, Corollary 3.15] (cf. [GKPS21, Proposition 3.14, Proposition 3.9(v) and Remark 3.11] for the case k = 2).

LEMMA 2.1. — For any two conical functions F and G we have

$$|\operatorname{ind}(F \oplus G) - \operatorname{ind}(F) - \operatorname{ind}(G)| \leq 1$$
,

and

$$\operatorname{ind}(F \oplus G) = \operatorname{ind}(F) + \operatorname{ind}(G)$$

if either k = 2 or ind(F) is even or ind(G) is even.

For a contact isotopy  $\{\phi_t\}_{t \in [0,1]}$  of  $(L_k^{2n-1}, \xi_0)$  we define

$$\mu(\{\phi_t\}_{t \in [0,1]}) = \operatorname{ind}(F_0) - \operatorname{ind}(F_1),$$

where  $F_t, t \in [0, 1]$ , is any based family of conical generating functions for  $\{\phi_t\}_{t \in [0,1]}$ . It is proved in [GKPS21, Proposition 2.20] that any two based families of conical generating functions for  $\{\phi_t\}_{t \in [0,1]}$  are equivalent, where we consider on the set of based families of conical generating functions the smallest equivalence relation under which two such families are equivalent if they differ by a stabilization (i.e. replacing  $F_t : \mathbb{R}^{2n} \times \mathbb{R}^{2nN} \to \mathbb{R}$  by  $F_t \oplus Q : \mathbb{R}^{2n} \times \mathbb{R}^{2nN'} \to \mathbb{R}$  for a non-degenerate  $\mathbb{Z}_k$ -invariant quadratic form Q on  $\mathbb{R}^{2nN'}$ ) or by a 1-parameter family of fibre preserving conical homeomorphism that restrict to diffeomorphisms between neighborhoods of fibre critical points other than the origin. Since, for k > 2,  $\operatorname{ind}(Q)$  is even for every  $\mathbb{Z}_k$ -invariant quadratic form Q ([GKPS21, Remark 3.13]), it thus follows from Lemma 2.1 that  $\mu(\{\phi_t\}_{t \in [0,1]})$  is well-defined, i.e. it does not depend on the choice of a based family of conical generating functions. Moreover, it is proved in [GKPS21] (as a consequence of [GKPS21, Proposition 2.21]) that  $\mu$  descends to a map

$$\mu: \widetilde{\operatorname{Cont}_0}\left(L_k^{2n-1}, \xi_0\right) \to \mathbb{Z}.$$

Example 2.2. — By definition we have  $\mu(id) = 0$ . By [GKPS21, Example 4.1], if  $\tilde{\phi}$  is small enough in the  $C^1$ -topology then  $0 \leq \mu(\tilde{\phi}) \leq 2n$ , and if moreover  $\tilde{\phi}$  is positive then  $\mu(\tilde{\phi}) = 2n$ . In particular, for  $\epsilon > 0$  small enough we have  $\mu(\tilde{r_{\epsilon}}) = 2n$ . By [GKPS21, Example 4.13], for every integer m we have  $\mu(\tilde{r_{2\pi m}}) = 2nm$ .

It is proved in [GKPS21, Theorem 1.4(i) and Remark 1.7] that for any two elements  $\widetilde{\phi}$  and  $\widetilde{\psi}$  of  $\widetilde{\text{Cont}}_0(L_k^{2n-1}, \xi_0)$  we have

(2.1) 
$$\left| \mu(\widetilde{\phi} \cdot \widetilde{\psi}) - \mu(\widetilde{\phi}) - \mu(\widetilde{\psi}) \right| \leq 2n+1,$$

and

(2.2) 
$$\left| \mu(\widetilde{\phi} \cdot \widetilde{\psi}) - \mu(\widetilde{\phi}) - \mu(\widetilde{\psi}) \right| \leq 2n$$

if k = 2; in particular,  $\mu : \widetilde{\text{Cont}}_0(L_k^{2n-1}, \xi_0) \to \mathbb{Z}$  is a quasimorphism. The non-linear Maslov index also satisfies the following triangle inequality, which is not proved in [GKPS21].

PROPOSITION 2.3. — For any two elements  $\tilde{\phi}$  and  $\tilde{\psi}$  of  $\widetilde{\text{Cont}}_0(L_k^{2n-1}, \xi_0)$  we have  $\mu(\tilde{\phi} \cdot \tilde{\psi}) \leq \mu(\tilde{\phi}) + \mu(\tilde{\psi}) + 1$ ,

and

$$\mu(\widetilde{\phi}\cdot\widetilde{\psi})\leqslant\mu(\widetilde{\phi})+\mu(\widetilde{\psi})$$

if either k = 2 or  $\mu(\tilde{\phi})$  is even or  $\mu(\tilde{\psi})$  is even.

Before proving Proposition 2.3, recall ([GKPS21, Proposition 2.10]) that if  $F : \mathbb{R}^{2n} \times \mathbb{R}^{2nN_1} \to \mathbb{R}$  and  $G : \mathbb{R}^{2n} \times \mathbb{R}^{2nN_2} \to \mathbb{R}$  are conical generating functions for conical symplectomorphisms  $\Phi$  and  $\Psi$  respectively, then the function

$$F \, \sharp \, G \colon \mathbb{R}^{2n} \times \left( \mathbb{R}^{2n} \times \mathbb{R}^{2n} \times \mathbb{R}^{2nN_1} \times \mathbb{R}^{2nN_2} \right) \to \mathbb{R}$$

defined by

$$F \sharp G(q; \zeta_1, \zeta_2, \nu_1, \nu_2) = F(\zeta_1, \nu_1) + G(\zeta_2, \nu_2) - 2 \langle \zeta_2 - q, i(\zeta_1 - q) \rangle$$

is a conical generating function of  $\Psi \circ \Phi$ , and ([GKPS21, Proposition 2.26]) there is a linear  $(\mathbb{Z}_k \times \mathbb{R}_{>0})$ -equivariant injection

$$\iota \colon \mathbb{R}^{2n} \times \mathbb{R}^{2n} \times \mathbb{R}^{2nN_1} \times \mathbb{R}^{2nN_2} \to \mathbb{R}^{2n} \times \mathbb{R}^{2n} \times \mathbb{R}^{2n} \times \mathbb{R}^{2nN_1} \times \mathbb{R}^{2nN_2}$$

such that  $(F \sharp G) \circ \iota = F \oplus G$ . Since the cohomological index is monotone, i.e.  $ind(A) \leq ind(B)$  if  $A \subset B$  [GKPS21, Proposition 3.9(i)], we deduce that

(2.3) 
$$\operatorname{ind}(F \oplus G) \leq \operatorname{ind}(F \, \sharp \, G)$$
.

Recall also ([GKPS21, Lemma 4.2]) that if F and G are equivalent to the zero function then

$$\operatorname{ind}(F \,\sharp\, G) = \operatorname{ind}(F) + \operatorname{ind}(G)$$
.

Proof of Proposition 2.3. — Let  $F_t$  and  $G_t$  be based families of conical generating functions for contact isotopies  $\{\phi_t\}_{t \in [0,1]}$  and  $\{\psi_t\}_{t \in [0,1]}$  representing  $\tilde{\phi}$  and  $\tilde{\psi}$  respectively. Then, by [GKPS21, Proposition 2.10 and Remark 2.14],  $G_t \ \# F_t$  is a based family of conical generating functions for  $\{\phi_t \circ \psi_t\}_{t \in [0,1]}$ , and thus

$$\mu(\phi \cdot \psi) = \operatorname{ind}(G_0 \,\sharp F_0) - \operatorname{ind}(G_1 \,\sharp F_1) \,.$$

Since  $F_0$  and  $G_0$  are equivalent to the zero function, we have

$$\operatorname{ind}(G_0 \,\sharp\, F_0) = \operatorname{ind}(G_0) + \operatorname{ind}(F_0) \,.$$

Moreover, by (2.3) we have

$$-\operatorname{ind}(G_1 \,\sharp\, F_1) \leqslant -\operatorname{ind}(G_1 \oplus F_1).$$

Using Lemma 2.1 we thus deduce that

$$\mu(\tilde{\phi} \cdot \tilde{\psi}) \leq \operatorname{ind}(F_0) + \operatorname{ind}(G_0) - \operatorname{ind}(F_1) - \operatorname{ind}(G_1) + 1 = \mu(\tilde{\phi}) + \mu(\tilde{\psi}) + 1,$$

and that

$$\mu(\widetilde{\phi}\cdot\widetilde{\psi})\leqslant \mu(\widetilde{\phi})+\mu(\widetilde{\psi})$$

if either k = 2 or  $\operatorname{ind}(F_1)$  is even or  $\operatorname{ind}(G_1)$  is even, hence (since, for k > 2,  $\operatorname{ind}(F_0)$ and  $\operatorname{ind}(G_0)$  are even by [GKPS21, Remark 3.13]) if either k = 2 or  $\mu(\tilde{\phi})$  is even or  $\mu(\tilde{\psi})$  is even.

It is proved in [Giv90] that

$$\mu(\widetilde{\phi} \,\cdot\, \widetilde{\psi}) = \mu(\widetilde{\phi}) + \mu(\widetilde{\psi})$$

if either  $\mathcal{L}(\tilde{\phi})$  or  $\mathcal{L}(\tilde{\psi})$  are in  $\pi_1(\text{Cont}_0(\mathbb{S}^{2n-1}, \bar{\xi}_0))$ . For our applications we only need this property in the case when one of the factors is the Reeb flow.

PROPOSITION 2.4. — For every element  $\phi$  of  $\widetilde{\text{Cont}}_0(L_k^{2n-1},\xi_0)$  and every integer m we have

$$\mu\left(\widetilde{\phi} \cdot \widetilde{r_{2\pi m}}\right) = \mu\left(\widetilde{r_{2\pi m}} \cdot \widetilde{\phi}\right) = 2nm + \mu(\widetilde{\phi})$$

Proof. — Since  $\tilde{\phi} \cdot \tilde{r_{2\pi m}} = \tilde{r_{2\pi m}} \cdot \tilde{\phi}$ , it is enough to prove that  $\mu(\tilde{\phi} \cdot \tilde{r_{2\pi m}}) = 2nm + \mu(\tilde{\phi})$ . We represent  $\tilde{\phi} \cdot \tilde{r_{2\pi m}}$  by the concatenation

$$\{\varphi_t\}_{t \in [0,1]} = \{r_{4\pi m t}\}_{t \in [0,\frac{1}{2}]} \sqcup \{\phi_{2t-1}\}_{t \in [\frac{1}{2},1]}$$

where  $\{\phi_t\}_{t\in[0,1]}$  is a contact isotopy representing  $\tilde{\phi}$ , and we consider a based family  $F_t, t\in[0,1]$ , of conical generating functions for  $\{\varphi_t\}_{t\in[0,1]}$  so that  $F_t, t\in[0,\frac{1}{2}]$ , is a family of quadratic forms generating  $\{r_{4\pi m t}\}_{t\in[0,\frac{1}{2}]}$  (cf. [GKPS21, Proposition 4.9]). Since the lift of  $\{r_{4\pi m t}\}_{t\in[0,\frac{1}{2}]}$  to  $(\mathbb{S}^{2n-1}, \bar{\xi}_0)$  is a loop, by [GKPS21, Lemma 4.10] the quadratic form  $F_{\frac{1}{2}}$  is equivalent to the zero function, and so

$$\mu\left(\{\phi_t\}_{t\in[0,1]}\right) = \operatorname{ind}\left(F_{\frac{1}{2}}\right) - \operatorname{ind}(F_1).$$

We thus have

$$\mu\left(\widetilde{\phi} \cdot \widetilde{r_{2\pi m}}\right) = \operatorname{ind}(F_0) - \operatorname{ind}(F_1) = \operatorname{ind}(F_0) - \operatorname{ind}\left(F_{\frac{1}{2}}\right) + \operatorname{ind}\left(F_{\frac{1}{2}}\right) - \operatorname{ind}(F_1)$$
$$= \mu(\widetilde{r_{2\pi m}}) + \mu(\widetilde{\phi}) = 2nm + \mu\left(\{\phi_t\}_{t \in [0,1]}\right),$$

where the last equality follows from Example 2.2.

In the next section we also need the following result.

PROPOSITION 2.5. — For every  $\widetilde{\phi} \in \widetilde{\text{Cont}_0}(L_k^{2n-1}, \xi_0)$  and every  $T_0 \in \mathbb{R}$  there exists  $\epsilon > 0$  such that  $\mu(\widetilde{r_{-T}} \cdot \widetilde{\phi}) = \mu(\widetilde{r_{-T_0}} \cdot \widetilde{\phi})$  for every  $T \in [T_0, T_0 + \epsilon)$ .

Before proving Proposition 2.5, recall that the cohomological index is continuous ([GKPS21, Proposition 3.9(ii)]): every closed subset A of  $L_k^{2M-1}$  has a neighborhood  $\mathcal{U}$  such that if  $A \subset \mathcal{V} \subset \mathcal{U}$  then  $\operatorname{ind}(\mathcal{V}) = \operatorname{ind}(A)$ .

Proof of Proposition 2.5. — Let  $\{\phi_t\}_{t \in [0,1]}$  be a contact isotopy representing  $\phi$ , and let  $F_t$ ,  $t \in [0,1]$ , be a based family of conical generating functions for the concatenation

$$\left\{\phi_{2t}\right\}_{t \in \left[0, \frac{1}{2}\right]} \sqcup \left\{r_{-(2t-1)(T_0 + \epsilon')}\right\}_{t \in \left[\frac{1}{2}, 1\right]}$$

for some  $\epsilon' > 0$ . Then for every  $T \in [T_0, T_0 + \epsilon')$  we have

$$\mu\left(\widetilde{r_{-T}}\cdot\widetilde{\phi}\right) = \mathrm{ind}\left(\{f_0\leqslant 0\}\right) - \mathrm{ind}\left(\left\{f_{\frac{1}{2}\left(\frac{T}{T_0+\epsilon'}+1\right)}\leqslant 0\right\}\right)\ .$$

TOME 8 (2025)

By monotonicity of generating functions ([GKPS21, Proposition 2.23]) we can assume that  $\frac{df_t}{dt} \leq 0$  for all  $t \in [\frac{1}{2}, 1]$ , and so

$$\left\{f_{\frac{1}{2}\left(\frac{T}{T_{0}+\epsilon'}+1\right)}\leqslant0\right\}\subset\left\{f_{\frac{1}{2}\left(\frac{T'}{T_{0}+\epsilon'}+1\right)}\leqslant0\right\}$$

for  $T, T' \in [T_0, T_0 + \epsilon')$  with  $T \leq T'$ . By continuity of the cohomological index, there is a neighborhood  $\mathcal{U}$  of  $\{f_{\frac{1}{2}(\frac{T_0}{T_0 + \epsilon'} + 1)} \leq 0\}$  such that if

$$\left\{f_{\frac{1}{2}\left(\frac{T_{0}}{T_{0}+\epsilon'}+1\right)}\leqslant0\right\}\subset\mathcal{V}\subset\mathcal{U}$$

then

$$\operatorname{ind}(\mathcal{V}) = \operatorname{ind}\left(\left\{f_{\frac{1}{2}\left(\frac{T_0}{T_0+\epsilon'}+1\right)} \leqslant 0\right\}\right).$$

For every  $T \in [T_0, T_0 + \epsilon)$  with  $\epsilon < \epsilon'$  small enough we have

$$\left\{f_{\frac{1}{2}\left(\frac{T_0}{T_0+\epsilon'}+1\right)} \leqslant 0\right\} \subset \left\{f_{\frac{1}{2}\left(\frac{T}{T_0+\epsilon'}+1\right)} \leqslant 0\right\} \subset \mathcal{U},$$

and so

$$\operatorname{ind}\left(\left\{f_{\frac{1}{2}\left(\frac{T}{T_{0}+\epsilon'}+1\right)}\leqslant 0\right\}\right) = \operatorname{ind}\left(\left\{f_{\frac{1}{2}\left(\frac{T_{0}}{T_{0}+\epsilon'}+1\right)}\leqslant 0\right\}\right),$$

i.e.  $\mu(\widetilde{r_{-T}} \cdot \widetilde{\phi}) = \mu(\widetilde{r_{-T_0}} \cdot \widetilde{\phi})$  as we wanted.

It is proved in [GKPS21, Proposition 4.21] that if  $\{\phi_t\}_{t \in [0,1)}$  is a non-negative (respectively non-positive) contact isotopy then  $\mu([\{\phi_t\}_{t \in [0,1)}]) \ge 0$  (respectively  $\mu([\{\phi_t\}_{t \in [0,1)}]) \le 0$ ). We actually have the following result.

PROPOSITION 2.6. — If  $\tilde{\phi} \leq \tilde{\psi}$  then  $\mu(\tilde{\phi}) \leq \mu(\tilde{\psi})$ .

*Proof.* — Assume that  $\tilde{\phi} \leq \tilde{\psi}$ . Then  $\tilde{\psi}$  can be represented by the concatenation

$$\{\psi_t\}_{t \in [0,1]} = \{\phi_{2t}\}_{t \in [0,\frac{1}{2}]} \sqcup \{\chi_{2t-1}\}_{t \in [\frac{1}{2},1]}$$

of a contact isotopy  $\{\phi_t\}_{t \in [0,1]}$  representing  $\tilde{\phi}$  and a non-negative contact isotopy  $\{\chi_t\}_{t \in [0,1]}$ . Let  $F_t : \mathbb{R}^{2n} \times \mathbb{R}^{2nN} \to \mathbb{R}$  be a based family of generating functions for  $\{\psi_t\}_{t \in [0,1]}$ . By monotonicity of generating function ([GKPS21, Proposition 2.23]) we can assume that  $F_1 \ge F_{\frac{1}{2}}$  and so, by monotonicity of the cohomological index ([GKPS21, Proposition 3.9(i)]),  $\operatorname{ind}(F_1) \le \operatorname{ind}(F_{\frac{1}{2}})$ . We thus have

$$\mu(\tilde{\psi}) = \mu(\tilde{\phi}) + \operatorname{ind}\left(F_{\frac{1}{2}}\right) - \operatorname{ind}(F_{1}) \ge \mu(\tilde{\phi}).$$

Remark 2.7. — It follows from Proposition 2.5 and Proposition 2.6 that the map  $T \mapsto \mu(\widetilde{r_{-T}} \cdot \widetilde{\phi})$  is lower semi-continuous, i.e.  $\{T \in \mathbb{R} \mid \mu(\widetilde{r_{-T}} \cdot \widetilde{\phi}) \leq y\}$  is closed for every  $y \in \mathbb{R}$ .

If  $F_t$  is a based family of conical generating functions for a contact isotopy  $\{\phi_t\}_{t \in [0,1]}$ of  $(L_k^{2n-1}, \xi_0)$  then for every t there is a 1–1 correspondence between the critical points of critical value zero of  $f_t$  and the discriminant points of  $\phi_t$  that lift to discriminant points of  $\overline{\phi}_t$ , where  $\{\overline{\phi}_t\}_{t \in [0,1]}$  is the lift of  $\{\phi_t\}_{t \in [0,1]}$  to  $(\mathbb{S}^{2n-1}, \overline{\xi}_0)$ . Since the nonlinear Maslov index  $\mu(\{\phi_t\}_{t \in [0,1]})$  counts, with multiplicity given by the change in the cohomological index of the sublevel sets  $\{f_t \leq 0\}$ , the critical points of  $f_t$  with critical value zero as t varies in [0, 1], its value is related to the presence of discriminant points of  $\overline{\phi}_t$  for  $t \in [0, 1]$ . More precisely, we have the following result ([GKPS21, Theorem 1.4(iii)]).

PROPOSITION 2.8. — Let  $\{\phi_t\}_{t \in [0,1]}$  be a contact isotopy of  $(L_k^{2n-1}, \xi_0)$  starting at the identity, and let  $[t_0, t_1]$  be a subinterval of [0, 1]. If  $\mu(\{\phi_t\}_{t \in [0,t_0]}) \neq \mu(\{\phi_t\}_{t \in [0,t_1]})$ then there is  $\underline{t} \in [t_0, t_1]$  such that  $\overline{\phi}_{\underline{t}}$  has discriminant points. If moreover<sup>(1)</sup> the map  $s \mapsto \mu(\{\phi_t\}_{t \in [0,s]})$  is constant on  $[t_0, \underline{t})$  and on  $(\underline{t}, t_1]$  and all the discriminant points of  $\overline{\phi}_t$  are non-degenerate then

$$\left| \mu\left( \{\phi_t\}_{t \in [0,t_0]} \right) - \mu\left( \{\phi_t\}_{t \in [0,t_1]} \right) \right| \leq 1$$

In particular, it follows from Example 2.2, Proposition 2.4 and the first statement of Proposition 2.8 that for every real number T we have

(2.4) 
$$\mu(\widetilde{r_T}) = 2n \left\lceil \frac{T}{2\pi} \right\rceil$$

We also have the following result.

PROPOSITION 2.9. — Let  $\{\phi_t\}_{t \in [0,1]}$  be a contact isotopy of  $(L_k^{2n-1}, \xi_0)$  starting at the identity, and let  $[t_0, t_1]$  be a subinterval of [0, 1]. Assume that there is  $\underline{t} \in [t_0, t_1]$ such that  $\overline{\phi}_{\underline{t}}$  has discriminant points, and denote by  $\Delta(\phi_{\underline{t}}) \subset L_k^{2n-1}$  the set of discriminant points of  $\phi_{\underline{t}}$ . Assume also that the map  $s \mapsto \mu(\{\phi_t\}_{t \in [0,s]})$  is constant on  $[t_0, \underline{t}]$  and on  $(\underline{t}, t_1]$ . Then

$$\mu\left(\{\phi_t\}_{t\in[0,t_0]}\right) - \mu\left(\{\phi_t\}_{t\in[0,t_1]}\right) \Big| \leqslant \operatorname{ind}\left(\Delta(\phi_{\underline{t}})\right) + 1\,,$$

and

$$\left| \mu\left(\{\phi_t\}_{t \in [0,t_0]}\right) - \mu\left(\{\phi_t\}_{t \in [0,t_1]}\right) \right| \leq \operatorname{ind}\left(\Delta(\phi_{\underline{t}})\right)$$

if either k = 2 or if  $\{\phi_t\}_{t \in [0,1]}$  is negative and  $\mu(\{\phi_t\}_{t \in [0,t_0]})$  is even.

Before proving Proposition 2.9, recall that the cohomological index is subadditive in the following sense ([GKPS21, Proposition 3.9(iv)]): for any two closed subsets A and B of  $L_k^{2M-1}$  we have

$$\operatorname{ind}(A \cup B) \leq \operatorname{ind}(A) + \operatorname{ind}(B) + 1$$
,

and

$$\operatorname{ind}(A \cup B) \leq \operatorname{ind}(A) + \operatorname{ind}(B)$$

if either k = 2 or ind(A) is even or ind(B) is even.

<sup>&</sup>lt;sup>(1)</sup> In [GKPS21, Theorem 1.4(iii)] it is assumed that there is only one  $\underline{t} \in [0, 1]$  such that  $\bar{\phi}_{\underline{t}}$  has discriminant points. However, the proof works in the same way also if we only assume that  $s \mapsto \mu(\{\phi_t\}_{t \in [0,s]})$  is constant on  $[t_0, \underline{t})$  and on  $(\underline{t}, t_1]$ . Similarly for Proposition 2.9 below.

Proof of Proposition 2.9. — The first inequality and the second one in the case k = 2 are proved in [GKPS21, Proposition 4.15]. Suppose thus that  $\{\phi_t\}_{t\in[0,1]}$  is negative and  $\mu(\{\phi_t\}_{t\in[0,t_0]})$  is even. By monotonicity of generating functions ([GKPS21, Proposition 2.23]), there is a based family of conical generating functions  $F_t$  for  $\{\phi_t\}_{t\in[0,1]}$  with  $\frac{df_t}{dt} \leq 0$ . The proof of [GKPS21, Proposition 2.23] actually shows that for every t either  $\frac{df_t}{dt} < 0$  or  $\frac{df_t}{dt} = 0$ . We can thus assume that either  $\{f_{t_0} \leq 0\} = \{f_{\underline{t}} \leq 0\}$  or  $\{f_{t_0} \leq 0\}$  is included in the interior of  $\{f_{\underline{t}} \leq 0\}$ . In the first case, by continuity of the cohomological index and since the map  $s \mapsto \mu(\{\phi_t\}_{t\in[0,s]})$  is constant on  $(\underline{t}, t_1]$  we have  $\mu(\{\phi_t\}_{t\in[0,t_0]}) = \mu(\{\phi_t\}_{t\in[0,t_1]})$ , which implies the desired inequality

$$\left| \mu\left(\{\phi_t\}_{t \in [0,t_0]}\right) - \mu\left(\{\phi_t\}_{t \in [0,t_1]}\right) \right| \leq \operatorname{ind}\left(\Delta(\phi_{\underline{t}})\right).$$

In the second case, take  $\epsilon, \epsilon' > 0$  such that

$$\{f_{t_0} \leqslant 0\} \subset \{f_{\underline{t}} \leqslant -\epsilon\} \subset \{f_{\underline{t}-\epsilon'} \leqslant 0\} \subset \{f_{\underline{t}} \leqslant 0\}.$$

By monotonicity of the cohomological index and since  $s \mapsto \mu(\{\phi_t\}_{t \in [0,s]})$  is constant on  $[t_0, \underline{t})$  we then have

(2.5) 
$$\operatorname{ind}\left(\left\{f_{\underline{t}} \leqslant -\epsilon\right\}\right) = \operatorname{ind}\left(\left\{f_{t_0} \leqslant 0\right\}\right).$$

As in the proof of [GKPS21, Proposition 4.15] we have

$$(2.6) \qquad \left| \mu\left(\{\phi_t\}_{t \in [0,t_0]}\right) - \mu\left(\{\phi_t\}_{t \in [0,t_1]}\right) \right| \leq \operatorname{ind}\left(\{f_{\underline{t}} \leq \epsilon\}\right) - \operatorname{ind}\left(\{f_{\underline{t}} \leq -\epsilon\}\right)$$

and

(2.7) 
$$\operatorname{ind}\left(\{f_{\underline{t}} \leqslant \epsilon\}\right) \leqslant \operatorname{ind}\left(\{f_{\underline{t}} \leqslant -\epsilon\} \cup \mathcal{U}\right),$$

where  $\mathcal{U}$  is a neighborhood of the set C of critical points of  $f_{\underline{t}}$  of critical value zero that has the same cohomological index as C. Since

$$\mu\left(\{\phi_t\}_{t \in [0,t_0]}\right) = \operatorname{ind}\left(\{f_0 \leqslant 0\}\right) - \operatorname{ind}\left(\{f_{t_0} \leqslant 0\}\right)$$

is even (by assumption) and  $\operatorname{ind}(\{f_0 \leq 0\})$  is even (by [GKPS21, Remark 3.13]), using (2.5) we see that  $\operatorname{ind}(\{f_{\underline{t}} \leq -\epsilon\})$  is even and so, by subadditivity of the cohomological index,

$$\operatorname{ind}\left(\{f_{\underline{t}} \leqslant -\epsilon\} \cup \mathcal{U}\right) \leqslant \operatorname{ind}\left(\{f_{\underline{t}} \leqslant -\epsilon\}\right) + \operatorname{ind}(\mathcal{U}) = \operatorname{ind}\left(\{f_{\underline{t}} \leqslant -\epsilon\}\right) + \operatorname{ind}(C).$$

Since  $\operatorname{ind}(C) = \operatorname{ind}(\Delta(\phi_{\underline{t}}))$  by [GKPS21, Proposition 2.22], using (2.6) and (2.7) we thus conclude that

$$\mu\left(\{\phi_t\}_{t\in[0,t_0]}\right) - \mu\left(\{\phi_t\}_{t\in[0,t_1]}\right) \Big| \leq \operatorname{ind}\left(\Delta(\phi_{\underline{t}})\right). \qquad \Box$$

The following Poincaré duality property for the cohomological index is proved in [Car13, Proposition 4.1.15] for subsets of complex projective space. We adapt here the proof to the case of lens spaces.

LEMMA 2.10. — Assume that zero is a regular value of a function  $f: L_k^{2M-1} \to \mathbb{R}$ . Then

$$\operatorname{ind}\left(\{f \leq 0\}\right) + \operatorname{ind}\left(\{f \geq 0\}\right) = 2M.$$

Proof. — Let  $A = \{f \leq 0\}$  and  $B = \{f \geq 0\}$ . Since zero is a regular value of f, A and B are smooth submanifolds with boundary and thus deformation retract to sets A' and B' that are strictly included in  $\{f < 0\}$  and  $\{f > 0\}$  respectively. By continuity of the cohomological index, there are thus open subsets  $\mathcal{U}_A$  and  $\mathcal{U}_B$  of  $L_k^{2M-1}$  strictly included in  $\{f < 0\}$  and  $\{f > 0\}$  respectively with  $\operatorname{ind}(\mathcal{U}_A) = \operatorname{ind}(A)$  and  $\operatorname{ind}(\mathcal{U}_B) = \operatorname{ind}(B)$ . Assume first that  $\operatorname{ind}(A)$  is even. Since  $A \cup B = L_k^{2M-1}$ , by subadditivity of the cohomological index we have

$$\operatorname{ind}(A) + \operatorname{ind}(B) \ge 2M$$
.

Assume by contradiction that

(2.8) 
$$\operatorname{ind}(\mathcal{U}_A) + \operatorname{ind}(\mathcal{U}_B) = \operatorname{ind}(A) + \operatorname{ind}(B) \ge 2M + 1$$

Let  $\operatorname{ind}(A) = \operatorname{ind}(\mathcal{U}_A) = 2a$ . By definition of the cohomological index (cf. [GKPS21, Lemma 3.3]) and since Čech cohomology agrees with singular cohomology on open sets, the homomorphism  $H^{2a-1}(L_k^{2M-1};\mathbb{Z}_k) \to H^{2a-1}(\mathcal{U}_A;\mathbb{Z}_k)$  induced by the inclusion  $\mathcal{U}_A \hookrightarrow L_k^{2M-1}$  is injective. Since k is prime, the coefficient ring  $\mathbb{Z}_k$  is a field and so cohomology is the dual of homology, thus the homomorphism  $H_{2a-1}(\mathcal{U}_A;\mathbb{Z}_k) \to$  $H_{2a-1}(L_k^{2M-1};\mathbb{Z}_k)$  induced by the inclusion  $\mathcal{U}_A \hookrightarrow L_k^{2M-1}$  is surjective. Consider the commutative square

(2.9) 
$$\begin{array}{c} H_{2a-1}\left(L_{k}^{2M-1} \smallsetminus \operatorname{int}(B); \mathbb{Z}_{k}\right) \longrightarrow H_{2a-1}\left(L_{k}^{2M-1}; \mathbb{Z}_{k}\right) \\ \cong \uparrow \qquad \cong \uparrow \qquad \cong \uparrow \\ H^{2(M-a)}\left(L_{k}^{2M-1}, B; \mathbb{Z}_{k}\right) \xrightarrow{j_{B}} H^{2(M-a)}\left(L_{k}^{2M-1}; \mathbb{Z}_{k}\right) \end{array}$$

where the horizontal arrows are induced by the inclusions and the vertical ones are Poincaré duality isomorphisms (composed with excision  $H^{2(M-a)}(L_k^{2M-1}, B; \mathbb{Z}_k) \to$  $H^{2(M-a)}(L_k^{2M-1} \setminus \operatorname{int}(B), \partial B; \mathbb{Z}_k)$  for the vertical arrow on the left hand side). Since  $\mathcal{U}_A \subset L_k^{2M-1} \setminus \operatorname{int}(B)$ , surjectivity of the inclusion homomorphism  $H_{2a-1}(\mathcal{U}_A; \mathbb{Z}_k) \to$  $H_{2a-1}(L_k^{2M-1}; \mathbb{Z}_k)$  implies that the homomorphism on the top horizontal line of the diagram is also surjective. Thus  $j_B$  is surjective, and so there exists a class

$$u \in H^{2(M-a)}\left(L_k^{2M-1}, B; \mathbb{Z}_k\right)$$

such that  $j_B(u) = \beta^{M-a}$ , where  $\beta$  denotes a generator of  $H^2(L_k^{2M-1}; \mathbb{Z}_k)$ . By (2.8) we have  $\operatorname{ind}(\mathcal{U}_B) \ge 2M + 1 - 2a$ , and so by a similar argument there exists a class

$$v \in H^{2a-1}\left(L_k^{2M-1}, A; \mathbb{Z}_k\right)$$

such that  $j_A(v) = \alpha \beta^{a-1}$ , where

$$j_A: H_{2a-1}\left(L_k^{2M-1}, A; \mathbb{Z}_k\right) \to H_{2a-1}\left(L_k^{2M-1}; \mathbb{Z}_k\right)$$

is the homomorphism induced by the inclusion and where  $\alpha$  denotes a generator of  $H^1(L_k^{2M-1}; \mathbb{Z}_k)$ . We then obtain a contradiction: on the one hand  $v \cup u \in$  $H^{2M-1}(L_k^{2M-1}, A \cup B; \mathbb{Z}_k)$  is zero since  $A \cup B = L_k^{2M-1}$ , on the other hand by naturality of the cup product we have

$$j_{A\cup B}(v\cup u) = j_A(v)\cup j_B(u) = \alpha\beta^{M-1} \neq 0.$$

TOME 8 (2025)

This finishes the proof in the case when ind(A) is even. If ind(B) is even a similar argument also gives the result. Suppose thus that ind(A) and ind(B) are odd. Then ind(A) + ind(B) is even, and so it is enough to prove

$$2M - 1 \leq \operatorname{ind}(A) + \operatorname{ind}(B) \leq 2M$$
.

The first inequality follows from subadditivity of the cohomological index. For the second one, suppose by contradiction that

$$\operatorname{ind}(\mathcal{U}_A) + \operatorname{ind}(\mathcal{U}_B) = \operatorname{ind}(A) + \operatorname{ind}(B) \ge 2M + 1$$
,

and let  $\operatorname{ind}(A) = 2a - 1$ . By a similar argument as above, there exist cohomology classes u in  $H^{2M-2a+1}(L_k^{2M-1}, B; \mathbb{Z}_k)$  and v in  $H^{2a-2}(L_k^{2M-1}, A; \mathbb{Z}_k)$  such that  $j_B(u) = \alpha \beta^{M-a}$  and  $j_A(v) = \beta^{a-1}$ . Since  $j_B(u) \cup j_A(v) = \alpha \beta^{M-1} \neq 0$ , this leads to the same contradiction as above.

Applying Lemma 2.10 we obtain the following result.

PROPOSITION 2.11. — For every  $\tilde{\phi} \in \widetilde{\text{Cont}}_0(L_k^{2n-1}, \xi_0)$  such that  $\Pi(\mathcal{L}(\tilde{\phi}))$  does not have discriminant points we have

$$\mu(\widetilde{\phi}) + \mu(\widetilde{\phi}^{-1}) = 2n \,.$$

Proof. — Let  $F_t : \mathbb{R}^{2n} \times \mathbb{R}^{2nN} \to \mathbb{R}$  be a based family of conical generating functions for a contact isotopy  $\{\phi_t\}_{t \in [0,1]}$  representing  $\tilde{\phi}$ . Then  $-F_t$  is a based family of conical generating functions for  $\{\phi_t^{-1}\}_{t \in [0,1]}$ , which represent  $\tilde{\phi}^{-1}$ . Thus

$$\mu(\widetilde{\phi}) + \mu(\widetilde{\phi}^{-1}) = \operatorname{ind}(F_0) - \operatorname{ind}(F_1) + \operatorname{ind}(-F_0) - \operatorname{ind}(-F_1).$$

Since  $F_0$  is equivalent to the zero function, up to a fibre preserving conical homeomorphism it is equal to a  $\mathbb{Z}_k$ -invariant quadratic form  $Q_0$ . Since  $F_0$  generates the identity, by [GKPS21, Proposition 2.22] the nullity of  $Q_0$  is equal to 2n. Let  $\iota$  be the dimension of the maximal subspace on which  $Q_0$  is negative definite. Then

$$\operatorname{ind}(F_0) + \operatorname{ind}(-F_0) = \operatorname{ind}(Q_0) + \operatorname{ind}(-Q_0) = (2n+\iota) + (2n+(2nN-\iota)) = 4n+2nN.$$

On the other hand, since  $\Pi(\mathcal{L}(\tilde{\phi}))$  has no discriminant points we have that zero is a regular value of  $F_1$  and so we can apply Lemma 2.10 to obtain

$$\operatorname{ind}(F_1) + \operatorname{ind}(-F_1) = 2n + 2nN$$

We conclude that  $\mu(\tilde{\phi}) + \mu(\tilde{\phi}^{-1}) = 2n$ .

So far we have assumed that k is prime. Suppose now that k is not prime, and let k' be the smallest prime that divides k. As in [GKPS21, Remark 1.4], we define the non-linear Maslov index

$$\mu : \widetilde{\operatorname{Cont}_0}\left(L_k^{2n-1}, \xi_0\right) \to \mathbb{Z}$$

by pulling back  $\mu : \widetilde{\operatorname{Cont}}_0(L_{k'}^{2n-1},\xi_0) \to \mathbb{Z}$  by the natural map  $\widetilde{\operatorname{Cont}}_0(L_k^{2n-1},\xi_0) \to \widetilde{\operatorname{Cont}}_0(L_{k'}^{2n-1},\xi_0)$ . This general non-linear Maslov index then satisfies the following properties.

PROPOSITION 2.12 (Non-linear Maslov index for general k). — For any integer  $k \ge 2$  and n-tuple  $\underline{w} = (w_1, \dots, w_n)$  of positive integers relatively prime to k, the non-linear Maslov index

$$\mu: \widetilde{\operatorname{Cont}}_0\left(L_k^{2n-1}(\underline{w}), \xi_0\right) \to \mathbb{Z}$$

satisfies the following properties:

- (i) Identity and small elements: if  $\tilde{\phi}$  is small enough in the  $C^1$ -topology then  $0 \leq \mu(\tilde{\phi}) \leq 2n$ , and if moreover  $\tilde{\phi}$  is positive then  $\mu(\tilde{\phi}) = 2n$ . Moreover,  $\mu(\tilde{id}) = 0$ .
- (ii) Quasimorphism property:

$$\mu(\widetilde{\phi}\cdot\widetilde{\psi})-\mu(\widetilde{\phi})-\mu(\widetilde{\psi}) \mid \leq 2n+1 \,,$$

and

$$\left| \mu(\widetilde{\phi} \cdot \widetilde{\psi}) - \mu(\widetilde{\phi}) - \mu(\widetilde{\psi}) \right| \leq 2n$$

if k is even; in particular,  $\mu$  is a quasimorphism.

(iii) Triangle inequality:

$$\mu(\tilde{\phi} \cdot \tilde{\psi}) \leqslant \mu(\tilde{\phi}) + \mu(\tilde{\psi}) + 1$$

and

$$\mu(\widetilde{\phi} \cdot \widetilde{\psi}) \leqslant \mu(\widetilde{\phi}) + \mu(\widetilde{\psi})$$

if either k is even or  $\mu(\tilde{\phi})$  is even or  $\mu(\tilde{\psi})$  is even.

- (iv) Monotonicity: if  $\tilde{\phi} \leq \tilde{\psi}$  then  $\mu(\tilde{\phi}) \leq \mu(\tilde{\psi})$ .
- (v) Relation with discriminant points: for any contact isotopy  $\{\phi_t\}_{t \in [0,1]}$  starting at the identity and any subinterval  $[t_0, t_1]$  of [0, 1], if  $\mu(\{\phi_t\}_{t \in [0, t_0]}) \neq \mu(\{\phi_t\}_{t \in [0, t_1]})$  then there is  $\underline{t} \in [t_0, t_1]$  such that  $\overline{\phi}_{\underline{t}}$  has discriminant points. Assume that the map  $s \mapsto \mu(\{\phi_t\}_{t \in [0,s]})$  is constant on  $[t_0, \underline{t})$  and on  $(\underline{t}, t_1]$ , and denote by  $\Delta_{k'}(\phi_{\underline{t}}) \subset L_{k'}^{2n-1}(\underline{w})$  the set of discriminant points at time  $\underline{t}$  of the lift of  $\{\phi_t\}$  to  $L_{k'}^{2n-1}(\underline{w})$ , where k' is the smallest prime dividing k. Then

$$\left| \mu(\{\phi_t\}_{t \in [0,t_0]}) - \mu(\{\phi_t\}_{t \in [0,t_1]}) \right| \leq \operatorname{ind} \left( \Delta_{k'}(\phi_{\underline{t}}) \right) + 1,$$

and

$$\left| \mu(\{\phi_t\}_{t \in [0,t_0]}) - \mu(\{\phi_t\}_{t \in [0,t_1]}) \right| \leq \operatorname{ind} \left( \Delta_{k'}(\phi_{\underline{t}}) \right)$$

if either k is even or if  $\{\phi_t\}_{t\in[0,1]}$  is negative and  $\mu(\{\phi_t\}_{t\in[0,t_0]})$  is even. If moreover all the discriminant points of  $\overline{\phi}_{\underline{t}}$  are non-degenerate then

$$\left| \mu(\{\phi_t\}_{t \in [0,t_0]}) - \mu(\{\phi_t\}_{t \in [0,t_1]}) \right| \leq 1.$$

(vi) Reeb flow: for every real number T we have

$$\mu(\widetilde{r_T}) = 2n \left\lceil \frac{T}{2\pi} \right\rceil$$

(vii) Composition with the Reeb flow: for every integer m we have

$$\mu\left(\widetilde{\phi} \cdot \widetilde{r_{2\pi m}}\right) = \mu\left(\widetilde{r_{2\pi m}} \cdot \widetilde{\phi}\right) = 2nm + \mu(\widetilde{\phi})$$

- (viii) Lower semi-continuity: the map  $T \mapsto \mu(\widetilde{r_{-T}} \cdot \widetilde{\phi})$  is lower semi-continuous.
  - (ix) Poincaré duality: for every  $\tilde{\phi}$  such that  $\Pi(\mathcal{L}(\tilde{\phi}))$  has no discriminant points we have

$$\mu(\widetilde{\phi}) + \mu(\widetilde{\phi}^{-1}) = 2n \,.$$

Proof. — In the case when k is prime all the properties have been discussed above in Example 2.2, Equations (2.1) and (2.2), Proposition 2.3, Proposition 2.6, Proposition 2.8, Proposition 2.9, Equation (2.4), Proposition 2.4, Remark 2.7 and Proposition 2.11. If k is not prime and k' is the smallest prime that divides k then the properties of  $\mu : Cont_0(L_{k'}^{2n-1}(\underline{w}), \xi_0) \to \mathbb{Z}$  imply the corresponding properties for the pullback  $\mu : Cont_0(L_k^{2n-1}(\underline{w}), \xi_0) \to \mathbb{Z}$  by the natural map  $Cont_0(L_k^{2n-1}(\underline{w}), \xi_0) \to \mathbb{Z}$  by the natural map  $Cont_0(L_k^{2n-1}(\underline{w}), \xi_0) \to \mathbb{Z}$  by the natural map  $L_k^{2n-1}(\underline{w}), \xi_0 \to \mathbb{Z}$  by the natural map  $L_k^{2n-1}(\underline{w}), \xi_0$  be the natural map  $L_k^{2n-1}(\underline{w}), \xi_0$ 

#### **3.** Spectral selectors

For any  $j \in \mathbb{Z}$  we define the *j*-th spectral selector on  $\widetilde{\mathrm{Cont}}_0(L_k^{2n-1}(\underline{w}),\xi_0)$  by

$$c_j(\widetilde{\phi}) = \inf \left\{ T \in \mathbb{R} \mid \mu(\widetilde{r_{-T}} \cdot \widetilde{\phi}) \leqslant -j \right\}.$$

By Proposition 2.12 (iv), (ii) and (vi), the function  $T \mapsto \mu(\widetilde{r}_{-T} \cdot \phi)$  is non-increasing and tends to  $\mp \infty$  as  $T \to \pm \infty$ , thus  $c_j(\phi)$  is a well-defined real number. By Proposition 2.12 (viii) the infimum is in fact a minimum, in particular

(3.1) 
$$\left[c_{j}(\widetilde{\phi})\right]_{T_{\underline{w}}} = \min\left\{N \in T_{\underline{w}} \cdot \mathbb{Z} \mid \mu(\widetilde{r_{-N}} \cdot \widetilde{\phi}) \leqslant -j\right\}.$$

It follows from the definition that the sequence  $\{c_j\}$  is non-decreasing. In the rest of this section we prove the other properties listed in Theorem 1.1. We say that a contactomorphism of  $(L_k^{2n-1}(\underline{w}), \xi_0)$  is non-degenerate with respect

We say that a contactomorphism of  $(L_k^{2n-1}(\underline{w}), \xi_0)$  is non-degenerate with respect to  $\alpha_0$  if all its translated points with respect to  $\alpha_0$  are non-degenerate for all their translations. We then have the following lemma.

LEMMA 3.1. — The set of contactomorphisms contact isotopic to the identity that are non-degenerate with respect to  $\alpha_0$  is dense in  $\operatorname{Cont}_0(L_k^{2n-1}(\underline{w}), \xi_0)$  for the  $\mathcal{C}^1$ -topology.

Proof. — We first prove the following result. Let  $\Lambda_0$  be a closed Legendrian submanifold of a contact manifold  $(M, \xi = \ker(\alpha))$ , and denote by  $\mathcal{L}eg(\Lambda_0)$  its Legendrian isotopy class. Then for any collection  $N = \{N_l, l \in \mathbb{Z}\}$  of submanifolds of M the set  $\mathcal{L}eg_N(\Lambda_0)$  of elements of  $\mathcal{L}eg(\Lambda_0)$  transverse to  $N_l$  for all l is dense in  $\mathcal{L}eg(\Lambda_0)$  for the  $\mathcal{C}^1$ -topology. Indeed, let  $\Lambda$  be an element of  $\mathcal{L}eg(\Lambda_0)$  and let  $\mathcal{U}(\Lambda) \subset M$  be a Weinstein neighborhood of  $\Lambda$ . Fix a diffeomorphism  $\Psi$  from  $\mathcal{U}(\Lambda)$  to an open neighborhood  $\mathcal{U}(j^{10})$  of the zero section of  $J^1\Lambda$  such that  $\Psi(\Lambda) = j^{10}$  and  $\Psi^*(dz - \lambda_{can}) = \alpha$ . Denote for each l by  $N'_l$  the submanifold  $\Psi(N_l \cap \mathcal{U}(\Lambda))$  of  $\mathcal{U}(j^{10})$ , and let  $N' = \{N'_l, l \in \mathbb{Z}\}$ . Since the map  $j^1 : \mathcal{C}^{\infty}(\Lambda) \to \mathcal{L}eg(j^{10})$  that associates to a function its 1-jet is a local homeomorphism with respect to the  $C^2$ -topology on  $\mathcal{C}^{\infty}(\Lambda)$  and the  $\mathcal{C}^1$ -topology on  $\mathcal{L}eg(j^{10})$  (cf. for instance [Tsu08, Section 3]), for a sufficiently small  $\mathcal{C}^2$ -neighborhood  $\mathcal{U}(0)$  of the zero function in  $\mathcal{C}^{\infty}(\Lambda)$  the 1-jet of any  $f \in \mathcal{U}(0)$  is in  $\mathcal{U}(j^{1}0)$  and the map  $\mathcal{U}(0) \to \mathcal{L}eg(\Lambda_0)$  that sends f to  $\Psi^{-1}(j^{1}f)$  is a local homeomorphism. By Thom's transversality theorem (see for instance [GG73, Corollary 4.10]), the subset  $T_{N'}^1$  of  $\mathcal{U}(0)$  consisting of functions with 1-jet transverse to  $N'_l$  for all l is dense in  $\mathcal{U}(0)$  for the  $\mathcal{C}^{\infty}$ -topology, hence also for the  $\mathcal{C}^2$ -topology. Thus for any  $\mathcal{C}^1$ -neighborhood  $\mathcal{U}$  of  $\Lambda$  in  $\mathcal{L}eg(\Lambda_0)$  there exists  $f \in T_{N'}^1$  such that  $\Psi^{-1}(j^1f) \in \mathcal{U}$ . The Legendrian  $\Psi^{-1}(j^1f)$  intersects each  $N_l$  transversely, and so belongs to  $\mathcal{L}eg_N(\Lambda_0)$ . This shows that  $\mathcal{L}eg_N(\Lambda_0)$  is  $\mathcal{C}^1$ -dense in  $\mathcal{L}eg(\Lambda_0)$ .

Using this, we now prove that the set of contactomorphisms of  $(L_k^{2n-1}(\underline{w}), \xi_0)$ contact isotopic to the identity that are non-degenerate with respect to  $\alpha_0$  is  $\mathcal{C}^{1}$ dense in  $\operatorname{Cont}_0(L_k^{2n-1}(\underline{w}), \xi_0)$ . Consider the contact product of  $(L_k^{2n-1}(\underline{w}), \xi_0)$ , i.e. the product  $L_k^{2n-1}(\underline{w}) \times L_k^{2n-1}(\underline{w}) \times \mathbb{R}$  endowed with the contact structure given by the kernel of the contact form  $\pi_2^* \alpha_0 - e^{\theta} \pi_1^* \alpha_0$ , where  $\pi_1$  and  $\pi_2$  denote the projections on the first and second factor respectively and where  $\theta$  is the coordinate in  $\mathbb{R}$ . Denote by  $\operatorname{gr}_{\alpha_0}(\phi)$  the graph with respect to  $\alpha_0$  of a contactomorphism  $\phi$  of  $(L_k^{2n-1}(\underline{w}), \xi_0)$ , i.e. the Legendrian submanifold

$$\operatorname{gr}_{\alpha_0}(\phi) = \left\{ \left( p, \phi(p), g(p) \right) \mid p \in L_k^{2n-1}(\underline{w}) \right\}$$

of the contact product, where g is the conformal factor of  $\phi$  with respect to  $\alpha_0$ . Denote by  $\{R_t\}$  the Reeb flow of  $\pi_2^* \alpha_0 - e^{\theta} \pi_1^* \alpha_0$ . Let  $0 = t_0 < \cdots < t_m = 2\pi$  be a decomposition of the time interval  $[0, 2\pi]$  such that, for some  $\epsilon > 0$ ,  $N_l := \bigcup_{t \in (t_l - \epsilon, t_{l+1} + \epsilon)} R_t(\operatorname{gr}_{\alpha_0}(\operatorname{id}))$  is a submanifold of  $L_k^{2n-1}(\underline{w}) \times L_k^{2n-1}(\underline{w}) \times \mathbb{R}$  for every  $l \in \{0, \cdots, m-1\}$ . A contactomorphism  $\phi$  of  $(L_k^{2n-1}(\underline{w}), \xi_0)$  is non-degenerate with respect to  $\alpha_0$  if and only if  $\operatorname{gr}_{\alpha_0}(\phi)$  is transverse to  $N_l$  for all l. In other words, using the notation of the first part of the proof,  $\phi$  is non-degenerate with respect to  $\alpha_0$  if and only if  $\operatorname{gr}_{\alpha_0}(\operatorname{id})$ , where  $N = \{N_0, \cdots, N_{m-1}\}$ . Our result thus follows from the first part of the proof and the fact that the map

$$\operatorname{gr}_{\alpha_0}:\operatorname{Cont}_0\left(L_k^{2n-1}(\underline{w}),\xi_0\right)\to \mathcal{L}eg(\operatorname{gr}_{\alpha_0}(\operatorname{id}))$$

that associates to a contactomorphism its graph is a local homeomorphism with respect to the  $C^1$ -topologies.

We also remark the following fact.

LEMMA 3.2. — For any element  $\tilde{\phi}$  of  $\widetilde{\text{Cont}}_0(L_k^{2n-1}(\underline{w}), \xi_0)$ , the sets  $\mathcal{A}(\tilde{\phi})$  and  $\overline{\mathcal{A}}(\tilde{\phi})$  are closed and nowhere dense in  $\mathbb{R}$ .

Proof. — Let  $F_t : \mathbb{R}^{2n} \times \mathbb{R}^{2nN} \to \mathbb{R}, t \in [0,1]$ , be a based family of conical generating functions for the concatenation

$$\{\phi_{2t}\}_{t \in [0,\frac{1}{2}]} \sqcup \{r_{-2\pi(2t-1)}\}_{t \in [\frac{1}{2},1]}.$$

For every  $t \in [\frac{1}{2}, 1]$ , the  $\mathbb{Z}_k$ -orbits of translated points of  $\overline{\phi}_1$  of translation  $2\pi(2t-1)$  are in 1–1 correspondence with the critical points of critical value zero of  $f_t$ :  $L_k^{2n(N+1)-1} \to \mathbb{R}$ . Consider the function

$$f: L_k^{2n(N+1)-1} \times \left[\frac{1}{2}, 1\right] \to \mathbb{R}, \ (x, t) \mapsto f_t(x).$$

TOME 8 (2025)

As in [Thé98, Section 5.2 and Lemma 4.4], zero is a regular value of f and so  $f^{-1}(0)$  is a submanifold of  $L_k^{2n(N+1)-1} \times [\frac{1}{2}, 1]$ . Let  $p: f^{-1}(0) \to [\frac{1}{2}, 1]$  be the composition of the inclusion  $f^{-1}(0) \hookrightarrow L_k^{2n(N+1)-1} \times [\frac{1}{2}, 1]$  with the projection on the second factor. Then  $(x,t) \in f^{-1}(0)$  is a critical point of p (of critical value t) if and only if x is a critical point of  $f_t$  (of critical value zero). Thus the  $\mathbb{Z}_k$ -orbits of translated points of  $\bar{\phi}_1$  of translation  $2\pi(2t-1)$  are in 1–1 correspondence with the critical points of p of critical value t. It follows that

$$\bar{\mathcal{A}}(\tilde{\phi}) = 2\pi \Big( 2p \Big( \operatorname{Crit}(p) \Big) - 1 \Big) + 2\pi \cdot \mathbb{Z},$$

and so  $\overline{\mathcal{A}}(\phi)$  is closed and nowhere dense. Since

$$\bar{\mathcal{A}}(\tilde{\phi}) + T_{\underline{w}} \cdot \mathbb{Z} \subseteq \mathcal{A}(\tilde{\phi}) \subseteq \bar{\mathcal{A}}(\tilde{\phi}) + \frac{2\pi}{k} \cdot \mathbb{Z}$$

we deduce that  $\mathcal{A}(\tilde{\phi})$  is also closed and nowhere dense.

We now prove that the spectral selectors satisfy the properties listed in Theorem 1.1.

#### Spectrality

We have to show that

$$c_j(\widetilde{\phi}) \in \bar{\mathcal{A}}(\widetilde{\phi})$$

for every  $\tilde{\phi}$  in  $\widetilde{\text{Cont}}_0(L_k^{2n-1}(\underline{w}), \xi_0)$ . Suppose by contradiction that this is not the case. Since  $\bar{\mathcal{A}}(\tilde{\phi})$  is closed (by Lemma 3.2), there is then  $\epsilon > 0$  such that  $[c_j(\tilde{\phi}) - \epsilon, c_j(\tilde{\phi}) + \epsilon] \subset \mathbb{R} \smallsetminus \bar{\mathcal{A}}(\tilde{\phi})$ . By the first statement of Proposition 2.12 (v) we thus have

$$\mu\left(\widetilde{r_{-(c_j(\widetilde{\phi})-\epsilon)}}\cdot\widetilde{\phi}\right) = \mu\left(\widetilde{r_{-(c_j(\widetilde{\phi})+\epsilon)}}\cdot\widetilde{\phi}\right),$$

but this contradicts the definition of  $c_j(\phi)$ .

#### Normalization

It follows from Proposition 2.12 (vi) that

$$c_{j}(\tilde{id}) = \begin{cases} \cdots \\ -4\pi & \text{for } j = -6n + 1, \cdots, -4n \\ -2\pi & \text{for } j = -4n + 1, \cdots, -2n \\ 0 & \text{for } j = -2n + 1, \cdots, 0 \\ 2\pi & \text{for } j = 1, \cdots, 2n \\ 4\pi & \text{for } j = 2n + 1, \cdots, 4n \\ \cdots \end{cases}$$

In particular,  $c_0(\tilde{id}) = 0$ .

#### Relation with translated points

If all the translated points of  $\Pi(\mathcal{L}(\tilde{\phi}))$  are non-degenerate, it follows from the last statement of Proposition 2.12(v) that the spectral selectors  $\{c_j(\tilde{\phi}), j \in \mathbb{Z}\}$  are all distinct. Suppose now that

$$c_{j-1}(\widetilde{\phi}) < c_j(\widetilde{\phi}) = c_{j+1}(\widetilde{\phi}) = \dots = c_{j+m}(\widetilde{\phi}) = T < c_{j+m+1}(\widetilde{\phi})$$

for some j and  $1 \leq m \leq 2n - 1$ . Then for  $\epsilon > 0$  small enough we have

$$\left| \mu \left( \widetilde{r_{-(T-\epsilon)}} \cdot \widetilde{\phi} \right) - \mu \left( \widetilde{r_{-(T+\epsilon)}} \cdot \widetilde{\phi} \right) \right| = m+1$$

and  $\mu(\widetilde{r_{-(T-\epsilon)}} \cdot \widetilde{\phi}) = -j + 1$ . It thus follows from Proposition 2.12(v) that if either k is even or j is odd or m > 1 then  $\Pi(\mathcal{L}(\widetilde{\phi}))$  has infinitely many translated points of translation T. If moreover k is prime then Proposition 2.12(v) also implies that the set of translated points of translation T of  $\Pi(\widetilde{\phi})$  has cohomological index greater or equal than m, and greater or equal than m + 1 if either k = 2 or j is odd.

#### Non-degeneracy

Assume that

$$c_{-2n+1}(\widetilde{\phi}) = c_0(\widetilde{\phi}) = 0.$$

Then for  $\epsilon > 0$  small enough we have

$$\left| \mu(\widetilde{r_{\epsilon}} \cdot \widetilde{\phi}) - \mu(\widetilde{r_{-\epsilon}} \cdot \widetilde{\phi}) \right| = 2m$$

and  $\mu(\tilde{r}_{\epsilon} \cdot \tilde{\phi}) = 2n$ . By Proposition 2.12 (v) we then conclude that  $\Pi(\tilde{\phi})$  is the identity. Since, by spectrality,  $0 \in \bar{\mathcal{A}}(\tilde{\phi})$ , we have in fact that  $\Pi(\mathcal{L}(\tilde{\phi}))$  is the identity.

#### Composition with the Reeb flow

By definition of the spectral selectors, for every  $\tilde{\phi}$  and every  $T \in \mathbb{R}$  we have

$$c_{j}(\widetilde{r_{T}} \cdot \widetilde{\phi}) = \inf \left\{ \underline{T} \in \mathbb{R} \mid \mu\left(\widetilde{r_{-\underline{T}}} \cdot \widetilde{r_{T}} \cdot \widetilde{\phi}\right) \leqslant -j \right\} \\= \inf \left\{ \underline{T} - T \in \mathbb{R} \mid \mu\left(\widetilde{r_{-\underline{T}+T}} \cdot \widetilde{\phi}\right) \leqslant -j \right\} + T \\= c_{j}(\widetilde{\phi}) + T \,.$$

#### Periodicity

Using the previous property and Proposition 2.12 (vii) we have

$$c_j(\widetilde{\phi}) + 2\pi = c_j\left(\widetilde{r_{2\pi}} \cdot \widetilde{\phi}\right) = c_{j+2n}(\widetilde{\phi}).$$

TOME 8 (2025)

#### Monotonicity

Suppose that  $\widetilde{\phi} \leq \widetilde{\psi}$ . Then  $\widetilde{r_{-T}} \cdot \widetilde{\phi} \leq \widetilde{r_{-T}} \cdot \widetilde{\psi}$  and so, by Proposition 2.12 (iv),  $\mu\left(\widetilde{r_{-T}} \cdot \widetilde{\phi}\right) \leq \mu\left(\widetilde{r_{-T}} \cdot \widetilde{\psi}\right)$ .

This implies that  $c_j(\tilde{\phi}) \leq c_j(\tilde{\psi})$ .

#### Continuity

Suppose that  $\tilde{\phi} \cdot \tilde{\psi}^{-1}$  is represented by a contact isotopy with Hamiltonian function  $H_t : L_k^{2n-1}(\underline{w}) \to \mathbb{R}$  with respect to  $\alpha_0$ . Let  $m(t) = \min H_t$  and  $M(t) = \max H_t$ . The flows of m and M are respectively  $\{r_{\int_0^t m}\}$  and  $\{r_{\int_0^t M}\}$ , thus

$$\widetilde{r_{\int_0^1 m}} \leqslant \widetilde{\phi} \cdot \widetilde{\psi}^{-1} \leqslant \widetilde{r_{\int_0^1 M}}$$

and so

$$\widetilde{r_{\int_0^1 m}} \cdot \widetilde{\psi} \leqslant \widetilde{\phi} \leqslant \widetilde{r_{\int_0^1 M}} \cdot \widetilde{\psi}.$$

By the composition with the Reeb flow property and monotonicity we thus have

(3.2) 
$$\int_0^1 \min H_t \, dt \leqslant c_j(\tilde{\phi}) - c_j(\tilde{\psi}) \leqslant \int_0^1 \max H_t \, dt \, .$$

We now show that (3.2) implies that each  $c_j$  is continuous with respect to the  $\mathcal{C}^1$ topology. Notice first that the Shelukhin–Hofer norm  $\nu_{\alpha} : \widetilde{\text{Cont}}_0(L_k^{2n-1}(\underline{w}), \xi_0) \to \mathbb{R}$ , which is defined by

$$\nu_{\alpha}(\widetilde{\phi}) = \inf \int_{0}^{1} \max |H_t| \, dt$$

with the infimum taken over all contact Hamiltonian functions  $H_t$  whose flow represents  $\tilde{\phi}$ , is continuous with respect to the  $\mathcal{C}^1$ -topology. This follows from the main result of [AA23], or can be seen directly as follows. Since  $\operatorname{Cont}_0(L_k^{2n-1}(\underline{w}), \xi_0)$  is a topological group, it is enough to show that  $\nu_{\alpha}$  is  $\mathcal{C}^1$ -continuous at the identity, i.e. that for every  $\epsilon > 0$  there is a  $\mathcal{C}^1$ -neighborhood  $\tilde{\mathcal{U}}$  of id in  $\operatorname{Cont}_0(L_k^{2n-1}(\underline{w}), \xi_0)$  such that for every  $\tilde{\phi} \in \tilde{\mathcal{U}}$  we have  $\nu_{\alpha}(\tilde{\phi}) < \epsilon$ . As in the proof of Lemma 3.1 we consider the product  $L_k^{2n-1}(\underline{w}) \times L_k^{2n-1}(\underline{w}) \times \mathbb{R}$  endowed with the contact structure given by the kernel of the contact form  $\pi_2^*\alpha_0 - e^{\theta}\pi_1^*\alpha_0$ . Applying the Weinstein theorem we can find a neighborhood  $\mathcal{U}(\operatorname{gr}_{\alpha_0}(\operatorname{id}))$  of  $\operatorname{gr}_{\alpha_0}(\operatorname{id})$ , a neighborhood  $\mathcal{U}(j^{10})$  of the zero section of  $J^1 L_k^{2n-1}(\underline{w})$  of the form  $\mathcal{U}(j^{10}) = \underline{\mathcal{U}}(j^{10}) \times (-\epsilon', \epsilon')$  with  $\epsilon' < \epsilon$  and a diffeomorphism  $\Psi$  from  $\mathcal{U}(\operatorname{gr}_{\alpha_0}(\operatorname{id}))$  to  $\mathcal{U}(j^{10})$  with  $\Psi(\operatorname{gr}_{\alpha_0}(\operatorname{id})) = j^{10}$  and  $\Psi^*(dz - \lambda_{\operatorname{can}}) = \pi_2^*\alpha_0 - e^{\theta}\pi_1^*\alpha_0$ . Since the map  $j^1: \mathcal{C}^{\infty}(L_k^{2n-1}(\underline{w})) \to \mathcal{L}eg(j^{10})$  that associates to a function its 1-jet is a local homeomorphism with respect to the  $\mathcal{C}^2$ -neighborhood  $\mathcal{U}(0)$  of the zero function in  $\mathcal{C}^{\infty}(L_k^{2n-1}(\underline{w}))$  such that  $j^1f \in \mathcal{U}(j^{10})$  for any  $f \in \mathcal{U}(0)$ , and the map  $\mathcal{U}(0) \mapsto \mathcal{L}eg(\operatorname{gr}_{\alpha_0}(\operatorname{id}))$  that sends f to  $\Psi^{-1}(j^1f)$  is a local homeomorphism. Since the map  $\operatorname{gr}_{\alpha_0}$ :  $\operatorname{Cont}_k(L_k^{2n-1}(\underline{w}), \xi_0) \to \mathcal{L}eg(\operatorname{gr}_{\alpha_0}(\operatorname{id}))$  that associates to a contactomorphism its graph is a local homeomorphism with

respect to the  $\mathcal{C}^1$ -topologies, we obtain a map  $\mathcal{U}(0) \to \operatorname{Cont}_0(L_k^{2n-1}(\underline{w}), \xi_0)$  that associates to a function f a contactomorphism  $\phi$  with  $j^1 f = \Psi(\operatorname{gr}_{\alpha_0}(\phi))$ , which is a homeomorphism on its image  $\mathcal{U}$ . Since  $\mathcal{U}(0)$  is convex,  $\mathcal{U}$  is simply connected; let thus  $\widetilde{\mathcal{U}}$  be the open neighborhood of id in  $\operatorname{Cont}_0(L_k^{2n-1}(\underline{w}), \xi_0)$  that projects homeomorphically to  $\mathcal{U}$ . Consider now any  $\widetilde{\phi}$  in  $\widetilde{\mathcal{U}}$ . Let  $f: L_k^{2n-1}(\underline{w}) \to (-\epsilon, \epsilon)$  be the function in  $\mathcal{U}(0)$  such that  $j^1 f = \Psi(\operatorname{gr}_{\alpha_0}(\Pi(\widetilde{\phi})))$ . Since  $\mathcal{U}(0)$  is convex,  $f_t := tf$  is in  $\mathcal{U}(0)$  for every  $t \in [0, 1]$ , and so for every  $t \in [0, 1]$  there is  $\phi_t \in \mathcal{U}$  with  $j^1 f_t = \Psi(\operatorname{gr}_{\alpha_0}(\phi_t))$ . Consider the two Legendrian isotopies  $j^1 f_t$  and  $\operatorname{gr}_{\alpha_0}(\phi_t)$ , with parametrisations given respectively by

$$i_1: [0,1] \times j^1 0 \to J^1 L_k^{2n-1}(\underline{w}), \ i_1(t,(x,0,0)) = j^1 f_t(x)$$

and

 $i_2: [0,1] \times \operatorname{gr}_{\alpha_0}(\operatorname{id}) \to L_k^{2n-1}(\underline{w}) \times L_k^{2n-1}(\underline{w}) \times \mathbb{R}, \ i_2(t,(x,x,0)) = (x,\phi_t(x),g_t(x)),$ where  $a_t$  is the conformal factor of  $\phi_t$ . Let  $H_t$  be the contact Hamiltonian function

where  $g_t$  is the conformal factor of  $\phi_t$ . Let  $H_t$  be the contact Hamiltonian function of the contact isotopy  $\{\phi_t\}$ . Then

$$H_t(\phi_t(x)) = \left(\pi_2^* \alpha_0 - e^{\theta} \pi_1^* \alpha_0\right) \left(\frac{d}{dt} i_2(t, (x, x, 0))\right)$$
$$= \left(dz - \lambda_{\text{can}}\right) \left(\frac{d}{dt} i_1(t, (x, 0, 0))\right) = f(x),$$

and so  $|H_t| < \epsilon$ . Moreover  $[\{\phi_t\}_{t \in [0,1]}] = \tilde{\phi}$ , because  $\tilde{\phi} \cdot [\{\phi_t\}_{t \in [0,1]}]^{-1}$  can be represented by a loop in  $\mathcal{U}$ , which is simply connected. We thus conclude that  $\nu_{\alpha}(\tilde{\phi}) < \epsilon$ , as we wanted.

Using (3.2) and the fact that  $\nu_{\alpha}$  is  $\mathcal{C}^1$ -continuous we now deduce that each  $c_j$  is  $\mathcal{C}^1$ -continuous. Let  $\tilde{\phi} \in \widetilde{\text{Cont}}_0(L_k^{2n-1}(\underline{w}), \xi_0)$ . By  $\mathcal{C}^1$ -continuity of  $\nu_{\alpha}$ , for any  $\epsilon > 0$  there is a  $\mathcal{C}^1$ -neighborhood  $\widetilde{\mathcal{U}}$  of  $\widetilde{\text{id}}$  in  $\widetilde{\text{Cont}}_0(L_k^{2n-1}(\underline{w}), \xi_0)$  such that  $\nu_{\alpha}|_{\widetilde{\mathcal{U}}} < \epsilon$ . Then  $\widetilde{\mathcal{V}} := \widetilde{\mathcal{U}} \cdot \widetilde{\phi}$  is a  $\mathcal{C}^1$ -neighborhood of  $\widetilde{\phi}$  such that for every  $\widetilde{\psi} \in \widetilde{\mathcal{V}}$  we have

$$\nu_{\alpha}(\widetilde{\phi}\cdot\widetilde{\psi}^{-1}) = \nu_{\alpha}(\widetilde{\psi}\cdot\widetilde{\phi}^{-1}) < \epsilon.$$

This implies that there is a contact Hamiltonian function  $H_t$  whose flow represents  $\tilde{\phi} \cdot \tilde{\psi}^{-1}$  and satisfies  $\int_0^1 \max |H_t| dt < \epsilon$ . Using (3.2) we thus conclude that for every  $\tilde{\psi} \in \tilde{\mathcal{V}}$  we have  $|c_j(\tilde{\phi}) - c_j(\tilde{\psi})| < \epsilon$ , and so that  $c_j$  is  $\mathcal{C}^1$ -continuous.

#### Triangle inequality

We have to prove that if either k is even or j is even then

$$c_{j+l}(\tilde{\phi} \cdot \tilde{\psi}) \leqslant c_j(\tilde{\phi}) + \left[c_l(\tilde{\psi})\right]_{T_{\underline{w}}}$$

Let  $T = c_j(\tilde{\phi})$  and  $N = \lceil c_l(\tilde{\psi}) \rceil_{T_{\underline{w}}}$ . Since

$$\Pi: \widetilde{\operatorname{Cont}}_0\left(L_k^{2n-1}(\underline{w}), \xi_0\right) \to \operatorname{Cont}_0\left(L_k^{2n-1}(\underline{w}), \xi_0\right)$$

is a local homeomorphism, by the continuity property of the spectral selectors and Lemma 3.1 we can assume that  $\Pi(\tilde{\phi})$  is non-degenerate with respect to  $\alpha_0$ . By the relation with translated points property, the spectral selectors of  $\tilde{\phi}$  are thus all distinct, and so  $\mu(\tilde{r}_{-T} \cdot \tilde{\phi}) = -j$ . Using the fact that  $\tilde{r}_{-N}$  commutes with  $\tilde{\phi}$  (since  $\{r_t\}$  has period  $T_w$ ) and the triangle inequality for the non-linear Maslov index (Proposition 2.12 (iii)) we thus have

$$\mu \left( \widetilde{r_{-(T+N)}} \cdot \widetilde{\phi} \cdot \widetilde{\psi} \right)$$
  
=  $\mu \left( \left( \widetilde{r_{-T}} \cdot \widetilde{\phi} \right) \cdot \left( \widetilde{r_{-N}} \cdot \widetilde{\psi} \right) \right) \leq \mu \left( \widetilde{r_{-T}} \cdot \widetilde{\phi} \right) + \mu \left( \widetilde{r_{-N}} \cdot \widetilde{\psi} \right) \leq -(j+l) .$ 

By definition of  $c_{i+l}$  we conclude that

$$c_{j+l}\left(\widetilde{\phi}\cdot\widetilde{\psi}\right) \leqslant T+N = c_j(\widetilde{\phi}) + \left[c_l(\widetilde{\psi})\right]_{T_{\underline{w}}}.$$

#### Conjugation invariance

We have to prove that

(3.3) 
$$\left[c_j\left(\tilde{\psi}\cdot\tilde{\phi}\cdot\tilde{\psi}^{-1}\right)\right]_{T_{\underline{w}}} = \left[c_j(\tilde{\phi})\right]_{T_{\underline{w}}}$$

Assume first that  $\Pi(\tilde{\phi})$  does not have discriminant points. Let  $\{\psi_t\}_{t \in [0,1]}$  be a contact isotopy representing  $\tilde{\psi}$ , and consider the homotopy  $\tilde{\psi}_s = [\{\psi_{st}\}_{t \in [0,1]}]$  from  $\tilde{\psi}_0 = id$ to  $\tilde{\psi}_1 = \tilde{\psi}$ . By the continuity property, the map

$$s \mapsto c_j \left( \widetilde{\psi}_s \cdot \widetilde{\phi} \cdot \widetilde{\psi}_s^{-1} \right) \in \overline{\mathcal{A}} \left( \widetilde{\psi}_s \cdot \widetilde{\phi} \cdot \widetilde{\psi}_s^{-1} \right)$$

is continuous. Moreover,  $c_j(\tilde{\psi}_s \cdot \tilde{\phi} \cdot \tilde{\psi}_s^{-1}) \in \mathbb{R} \smallsetminus T_{\underline{w}} \cdot \mathbb{Z}$  for all  $s \in [0, 1]$ . Indeed, if we had  $c_j(\tilde{\psi}_{\underline{s}} \cdot \tilde{\phi} \cdot \tilde{\psi}_{\underline{s}}^{-1}) \in T_{\underline{w}} \cdot \mathbb{Z}$  for some  $\underline{s}$  then, by the spectrality property,  $\Pi(\tilde{\psi}_{\underline{s}} \cdot \tilde{\phi} \cdot \tilde{\psi}_{\underline{s}}^{-1})$  would have discriminant points. But this is absurd, because the discriminant points of  $\Pi(\tilde{\psi}_{\underline{s}} \cdot \tilde{\phi} \cdot \tilde{\psi}_{\underline{s}}^{-1})$  are in bijection with the discriminant points of  $\Pi(\tilde{\phi})$ . We thus obtain (3.3) in this case.

The general case can be obtained as follows. Given any  $\tilde{\phi} \in Cont_0(L_k^{2n-1}(\underline{w}), \xi_0)$ , since (by Lemma 3.2)  $\mathcal{A}(\tilde{\phi})$  is nowhere dense, there is a sequence  $(\epsilon_l)$  of positive real numbers with  $\epsilon_l \to 0$  such that, for every l,  $\Pi(\widetilde{r_{-\epsilon_l}} \cdot \tilde{\phi})$  does not have discriminant points. Pose  $\widetilde{\chi_l} = \widetilde{r_{-\epsilon_l}} \cdot \tilde{\phi}$ . By the first part of the proof we have

(3.4) 
$$\left[c_{j}\left(\widetilde{\psi}\cdot\widetilde{\chi_{l}}\cdot\widetilde{\psi}^{-1}\right)\right]_{T_{\underline{w}}}=\left[c_{j}(\widetilde{\chi_{l}})\right]_{T_{\underline{w}}}$$

for all l. Since  $(\tilde{\chi}_l)$  converges to  $\tilde{\phi}$  in the  $\mathcal{C}^1$ -topology and  $\tilde{\chi}_l \leq \tilde{\phi}$  for all l, by the continuity and monotonicity properties of the spectral selectors for l big enough we have  $\lceil c_j(\tilde{\phi}) \rceil_{T_{\underline{w}}} = \lceil c_j(\tilde{\chi}_l) \rceil_{T_{\underline{w}}}$  and  $\lceil c_j(\tilde{\psi} \cdot \tilde{\phi} \cdot \tilde{\psi}^{-1}) \rceil_{T_{\underline{w}}} = \lceil c_j(\tilde{\psi} \cdot \tilde{\chi}_l \cdot \tilde{\psi}^{-1}) \rceil_{T_{\underline{w}}}$ . Equation (3.4) thus gives the desired result (3.3).

#### Poincaré duality

We first notice that if  $\Pi(\tilde{\phi})$  does not have discriminant points then

(3.5) 
$$\left| c_j(\widetilde{\phi}) \right|_{T_{\underline{w}}} = \max \left\{ N \in T_{\underline{w}} \cdot \mathbb{Z} \mid \mu(\widetilde{r_{-N}} \cdot \widetilde{\phi}) > -j \right\}.$$

Indeed, by spectrality we have  $c_j(\tilde{\phi}) \notin T_{\underline{w}} \cdot \mathbb{Z}$  and thus  $\underline{N} := \lfloor c_j(\tilde{\phi}) \rfloor_{T_{\underline{w}}} < c_j(\tilde{\phi})$ . This implies that  $\mu(\widetilde{r_{-\underline{N}}} \cdot \tilde{\phi}) > -j$ , and so the inequality  $\leq$  in (3.5). On the other hand, the opposite inequality follows (without any assumption on  $\tilde{\phi}$ ) from (3.1) and the fact that  $\lfloor c_j(\tilde{\phi}) \rfloor_{T_{\underline{w}}} + T_{\underline{w}} \geq \lceil c_j(\tilde{\phi}) \rceil_{T_{\underline{w}}}$ .

We now prove the Poincaré duality property, i.e. that

(3.6) 
$$\left[c_{j}(\tilde{\phi})\right]_{T_{\underline{w}}} = -\left\lfloor c_{-j-(2n-1)}(\tilde{\phi}^{-1})\right\rfloor_{T_{\underline{w}}}$$

for any  $\tilde{\phi} \in \widetilde{\text{Cont}}_0(L_k^{2n-1}(\underline{w}), \xi_0)$ . Assume first that  $\Pi(\tilde{\phi})$  does not have discriminant points. Then Proposition 2.12 (ix) implies that

(3.7) 
$$\mu(\widetilde{r_N} \cdot \widetilde{\phi}) + \mu\left(\widetilde{\phi}^{-1} \cdot \widetilde{r_{-N}}\right) = 2n$$

for every N that is a multiple of  $T_{\underline{w}}$ . The Poincaré duality (3.6) then follows from (3.1), (3.5), (3.7) and the fact that  $\tilde{\phi}^{-1} \cdot \widetilde{r_{-N}} = \widetilde{r_{-N}} \cdot \tilde{\phi}^{-1}$  for every N that is a multiple of  $T_{\underline{w}}$ .

For a general  $\tilde{\phi}$ , as in the proof of conjugation invariance we can find a sequence  $(\tilde{\chi}_l)$  that converges to  $\tilde{\phi}$  in the  $C^1$ -topology and such that, for all  $l, \tilde{\chi}_l \leq \tilde{\phi}$  and  $\Pi(\tilde{\chi}_l)$  does not have discriminant points. By the first part of the proof we have  $\lceil c_j(\tilde{\chi}_l) \rceil_{T_{\underline{w}}} = -\lfloor c_{-j-(2n-1)}(\tilde{\chi}_l^{-1}) \rfloor_{T_{\underline{w}}}$ . By monotonicity and continuity of the spectral selectors we thus obtain (3.6) also in this case.

### 4. Non-shortening of the standard Reeb flow with respect to the discriminant and oscillation norms

Recall from [CS15] that, for any closed contact manifold  $(M, \xi)$ , the discriminant norm  $\nu_{\text{dis}}$  on the universal cover  $\widetilde{\text{Cont}}_0(M, \xi)$  of the identity component of the contactomorphism group is the word norm associated to the generating set  $\mathcal{D}$  formed by elements  $\phi$  that can be represented by an embedded contact isotopy, i.e. a contact isotopy  $\{\phi_t\}_{t \in [0,1]}$  such that  $\phi_t \circ \phi_s^{-1}$  has no discriminant points for all  $s \neq t \in [0,1]$ . Recall also from [Arl23] that the discriminant length of a contact isotopy  $\{\phi_t\}_{t \in [0,1]}$ is the minimal N such that there is a decomposition  $0 = t_0 < \cdots < t_N = 1$  of the time interval [0,1] with  $\{\phi_t\}_{t \in [t_j, t_{j+1}]}$  embedded for all  $j = 0, \cdots, N-1$ .

Consider the discriminant norm on  $\widetilde{\text{Cont}}_0(L_k^{2n-1}(\underline{w}),\xi_0)$ . For every positive real number T we have

(4.1) 
$$\nu_{\rm dis}(\widetilde{r_T}) \geqslant \frac{2n\lceil \frac{T}{2\pi}\rceil + 1}{2n+1}.$$

TOME 8 (2025)

Indeed, let  $N = \nu_{\text{dis}}(\widetilde{r_T})$  and write  $\widetilde{r_T} = \prod_{j=1}^N \widetilde{\phi}_j$  with  $\widetilde{\phi}_j \in \mathcal{D}$ . Then, by Proposition 2.12(i), (iii), (vi) and the first statement of (v) we have

$$2n\left\lceil \frac{T}{2\pi}\right\rceil = \mu(\widetilde{r_T}) \leqslant \sum_{j=1}^N \mu(\widetilde{\phi_j}) + N - 1 \leqslant 2nN + N - 1$$

Similarly, in the case of projective space we have

(4.2) 
$$\nu_{\rm dis}(\widetilde{r_T}) \geqslant \left\lceil \frac{T}{2\pi} \right\rceil$$

The estimates (4.1) and (4.2) are better than those obtained in [CS15, GKPS21], since in those references just the quasimorphism property of the non-liner Maslov index (Proposition 2.12(ii)) is used and not the triangle inequality (Proposition 2.12(iii)). However, they are still not optimal. Indeed, writing

$$0 < T_0 := \frac{T}{\left\lfloor \frac{k}{2\pi} T \right\rfloor + 1} < \frac{2\pi}{k}$$

we have

(4.3) 
$$\{r_{Tt}\}_{t \in [0,1]} = \{\underbrace{r_{T_0t} \circ \cdots \circ r_{T_0t}}_{\left\lfloor \frac{k}{2\pi}T \right\rfloor + 1} \}_{t \in [0,1]} .$$

Since the minimal period of a closed Reeb orbit of  $\alpha_0$  on  $L_k^{2n-1}(\underline{w})$  is  $\frac{2\pi}{k}$  we have that  $\{r_{T_0t}\}_{t \in [0,1]} \in \mathcal{D}$ , and so  $\{r_{Tt}\}_{t \in [0,1]}$  has discriminant length smaller or equal than  $\lfloor \frac{k}{2\pi}T \rfloor + 1$ ; this length is actually equal to  $\lfloor \frac{k}{2\pi}T \rfloor + 1$ , because for any interval  $[t_0, t_1]$  of length  $t_1 - t_0 \ge \frac{2\pi}{k}$  the contact isotopy  $\{r_t\}_{t \in [t_0, t_1]}$  is not embedded. For instance, in the case of projective space the discriminant length of  $\{r_{2\pi m t}\}_{t \in [0,1]}$  is thus 2m+1, while (4.2) only gives  $\nu_{\text{dis}}(\widetilde{r_{2\pi m}}) \ge m$ . In this section we prove the optimal estimates for the discriminant and oscillation lengths of the standard Reeb flow of lens spaces with equal weights using the spectral selectors defined in Section 3. The main advantage of using the spectral selectors is that while (by Proposition 2.12 (v)) the non-linear Maslov index only jumps in the presence of discriminant points of the lift of a contact isotopy of a lens space to the sphere, so that in particular for instance  $\mu(\widetilde{r_T}) = 2n \left\lceil \frac{T}{2\pi} \right\rceil$ , the spectral selectors allow to distinguish  $\widetilde{r_T}$  also for different values of T in  $[0, 2\pi]$ , indeed by Theorem 1.1 (v) we have for instance  $c_0(\widetilde{r_T}) = T$ .

We start with the following lemma.

LEMMA 4.1. — For any element 
$$\tilde{\phi}$$
 of  $\operatorname{Cont}_0(L_k^{2n-1}(\underline{w}), \xi_0)$ , if  $\tilde{\phi} \in \mathcal{D}$  then  
 $c_0(\tilde{\phi}) < T_w$ .

Proof. — If  $\tilde{\phi} \in \mathcal{D}$  then  $\tilde{\phi}$  can be represented by a contact isotopy  $\{\phi_t\}_{t \in [0,1]}$  such that  $\phi_t$  does not have discriminant points for all  $t \in (0,1]$ . Suppose by contradiction that  $c_0(\tilde{\phi}) \geq T_{\underline{w}}$ , and for  $s \in [0,1]$  let  $\tilde{\phi}_s = [\{\phi_{st}\}_{t \in [0,1]}]$ . Since  $c_0(\tilde{\phi}_1) = c_0(\tilde{\phi}) \geq T_{\underline{w}}$  and, by Theorem 1.1 (ii),  $c_0(\tilde{\phi}_0) = c_0(\tilde{id}) = 0$ , by continuity of  $c_0$  (Theorem 1.1 (viii)) there is a value of s in (0,1] such that  $c_0(\tilde{\phi}_s) = T_{\underline{w}}$ . But then, by spectrality (Theorem 1.1 (i)),  $T_{\underline{w}}$  belongs to  $\bar{\mathcal{A}}(\tilde{\phi}_s)$ . This means that  $\phi_s$  has discriminant points, which is a contradiction.

We can now prove that for every real number T the Reeb flow  $\{r_{Tt}\}_{t \in [0,1]}$  of the standard contact form  $\alpha_0$  on a lens space of the form  $L_k^{2n-1}(w, \cdots, w)$  is a geodesic for the discriminant norm. We have seen above that  $\{r_{Tt}\}_{t \in [0,1]}$  has discriminant length  $\lfloor \frac{k}{2\pi}T \rfloor + 1$ . In order to prove that it is a geodesic we thus have to show that

(4.4) 
$$\nu_{\rm dis}(\widetilde{r_T}) \ge \left\lfloor \frac{k}{2\pi} T \right\rfloor + 1$$

Let  $\nu_{\text{dis}}(\widetilde{r_T}) = N$ , and write  $\widetilde{r_T} = \prod_{j=1}^N \widetilde{\phi_j}$  with  $\widetilde{\phi_j} \in \mathcal{D}$  for all j. By Theorem 1.1 (v), (ix) and Lemma 4.1, and since  $T_{\underline{w}} = \frac{2\pi}{k}$  for  $\underline{w} = (w, \dots, w)$ , we then have

$$T = c_0(\widetilde{r_T}) \leqslant c_0(\widetilde{\phi_1}) + \sum_{j=2}^N \left[ c_0(\widetilde{\phi_j}) \right]_{\frac{2\pi}{k}} < N \, \frac{2\pi}{k} \, .$$

This implies that  $\nu_{\text{dis}}(\widetilde{r_T}) \ge \left\lfloor \frac{k}{2\pi}T \right\rfloor + 1$ , as we wanted. We now show that the standard Reeb flow  $\{r_{Tt}\}_{t \in [0,1]}$  on a lens space of the form

We now show that the standard Reeb flow  $\{r_{Tt}\}_{t\in[0,1]}$  on a lens space of the form  $L_k^{2n-1}(w, \dots, w)$  is a geodesic with respect to the oscillation norm. Recall from [CS15] that the oscillation pseudonorm  $\nu_{osc}$  on the universal cover  $Cont_0(M, \xi)$  of the identity component of the contactomorphism group of a closed contact manifold  $(M, \xi)$  is defined as follows. Let  $\mathcal{D}_+$  and  $\mathcal{D}_-$  be the sets of elements of  $Cont_0(M, \xi)$  that can be represented respectively by an embedded non-negative or non-positive contact isotopy. It is proved in [CS15] that every element  $\tilde{\phi}$  of  $Cont_0(M, \xi)$  can be written as  $\tilde{\phi} = \prod_{j=1}^N \tilde{\phi_j}$  with  $\tilde{\phi_j} \in \mathcal{D}_+$  or  $\tilde{\phi_j} \in \mathcal{D}_-$  for every j. We denote by  $\nu_+(\tilde{\phi})$  and  $\nu_-(\tilde{\phi})$  respectively the minimal number of elements of  $\mathcal{D}_+$  and minus the minimal number of elements of  $\mathcal{D}_+$  in such a decomposition. The oscillation pseudonorm is then defined by

$$\nu_{\rm osc}(\widetilde{\phi}) = \nu_+(\widetilde{\phi}) - \nu_-(\widetilde{\phi})$$

for  $\tilde{\phi} \neq i\tilde{d}$ , and  $\nu_{osc}(i\tilde{d}) = 0$ . By [CS15, Proposition 3.2], the oscillation pseudonorm on  $Cont_0(M, \xi)$  is non-degenerate if and only if  $(M, \xi)$  is orderable; it is thus a norm for lens spaces. Recall also from [Arl23] that the oscillation length of a contact isotopy  $\{\phi_t\}_{t \in [0,1]}$  is the sum of  $\mathcal{L}_+(\{\phi_t\}_{t \in [0,1]})$  and  $\mathcal{L}_-(\{\phi_t\}_{t \in [0,1]})$ , where  $\mathcal{L}_+(\{\phi_t\}_{t \in [0,1]})$  is the minimal  $N_+$  for which there is  $N \ge N_+$  and a decomposition  $0 = t_0 < \cdots < t_N =$ 1 with each  $\{\phi_t\}_{t \in [t_j, t_{j+1}]}$  embedded and non-negative or non-positive and exactly  $N_+$  of them non-negative, and  $\mathcal{L}_-(\{\phi_t\}_{t \in [0,1]})$  is the minimal  $N_-$  for which there is  $N \ge N_-$  and a decomposition  $0 = t_0 < \cdots < t_N = 1$  with each  $\{\phi_t\}_{t \in [t_j, t_{j+1}]}$ embedded and non-negative or non-positive and exactly  $N_-$  of them non-positive.

Consider now as above the class  $\widetilde{r_T} = [\{r_{Tt}\}_{t \in [0,1]}]$  of the Reeb flow of  $\alpha_0$  on a lens space  $(L_k^{2n-1}(\underline{w}), \xi_0)$ . Similarly as before, the oscillation length of  $\{r_{Tt}\}_{t \in [0,1]}$ is  $\lfloor \frac{k}{2\pi}T \rfloor + 1$ . The decomposition (4.3) shows moreover that  $\nu_{-}(\widetilde{r_T}) = 0$ , and thus  $\nu_{\text{osc}}(\widetilde{r_T}) = \nu_{+}(\widetilde{r_T})$ . In order to show that the Reeb flow  $\{r_{Tt}\}_{t \in [0,1]}$  of  $\alpha_0$  on a lens space of the form  $L_k^{2n-1}(w, \dots, w)$  is a geodesic with respect to the oscillation norm we thus have to show that

$$\nu_+(\widetilde{r_T}) \geqslant \left\lfloor \frac{k}{2\pi} T \right\rfloor + 1.$$

Let  $\nu_{+}(\widetilde{r_{T}}) = N_{+}$ , and write  $\widetilde{r_{T}} = \prod_{j=1}^{N} \widetilde{\phi_{j}}$  with  $\widetilde{\phi_{j}} \in \mathcal{D}_{\pm}$  for all j and with exactly  $N_{+}$  of the  $\widetilde{\phi_{j}}$  in  $\mathcal{D}_{+}$ . Denote such elements by  $\widetilde{\phi_{\sigma(1)}}, \cdots, \widetilde{\phi_{\sigma(N_{+})}}$ . Then  $\widetilde{r_{T}} \leq \prod_{j=1}^{N_{+}} \widetilde{\phi_{\sigma(j)}}$ , and so by Theorem 1.1(v), (vii), (ix) and Lemma 4.1, and since  $T_{\underline{w}} = \frac{2\pi}{k}$  for  $\underline{w} = (w, \cdots, w)$ , we have

$$T = c_0(\widetilde{r_T}) \leqslant c_0 \left(\prod_{j=1}^{N_+} \widetilde{\phi_{\sigma(j)}}\right) \leqslant c_0(\widetilde{\phi_{\sigma(1)}}) + \sum_{j=2}^{N_+} \left\lceil c_0\left(\widetilde{\phi_{\sigma(j)}}\right) \right\rceil_{\frac{2\pi}{k}} < \frac{2\pi}{k} \cdot N_+ \,.$$

This implies that  $\nu_+(\widetilde{r_T}) \ge \left\lfloor \frac{k}{2\pi} T \right\rfloor + 1$ , as we wanted.

Remark 4.2. — For a general lens space  $(L_k^{2n-1}(\underline{w}), \xi_0)$  the above discussion implies that the discriminant and oscillation norms of  $\widetilde{r_T} = [\{r_{Tt}\}_{t \in [0,1]}]$  are greater or equal than  $\lfloor \frac{T}{T_{\underline{w}}} \rfloor + 1$ , while the discriminant and oscillation lengths of  $\{r_{Tt}\}_{t \in [0,1]}$  are equal to  $\lfloor \frac{k}{2\pi}T \rfloor + 1$ . We do not know thus if the standard Reeb flow on general lens spaces is a geodesic for the discriminant and oscillation norms. The gap for general weights between the minimal period  $\frac{2\pi}{k}$  of a closed Reeb orbit and the period  $T_{\underline{w}}$  of the Reeb flow seems to suggest that it might be possible to shorten the Reeb flow. It would be interesting to investigate if this is indeed the case, or if other methods could be used to prove that the Reeb flow is still a geodesic.

#### 5. A spectral pseudonorm

Let 
$$c_{-} = c_{-2n+1}$$
 and  $c_{+} = c_{0}$ , and define  $\nu : \widetilde{\operatorname{Cont}}_{0}(L_{k}^{2n-1}(\underline{w}), \xi_{0}) \to T_{\underline{w}} \cdot \mathbb{Z}$  by  
 $\nu(\widetilde{\phi}) = \max\left\{\left[c_{+}(\widetilde{\phi})\right]_{T_{\underline{w}}}, -\left\lfloor c_{-}(\widetilde{\phi})\right\rfloor_{T_{\underline{w}}}\right\}.$ 

In this section we prove that  $\nu$  is a pseudonorm satisfying the properties stated in Corollary 1.6.

Recall that a pseudonorm  $\nu$  on a group G is said to be stably unbounded if there is an element  $\sigma$  of G such that  $\lim_{m\to\infty} \frac{\nu(\sigma^m)}{m} \neq 0$ , and is said to be compatible with a bi-invariant partial order  $\leq$  if id  $\leq \sigma_1 \leq \sigma_2$  implies  $\nu(\sigma_1) \leq \nu(\sigma_2)$ .

PROPOSITION 5.1. — The map  $\nu : \widetilde{\text{Cont}}_0(L_k^{2n-1}(\underline{w}), \xi_0) \to T_{\underline{w}} \cdot \mathbb{Z}$  is a stably unbounded conjugation invariant pseudonorm compatible with the partial order  $\leq$ .

Proof. — We first show that for every  $\tilde{\phi}$  we have  $\nu(\tilde{\phi}) \ge 0$ . Suppose by contradiction that  $\nu(\tilde{\phi}) < 0$ . Then  $\lceil c_+(\tilde{\phi}) \rceil_{\underline{T}_{\underline{w}}} < 0$ , thus  $c_+(\tilde{\phi}) < 0$ , and  $-\lfloor c_-(\tilde{\phi}) \rfloor_{\underline{T}_{\underline{w}}} < 0$ , thus  $c_-(\tilde{\phi}) > 0$ . But this contradicts the fact that, since the sequence  $c_j$  is non-decreasing,  $c_-(\tilde{\phi}) \le c_+(\tilde{\phi})$ . The triangle inequality  $\nu(\tilde{\phi} \cdot \tilde{\psi}) \le \nu(\tilde{\phi}) + \nu(\tilde{\psi})$  follows from Theorem 1.1 (ix) and (xi), and symmetry  $\nu(\tilde{\phi}) = \nu(\tilde{\phi}^{-1})$  from Theorem 1.1 (xi). This shows that  $\nu$  is a pseudonorm. Invariance by conjugation follows from Theorem 1.1 (x) and (xi). The pseudonorm  $\nu$  is stably unbounded, indeed Theorem 1.1 (v) implies that

$$\nu\left(\widetilde{r_{T_{\underline{w}}}}^{m}\right) = \nu\left(\widetilde{r_{mT_{\underline{w}}}}\right) = m T_{\underline{w}}$$

for every positive integer m, thus posing  $\sigma = \widetilde{r_{T_{\underline{w}}}}$  we have  $\lim_{m\to\infty} \frac{\nu(\sigma^m)}{m} = T_{\underline{w}} \neq 0$ . Finally, the fact that  $\nu$  is compatible with the partial order  $\leq$  follows from Theorem 1.1 (vii) and (xi).

It would be interesting to know if  $\nu$  is equivalent to the oscillation norm  $\nu_{osc}$ . In this direction, we prove the following inequality.

PROPOSITION 5.2. — For every element 
$$\tilde{\phi}$$
 of  $\widetilde{\text{Cont}}_0(L_k^{2n-1}(\underline{w}), \xi_0)$  we have  
 $\nu(\tilde{\phi}) \leq T_w \cdot \nu_{\text{osc}}(\tilde{\phi})$ .

Proof. — Let  $\nu_{+}(\tilde{\phi}) = N_{+}$ , and write  $\tilde{\phi} = \prod_{j=1}^{N} \widetilde{\phi_{j}}$  with all the  $\widetilde{\phi_{j}}$  in  $\mathcal{D}_{+}$  or  $\mathcal{D}_{-}$ and exactly  $N_{+}$  of them in  $\mathcal{D}_{+}$ . Denote such elements by  $\widetilde{\phi_{\sigma(1)}}, \cdots, \widetilde{\phi_{\sigma(N_{+})}}$ . Then  $\widetilde{\phi} \leq \prod_{j=1}^{N_{+}} \widetilde{\phi_{\sigma(j)}}$ , and thus by Theorem 1.1 (vii), (ix) and Lemma 4.1 we have

$$c_{+}(\widetilde{\phi}) \leqslant c_{+} \left(\prod_{j=1}^{N_{+}} \widetilde{\phi_{\sigma(j)}}\right) \leqslant \sum_{j=1}^{N_{+}} \left\lceil c_{+} \left(\widetilde{\phi_{\sigma(j)}}\right) \right\rceil_{T_{\underline{w}}} \leqslant T_{\underline{w}} \cdot N_{+}$$

Similarly, setting  $\nu_{-}(\tilde{\phi}) = -N_{-}$  we have  $c_{+}(\tilde{\phi}^{-1}) \leq T_{\underline{w}} \cdot N_{-}$  and so, by Theorem 1.1 (xi),

$$-\left\lfloor c_{-}(\widetilde{\phi})\right\rfloor_{T_{\underline{w}}} = \left\lceil c_{+}\left(\widetilde{\phi}^{-1}\right)\right\rceil_{T_{\underline{w}}} \leqslant T_{\underline{w}} \cdot N_{-} \,.$$

We deduce that

$$\nu(\widetilde{\phi}) \leqslant T_{\underline{w}} \max\left\{\nu_{+}(\widetilde{\phi}), -\nu_{-}(\widetilde{\phi})\right\} \leqslant T_{\underline{w}} \left(\nu_{+}(\widetilde{\phi}) - \nu_{-}(\widetilde{\phi})\right) = T_{\underline{w}} \cdot \nu_{\mathrm{osc}}(\widetilde{\phi}) \,. \qquad \Box$$

We do not know whether the pseudonorm  $\nu$  is non-degenerate, i.e. whether  $\nu(\tilde{\phi}) = 0$ if and only if  $\tilde{\phi} = i\tilde{d}$ . Indeed, by the definition of  $\nu$  we have that  $\nu(\tilde{\phi}) = 0$  if and only if  $c_+(\tilde{\phi}) = c_-(\tilde{\phi}) = 0$ , which by Theorem 1.1(iv) only implies that  $\Pi(\tilde{\phi})$  is the identity. On the other hand, the induced conjugation invariant pseudonorm on  $\operatorname{Cont}_0(L_k^{2n-1}(\underline{w}), \xi_0)$ , i.e. the pseudonorm  $\nu_*$  defined by

$$\nu_*(\phi) = \inf \left\{ \nu(\widetilde{\phi}) \mid \Pi(\widetilde{\phi}) = \phi \right\},$$

is non-degenerate, hence a norm. However, this norm is bounded (hence equivalent to the trivial norm, since it is discrete), as shown in the following proposition.

PROPOSITION 5.3. — For every 
$$\phi \in \text{Cont}_0(L_k^{2n-1}(\underline{w}), \xi_0)$$
 we have  
 $\nu_*(\phi) \leq 2\pi + T_{\underline{w}}.$ 

Proof. — We show that

(5.1) 
$$\nu_* \left( \Pi(\tilde{\phi}) \right) \leqslant \left[ c_+(\tilde{\phi}) \right]_{T_{\underline{w}}} - \left\lfloor c_-(\tilde{\phi}) \right\rfloor_{T_{\underline{w}}} \leqslant 2\pi + T_{\underline{w}}$$

for every  $\tilde{\phi} \in \widetilde{\text{Cont}_0}(L_k^{2n-1}(\underline{w}), \xi_0)$ . Using periodicity of the spectral selectors (Theorem 1.1 (vi)) and the fact that the sequence of spectral selectors  $c_j$  is non-decreasing we have  $c_+(\tilde{\phi}) \leq c_-(\tilde{\phi}) + 2\pi$ , which implies the second inequality in (5.1). For the first inequality, it is enough to find  $N \in T_{\underline{w}} \cdot \mathbb{Z}$  such that  $\nu(\widetilde{r_{-N}} \cdot \tilde{\phi}) = \lceil c_+(\tilde{\phi}) \rceil_{T_{\underline{w}}} - \lfloor c_-(\tilde{\phi}) \rfloor_{T_{\underline{w}}}$ .

Suppose first that  $\nu(\tilde{\phi}) = \lceil c_+(\tilde{\phi}) \rceil_{T_{\underline{w}}}$ , and pose  $N = \lceil c_+(\tilde{\phi}) \rceil_{T_{\underline{w}}}$ . By Theorem 1.1 (v) we then have

$$\nu(\widetilde{r_{-N}} \cdot \widetilde{\phi}) = \max\left\{ \left[ c_{+}(\widetilde{r_{-N}} \cdot \widetilde{\phi}) \right]_{T_{\underline{w}}}, - \left[ c_{-}(\widetilde{r_{-N}} \cdot \widetilde{\phi}) \right]_{T_{\underline{w}}} \right\}$$
$$= \max\left\{ 0, \left[ c_{+}(\widetilde{\phi}) \right]_{T_{\underline{w}}} - \left[ c_{-}(\widetilde{\phi}) \right]_{T_{\underline{w}}} \right\} = \left[ c_{+}(\widetilde{\phi}) \right]_{T_{\underline{w}}} - \left[ c_{-}(\widetilde{\phi}) \right]_{T_{\underline{w}}}.$$

Similarly, if  $\nu(\tilde{\phi}) = -\lfloor c_{-}(\tilde{\phi}) \rfloor_{T_{\underline{w}}}$  then, posing  $N = \lfloor c_{-}(\tilde{\phi}) \rfloor_{T_{\underline{w}}}$  we have

$$\nu(\widetilde{r_{-N}} \cdot \widetilde{\phi}) = \left[ c_+(\widetilde{\phi}) \right]_{T_{\underline{w}}} - \left\lfloor c_-(\widetilde{\phi}) \right\rfloor_{T_{\underline{w}}}.$$

Remark 5.4. — It follows from [AA23, Corollary 4.12] that on the universal cover of the identity component of the contactomorphism group of the unit cotangent bundle of the torus  $\mathbb{T}^n$  for  $n \ge 2$  the difference of the invariants  $c_+$  and  $c_-$  defined in [AA23] is unbounded. This difference with respect to (5.1) might be related to the fact that the identity component of the contactomorphism group of the unit cotangent bundle of the torus does not contain positive loops. It would be interesting to investigate if on the other hand the difference of the invariants  $c_+$  and  $c_-$  of [AA23] is bounded on  $\widetilde{\text{Cont}}_0(L_k^{2n-1}(\underline{w}), \xi_0)$ . This would then imply as in Proposition 5.3 that the induced norm on  $\operatorname{Cont}_0(L_k^{2n-1}(\underline{w}), \xi_0)$  is bounded, and therefore answer partially a question in [FPR18, Example 2.21].

Remark 5.5. — If  $\nu: G \to \mathbb{R}_{\geq 0}$  is a pseudonorm on a group G then, for any c > 0, the map  $\nu': G \to \mathbb{R}_{\geq 0}$  defined by

$$\nu'(g) := \begin{cases} \max\{\nu(g), c\} & \text{if } g \neq \text{id} \\ 0 & \text{if } g = \text{id} \end{cases}$$

is a norm. Moreover,  $\nu'$  is invariant by conjugation if and only if so is  $\nu$ . This trick (which is similar to one used in [BIP08]) can be applied to our pseudonorm  $\nu$ , with  $c = T_{\underline{w}}$ , to obtain a stably unbounded conjugation invariant norm  $\nu'$  on  $\widetilde{\text{Cont}}_0(L_k^{2n-1}(\underline{w}), \xi_0)$ . Since  $\nu$  takes values in  $T_{\underline{w}} \cdot \mathbb{Z}$ , if  $\nu$  is already a norm then  $\nu' \equiv \nu$ . Proposition 5.2 holds also for  $\nu'$ , indeed for any element  $\tilde{\phi} \neq id$  we have

$$\nu'(\widetilde{\phi}) = \max\left\{ \left[ c_{+}(\widetilde{\phi}) \right]_{T_{\underline{w}}}, -\left\lfloor c_{-}(\widetilde{\phi}) \right\rfloor_{T_{\underline{w}}}, T_{\underline{w}} \right\}$$
$$\leqslant T_{\underline{w}} \cdot \max\left\{ \nu_{\text{osc}}(\widetilde{\phi}), 1 \right\}$$
$$= T_{w} \cdot \nu_{\text{osc}}(\widetilde{\phi}).$$

#### BIBLIOGRAPHY

- [AA23] Simon Allais and Pierre-Alexandre Arlove, Spectral selectors and contact orderability, 2023, https://arxiv.org/abs/2309.10578. ↑72, 74, 90, 98
- [All22a] Simon Allais, On the Hofer-Zehnder conjecture on CP<sup>d</sup> via generating functions, Internat. J. Math. 33 (2022), no. 10-11, article no. 2250072. ↑71
- [All22b] \_\_\_\_\_, On the minimal number of translated points in contact lens spaces, Proc. Amer. Math. Soc. 150 (2022), 2685–2693. ↑72

- [AM18] Peter Albers and Will J. Merry, Orderability, contact non-squeezing, and Rabinowitz Floer homology, J. Symplectic Geom. 16 (2018), no. 6, 1481–1547. ↑68
- [Arl23] Pierre-Alexandre Arlove, Geodesics of norms on the contactomorphism group of  $\mathbb{R}^{2n} \times S^1$ , J. Fixed Point Theory Appl. **25** (2023), no. 4, article no. 80.  $\uparrow$ 68, 73, 93, 95
- [Bhu01] Mohan Bhupal, A partial order on the group of contactomorphisms of  $\mathbb{R}^{2n+1}$  via generating functions, Turkish J. Math. **25** (2001), 125–135.  $\uparrow 68$
- [BIP08] Dmitri Burago, Sergei Ivanov, and Leonid Polterovich, Conjugation-invariant norms on groups of geometric origin, Groups of diffeomorphisms in honor of Shigeyuki Morita on the occasion of his 60th birthday, Advanced Studies in Pure Mathematics, vol. 52, Mathematical Society of Japan, 2008, pp. 221–250. ↑98
- [Car13] André Carneiro, A geometric construction of a Calabi quasimorphism on projective space, Ph.D. thesis, Columbia University, New York, USA, 2013. ↑71, 82
- [CS15] Vincent Colin and Sheila Sandon, The discriminant and oscillation lengths for contact and Legendrian isotopies, J. Eur. Math. Soc. 17 (2015), 1657–1685. ↑68, 73, 74, 93, 94, 95
- [EP00] Yakov Eliashberg and Leonid Polterovich, Partially ordered groups and geometry of contact transformations, Geom. Funct. Anal. 10 (2000), 1448–1476. ↑68
- [EP03] Michael Entov and Leonid Polterovich, Calabi quasimorphism and quantum homology, Int. Math. Res. Not. 2003 (2003), no. 30, 1635–1676. ↑71
- [FPR18] Maia Fraser, Leonid Polterovich, and Daniel Rosen, On Sandon-type metrics for contactomorphism groups, Ann. Math. Qué. 42 (2018), 191–214. ↑98
- [FSZ23] Maia Fraser, Sheila Sandon, and Bingyu Zhang, Contact non-squeezing at large scale via generating functions, to appear in the Journal of Fixed Point Theory and Applications, 2023, https://arxiv.org/abs/2310.11993. ↑68
- [GG73] Martin Golubitsky and Victor Guillemin, Stable mappings and their singularities, Graduate Texts in Mathematics, vol. 14, Springer, 1973. ↑87
- [Giv90] Alexander B. Givental, Nonlinear generalization of the Maslov index, Theory of singularities and its applications, Advances in Soviet Mathematics, vol. 1, American Mathematical Society, 1990, pp. 71–103. ↑68, 69, 71, 72, 79
- [GKPS21] Gustavo Granja, Yael Karshon, Milena Pabiniak, and Sheila Sandon, Givental's nonlinear Maslov index on lens spaces, Int. Math. Res. Not. 2021 (2021), 18225–18299. ↑68, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 94
- [Mil08] Isidora Milin, Orderability of contactomorphism groups of lens spaces, Ph.D. thesis, Stanford University, Stanford, USA, 2008. ↑68
- [San10] Sheila Sandon, An integer-valued bi-invariant metric on the group of contactomorphisms of  $\mathbb{R}^{2n} \times S^1$ , J. Topol. Anal. **2** (2010), 327–339.  $\uparrow 68$
- [San11a] \_\_\_\_\_, Contact homology, capacity and non-squeezing in  $\mathbb{R}^{2n} \times S^1$  via generating functions, Ann. Inst. Fourier **61** (2011), 145–185.  $\uparrow$ 68, 71, 72
- [San11b] \_\_\_\_\_, Equivariant homology for generating functions and orderability of lens spaces, J. Symplectic Geom. 9 (2011), 123–146. ↑68
- [San13] \_\_\_\_\_, A Morse estimate for translated points of contactomorphisms of spheres and projective spaces, Geom. Dedicata 165 (2013), 95–110. ↑68
- [Thé98] David Théret, Rotation numbers of Hamiltonian isotopies in complex projective spaces, Duke Math. J. 94 (1998), 13–27. ↑69, 71, 88
- [Tsu08] Takashi Tsuboi, On the simplicity of the group of contactomorphisms, Groups of diffeomorphisms, Advanced Studies in Pure Mathematics, vol. 52, Mathematical Society of Japan, 2008, pp. 491–504. ↑86
- [Vit92] Claude Viterbo, Symplectic topology as the geometry of generating functions, Math. Ann.
   292 (1992), 685–710. ↑71

Manuscript received on 4th April 2024, accepted on 30th September 2024.

Recommended by Editors C. Sorger and V. Colin. Published under license CC BY 4.0.



eISSN: 2644-9463 This journal is a member of Centre Mersenne.



Simon ALLAIS Université de Strasbourg, IRMA UMR 7501, F-67000 Strasbourg (France) simon.allais@math.unistra.fr

Pierre-Alexandre ARLOVE Ruhr-Universität Bochum, Fakultät für Mathematik, 44780 Bochum (Germany) pierre-alexandre.arlove@ruhr-uni-bochum.de

Sheila SANDON Université de Strasbourg, CNRS, IRMA UMR 7501, F-67000 Strasbourg (France) sandon@math.unistra.fr