



ANNALES
HENRI LEBESGUE

LOUIS IOOS

VICTORIA KAMINKER

LEONID POLTEROVICH

DOR SHMOISH

SPECTRAL ASPECTS OF THE BEREZIN TRANSFORM

ASPECTS SPECTRAUX DE LA
TRANSFORMÉE DE BEREZIN

ABSTRACT. — We discuss the Berezin transform, a Markov operator associated to positive operator valued measures (POVMs), in a number of contexts including the Berezin–Toeplitz quantization, Donaldson’s dynamical system on the space of Hermitian inner products on a complex vector space, representations of finite groups, and quantum noise. In particular, we calculate the spectral gap for quantization in terms of the fundamental tone of the phase space. Our results confirm a prediction of Donaldson for the spectrum of the Q -operator on Kähler manifolds with constant scalar curvature, and yield exponential convergence of Donaldson’s iterations to the fixed point. Furthermore, viewing POVMs as data clouds, we study their spectral features via geometry of measure metric spaces and the diffusion distance.

Keywords: Berezin–Toeplitz quantization, Berezin transform, Laplace–Beltrami operator, balanced metric, Positive Operator Valued Measure.

2020 *Mathematics Subject Classification*: 53D50, 81P15.

DOI: <https://doi.org/10.5802/ahl.63>

(*) L.I. partially supported by the European Research Council Starting grant 757585. V.K., L.P. and D.S. partially supported by the European Research Council Advanced grant 338809.

RÉSUMÉ. — Nous étudions la transformée de Berezin, un opérateur de Markov associé aux mesures à valeur opérateurs positifs (POVM), dans un certain nombre de contextes incluant la quantification de Berezin–Toeplitz, le système dynamique de Donaldson sur l’espace des produits Hermitiens d’un espace vectoriel complexe, les représentations de groupes finis, et le bruit quantique. En particulier, nous calculons le trou spectral de la quantification en termes de l’harmonique fondamentale de l’espace des phases. Nos résultats confirment une prédiction de Donaldson à propos du spectre de l’opérateur Q sur les variétés de Kähler à courbure scalaire constante, et impliquent la convergence exponentielle du système dynamique de Donaldson vers son point fixe. De plus, en regardant un POVM comme un nuage de points, nous étudions ses propriétés spectrales à travers la géométrie des espaces métriques mesurés et la distance de diffusion.

1. Introduction

Given a function f on a classical phase space X , let us first quantize it and then dequantize. This operation on functions, $f \mapsto \mathcal{B}f$, is called *the Berezin transform*. As a result of this operation, the function f blurs on the phase space. The intuition behind this is as follows⁽¹⁾: assume that f is the Dirac delta-function at a point $x \in X$. Its quantization is a coherent state at x , whose dequantization is approximately a Gaussian centered at x . In the framework of the Berezin–Toeplitz quantization of closed Kähler manifolds, \mathcal{B} is known to be a Markov operator with finite-dimensional image, and is closely related to the Laplace–Beltrami operator Δ of the Kähler manifold. In fact, the Berezin transform has the following asymptotic expansion as $\hbar \rightarrow 0$, due to Karabegov and Schlichenmaier [KS01]⁽²⁾:

$$(1.1) \quad \mathcal{B}_\hbar(f) = f - \frac{\hbar}{4\pi} \Delta f + \mathcal{O}(\hbar^2),$$

for every smooth function f on X , with remainder depending on f and where \hbar stands for the Planck constant (see Section 3 for notations and conventions).

We focus on the spectral properties of \mathcal{B} . For fixed \hbar , this operator factors through a finite-dimensional space and hence its spectrum consists of a finite collection of points lying in the interval $[0, 1]$. Moreover, multiplicities of positive eigenvalues are finite, and 1 is the maximal eigenvalue corresponding to the constant function. Write its spectrum (with multiplicities) in the form

$$1 = \gamma_0 \geq \gamma_1 \geq \gamma_2 \geq \dots \geq \gamma_k \geq \dots \geq 0.$$

The quantity $\gamma := 1 - \gamma_1$ is called *the spectral gap*, a fundamental characteristic of a Markov chain responsible for the rate of convergence to the stationary distribution. Our first result, Theorem 3.1, implies that in the context of the Berezin–Toeplitz quantization, the spectral gap γ of the Berezin transform equals

$$(1.2) \quad \gamma = \frac{\hbar}{4\pi} \lambda_1 + \mathcal{O}(\hbar^2),$$

⁽¹⁾ We thank S. Nonnenmacher for this explanation.

⁽²⁾ Note that after renormalization, there is a missing factor of $\frac{1}{2}$ in front of the second term of the analogous formula in [KS01, (1.2)].

where λ_1 stands for the first eigenvalue of Δ . Note that the upper bound on the gap readily follows from (1.1). The proof follows a work of Lebeau and Michel [LM10] on semiclassical random walks on manifolds with extra ingredients such as an asymptotic expansion for the Bergman kernel due to Dai, Liu and Ma [DLM06], a comparison between the Berezin transform and the heat operator motivated by the work of Liu and Ma [LM07], and a refined version of the above-mentioned Karabegov–Schlichenmaier asymptotic expansion [KS01]. In fact, Theorem 3.1 shows much more than (1.2), namely that one can approximate the full spectrum of Δ , as well as the associated eigenfunctions, with those of \mathcal{B} .

Let us point out that the proof of Theorem 3.1 can be extended to Berezin–Toeplitz quantization of closed symplectic manifolds, using the quantum spaces given by the eigenstates corresponding to the small eigenvalues of the renormalized Bochner Laplacian. This uses the associated generalized Bergman kernel of Ma and Marinescu [MM08] and asymptotic estimates refining those of Ma, Marinescu and Lu [LMM17] (see the discussion at the end of Section 3).

The Berezin transform is defined in the more general context of positive operator valued measures (POVMs). In fact, the Berezin–Toeplitz quantization is nothing else but the integration over a certain POVM on the phase space M with values in the space of quantum observables, and the dequantization is the dual operation [CP18b, Lan98]. In addition to quantization, POVMs appear in quantum mechanics in another setting: they model quantum measurements [BLPY16]. Interestingly enough, within this model the spectral gap of the Berezin transform corresponding to a POVM admits two different interpretations: it measures the minimal magnitude of quantum noise production, and it equals the spectral gap of the Markov chain corresponding to repeated quantum measurements (see Section 7 for details).

Another theme of this paper is related to Donaldson’s program [Don09] of developing approximate methods for detecting canonical metrics on Kähler manifolds. Interestingly enough, our study of the Berezin transform yields the asymptotic behaviour of the spectrum and of the eigenfunctions of the Q -operator, a geometric operator arising in this program, for Kähler metrics of constant scalar curvature. This behaviour, which was predicted by Donaldson in [Don09], is stated in Theorem 3.2 below.

Additionally, Donaldson discovered in [Don09] a remarkable class of dynamical systems on the space of all Hermitian products on a given complex vector space. Section 4 deals with the spectrum of the linearization of such a system at a fixed point. We show that it can be identified with the quantum channel associated to a certain POVM. Using the positivity of the associated spectral gap and under certain natural assumptions, we prove that this linearization is contracting, which confirms Donaldson’s prediction via numerical computations in [Don09, Section 3]. By the Grobman–Hartman theorem and earlier results of Donaldson, this implies in particular that the iterations of this system converge exponentially fast to its fixed point (see Theorem 4.4), and not only for “almost all initial conditions”, as predicted in [Don09, Section 4.1]. The use of Hartman’s theorem in a related context has been suggested by Fine in [Fin12b].

This naturally brings us, in Section 5, to a geometric viewpoint at POVMs. Following Oreshkov and Calsamiglia [OC09, VII.C], we encode them as probability measures in the space of quantum states \mathcal{S} equipped with the Hilbert–Schmidt metric. It turns out that the spectral gap admits a transparent description in terms of the geometry of such metric measure spaces and exhibits a robust behaviour under perturbations of POVMs in the Wasserstein metric. In a similar spirit, one can consider a POVM as a data cloud in \mathcal{S} , which leads us to a link between the spectral gap and the diffusion distance, a notion coming from geometric data analysis.

Section 6 contains a case study of POVMs associated to irreducible unitary representations of finite groups. In this case the spectrum of the Berezin transform and the diffusion distance associated to the corresponding Markov chain can be calculated explicitly via the character table of the group, and their properties reflect algebraic features. In particular, we prove that any non-trivial irreducible representation of a simple group has a strictly positive spectral gap (see Corollary 6.6).

2. Preliminaries

The mathematical model of quantum mechanics starts with a complex Hilbert space \mathcal{H} . In what follows we consider finite-dimensional Hilbert spaces only. Observables are represented by Hermitian operators whose space is denoted by $\mathcal{L}(\mathcal{H})$. Quantum states are provided by *density operators*, i.e., positive trace-one operators $\rho \in \mathcal{L}(\mathcal{H})$. They form a subset $\mathcal{S}(\mathcal{H}) \subset \mathcal{L}(\mathcal{H})$. *Notation:* We write $((A, B))$ for the scalar product $\text{tr}(AB^*) = \text{tr}(AB)$ on $\mathcal{L}(\mathcal{H})$.

Let Ω be a set equipped with a σ -algebra \mathcal{C} of its subsets. By default, we assume that Ω is a Polish topological space (i.e., it is homeomorphic to a complete metric space possessing a countable dense subset) and \mathcal{C} is the Borel σ -algebra.

An $\mathcal{L}(\mathcal{H})$ -valued *positive operator valued measure* W on (Ω, \mathcal{C}) , which we abbreviated to POVM, is a countably additive map $W: \mathcal{C} \rightarrow \mathcal{L}(\mathcal{H})$ which takes each subset $X \in \mathcal{C}$ to a positive operator $W(X) \in \mathcal{L}(\mathcal{H})$ and which is normalized by $W(\Omega) = \mathbb{1}$. According to [CDS07], every $\mathcal{L}(\mathcal{H})$ -valued POVM possesses a density with respect to some probability measure α on (Ω, \mathcal{C}) , that is having the form

$$(2.1) \quad dW(s) = nF(s)d\alpha(s) ,$$

where $n = \dim_{\mathbb{C}} \mathcal{H}$ and $F: \Omega \rightarrow \mathcal{S}(\mathcal{H})$ is a measurable function.

A POVM W given by formula (2.1) is called *pure* if

- (i) for every $s \in \Omega$ the state $F(s)$ is pure, i.e. a rank one projector;
- (ii) the map $F: \Omega \rightarrow \mathcal{S}(\mathcal{H})$ is one to one.

Pure POVMs, under various names, arise in several areas of mathematics including the Berezin–Toeplitz quantization, convex geometry (see [GM00] for the notion of an isotropic measure and [AS17] for the resolution of identity associated to John and Löwner ellipsoids), signal processing (see [EF02] for a link between tight frames and quantum measurements) and Hamiltonian group actions [FM05]. When Ω is a finite set, a pure POVM with a given measure α exists if and only if the measure $\alpha(\{s\})$

of each point $s \in \Omega$ is $\leq \frac{1}{n}$, see [FM05] for a detailed account on the structure of the moduli spaces of pure POVMs on finite sets up to unitary conjugations.

Let us introduce the main character of our story, the spectral gap of a POVM of the form (2.1). Define a map $T : L_1(\Omega, \alpha) \rightarrow \mathcal{L}(\mathcal{H})$ by

$$T(\phi) = \int_{\Omega} \phi \, dW = n \int_{\Omega} \phi(s) F(s) d\alpha(s) .$$

(here and below we work with spaces of real-valued functions). The dual map $T^* : \mathcal{L}(\mathcal{H}) \rightarrow L_{\infty}(\Omega, \alpha)$ is given by $T^*(A)(s) = n((F(s), A))$. Since $L_{\infty} \subset L_1$, we have an operator

$$(2.2) \quad \mathcal{E} = \frac{1}{n} T T^* : \mathcal{L}(\mathcal{H}) \rightarrow \mathcal{L}(\mathcal{H}) ,$$

$$\mathcal{E}(A) = n \int_{\Omega} ((F(s), A)) F(s) d\alpha(s) .$$

Observe that \mathcal{E} is a unital trace-preserving completely positive map. In the terminology of [Hay06, Example 5.4], this is an example of an entanglement-breaking quantum channel.

Furthermore, set

$$(2.3) \quad \mathcal{B} = \frac{1}{n} T^* T : L_1(\Omega, \alpha) \rightarrow L_{\infty}(\Omega, \alpha) ,$$

$$\mathcal{B}(\phi)(t) = n \int_{\Omega} \phi(s) ((F(s), F(t))) d\alpha(s) .$$

Observe that the image of \mathcal{B} is finite-dimensional as \mathcal{B} factors through $\mathcal{L}(\mathcal{H})$.

Write $(\phi, \psi) := \int_{\Omega} \phi \psi \, d\alpha$ for the scalar product on $L_2(\Omega, \alpha)$, and $\|\cdot\|$ for the associated norm. Note that \mathcal{B} is defined as an operator on $L_2(\Omega, \alpha)$ and its spectrum belongs to $[0, 1]$, with 1 being the maximal eigenvalue associated with the constant function.

Note now that positive eigenvalues of \mathcal{E} and \mathcal{B} coincide. Indeed, T^* maps isomorphically an eigenspace corresponding to a positive eigenvalue of \mathcal{E} to the eigenspace of \mathcal{B} corresponding to the same eigenvalue. Write

$$1 = \gamma_0 \geq \gamma_1 \geq \gamma_2 \geq \gamma_k \geq \dots \geq 0$$

for the eigenvalues of \mathcal{B} with multiplicities.

DEFINITION 2.1. — *The non-negative number*

$$\gamma(W) := 1 - \gamma_1 \geq 0$$

is called the spectral gap of the POVM W .

With slight abuse of terminology, we sometimes refer to $\gamma(W)$ as the spectral gap of the operators \mathcal{B} and \mathcal{E} .

Several aspects of this paper concern the positivity of such a spectral gap, and are related to the theory of Markov chains with state space Ω . The notion of the spectral gap, while seemingly being unnoticed in the context of POVMs, naturally appears in the study of Markov chains, where it is responsible for the rate of convergence to the stationary measure. In Section 3, Markov chains will provide a useful link

between Berezin–Toeplitz quantization and semiclassical random walks studied by Lebeau and Michel [LM10], to get a semi-classical estimation of the spectral gap of the Berezin–Toeplitz POVM. In fact, the result of Lebeau and Michel can be applied directly to get a lower bound on this spectral gap, giving a weak version of Theorem 3.1. Finally, as we have already mentioned, POVMs play a central role in the mathematical theory of quantum measurements, and, interestingly enough, Markov chains arise in the context of repeated quantum measurements, see Section 7. Let us recall some basic notions from the theory of Markov chains [BGL13, Rud13]. A *Markov kernel* on Ω is a map $x \mapsto \sigma_x$ sending a point $x \in \Omega$ to a probability measure σ_x on (Ω, \mathcal{C}) such that $x \mapsto \sigma_x(A)$ is a measurable function for every $A \in \mathcal{C}$. With every Markov kernel σ one associates a Markov chain, i.e., a sequence of Ω -valued random variables ζ_k , $k = 0, 1, \dots$ defined on the same probability space, such that for every n and every sequence $x_i \in \Omega$ the conditional probabilities satisfy

$$\mathbb{P}(\zeta_n \mid \zeta_{n-1} = x_{n-1}, \dots, \zeta_0 = x_0) = \sigma_{x_{n-1}}.$$

If ζ_0 is distributed according to a probability measure ν_0 on Ω , then ζ_1 is distributed according to ν_1 given by the formula

$$\nu_1(A) = \int_{\Omega} \sigma_x(A) d\nu_0(x), \quad \forall A \in \mathcal{C}.$$

If $\nu_0 = \nu_1$, we say that ν_0 is a stationary measure for the Markov chain.

The Markov kernel is called *reversible* with respect to a measure ν on Ω if

$$d\nu(x) d\sigma_x(y) = d\sigma_y(x) d\nu(y),$$

as measures on $\Omega \times \Omega$. In this case ν is a stationary measure of the Markov chain. Given a ν -reversible Markov kernel σ with the state space Ω , define *the Markov operator* \mathcal{A} on $L_1(\Omega, \nu)$ by

$$(2.4) \quad \mathcal{A}(\phi)(x) = \int_{\Omega} \phi(y) d\sigma_x(y).$$

Note that \mathcal{A} preserves positivity: $\mathcal{A}(\phi) \geq 0$ for $\phi \geq 0$, $\mathcal{A}(1) = 1$, and its operator norm is ≤ 1 . The reversibility readily yields that the Markov operator \mathcal{A} is self-adjoint on $L_2(\Omega, \nu)$. Denote by 1^\perp the orthogonal complement to the constant function 1 on Ω , i.e., the space of functions with zero mean. Then \mathcal{A} preserves 1^\perp . By definition, the spectral gap $\gamma(\mathcal{A})$ is defined as

$$(2.5) \quad \gamma(\mathcal{A}) = 1 - \|\mathcal{A}|_{1^\perp}\| = \inf_{\phi \neq 0} \frac{(\phi - \mathcal{A}\phi, \phi)}{(\phi, \phi) - (\phi, 1)^2}.$$

With this language, the operator \mathcal{B} given by (2.3) is a Markov operator with the Markov kernel

$$(2.6) \quad t \mapsto n((F(s), F(t))) d\alpha(s).$$

It is reversible with respect to the stationary measure α .

3. Spectral gap for quantization

3.1. Berezin transform vs. Laplace–Beltrami operator

Pure POVMs naturally appear in the context of Berezin–Toeplitz quantization of closed Kähler manifolds (X, ω) , which are *quantizable* in the sense that the cohomology class $[\omega]$ of the Kähler symplectic form $\omega \in \Omega^2(X, \mathbb{R})$ is integral. Recall that this last condition is equivalent to the existence of a holomorphic Hermitian line bundle (L, h) over X whose Chern connection has curvature $-2\pi i\omega$.

Let us briefly recall the construction of this quantization (see [BMS94, LF18, Sch10] for preliminaries). Let X be a quantizable closed Kähler manifold with $\dim_{\mathbb{C}} X = d$, and let (L, h) be a holomorphic Hermitian line bundle as above. Write L^p for the p^{th} tensor power of L , and h^p for the Hermitian metric on L^p induced by h , for any $p \in \mathbb{N}^*$ ⁽³⁾. Then the Hilbert space of quantum states in the space \mathcal{H}_p of global holomorphic sections of L^p , together with the L_2 -inner product induced by the Hermitian metric h^p on L^p and the *Liouville measure* dv_X associated to the canonical volume form $\omega^{\frac{d}{2}}$. We set $n_p = \dim_{\mathbb{C}} \mathcal{H}_p$. The quantity $\hbar = \frac{1}{p}$ plays the role of the Planck constant, so that the classical limit is given by $p \rightarrow +\infty$. For all $p \in \mathbb{N}^*$ large enough, we define a pure $\mathcal{L}(\mathcal{H}_p)$ -valued POVM on X through its density (2.1) by the formula

$$(3.1) \quad dW_p = n_p F_p d\alpha_p,$$

where the map $F_p : X \rightarrow \mathcal{S}(\mathcal{H}_p)$ sends a point $x \in X$ to the *coherent state projector* with kernel the space of sections vanishing at $x \in X$, and where the measure α_p is given at any $x \in X$ by

$$(3.2) \quad d\alpha_p(x) = \frac{R_p(x)}{n_p} dv_X(x),$$

with density $R_p : X \rightarrow \mathbb{R}$ called the *Rawnsley function*. From the viewpoint of complex geometry, the map F_p is given by the Kodaira map and the Rawnsley function is given by the value on the diagonal of the Bergman kernel, i.e. the Schwarz kernel with respect to dv_X of the orthogonal projection $\Pi_p : L^2(X, L^p) \rightarrow \mathcal{H}_p$. By the Kodaira embedding theorem, for all $p \in \mathbb{N}^*$ large enough, the map F_p is well defined and injective, and we have $R_p(x) \neq 0$ for all $x \in X$, so that the $\mathcal{L}(\mathcal{H}_p)$ -valued measure W_p defines a pure POVM in the sense of Section 2, called the *Berezin–Toeplitz POVM*.

In this context, the operator $\mathcal{B}_p := \frac{1}{n_p} T_p^* T_p$ given by formula (2.3) is known as the *Berezin transform*. Recall that for any $p \in \mathbb{N}^*$, the operator \mathcal{B}_p has a finite-dimensional image, and all its eigenvalues lie in the interval $[0, 1]$. There is a finite number of positive eigenvalues with multiplicities, while 0 has infinite multiplicity. Write

$$1 = \gamma_{0,p} \geq \gamma_{1,p} \geq \gamma_{2,p} \geq \dots \geq \gamma_{k,p} \geq \dots \geq 0$$

for the eigenvalues of \mathcal{B}_p with multiplicities.

⁽³⁾ Our convention is that the set of natural numbers \mathbb{N} contains 0. We write \mathbb{N}^* for strictly positive natural numbers.

Let $\Delta f = -\operatorname{div} \nabla f$ be the (positive) Laplace–Beltrami operator associated with the Kähler metric, acting on functions on X with eigenvalues

$$(3.3) \quad 0 = \lambda_0 < \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_k \leq \dots$$

THEOREM 3.1. — *For every integer $k \in \mathbb{N}$, we have the following asymptotic estimate as $p \rightarrow +\infty$,*

$$(3.4) \quad 1 - \gamma_{k,p} = \frac{1}{4\pi p} \lambda_k + \mathcal{O}(p^{-2}) .$$

Furthermore, every sequence in $p \in \mathbb{N}^*$ of $L_2(X, \alpha_p)$ -normalized eigenfunctions of \mathcal{B}_p corresponding to the eigenvalue $\gamma_{k,p}$ contains a subsequence converging to an eigenfunction of the Laplace–Beltrami operator corresponding to λ_k in the \mathcal{C}^∞ -sense.

The proof of Theorem 3.1 is given in Section 3.5. Note that in the context of Section 2, Theorem 3.1 is equivalent to the same statement via T_p^* for the operator $\mathcal{E}_p : \mathcal{L}(\mathcal{H}_p) \rightarrow \mathcal{L}(\mathcal{H}_p)$ defined from \mathcal{B}_p by the formula (2.2). Let us emphasize also that the remainder $\mathcal{O}(p^{-2})$ in (3.4) is not uniform in k .

The Berezin transform \mathcal{B}_p and its associated operator \mathcal{E}_p have prominent cousins, the $Q_{K,p}$ -operator and the Q_p -operator, respectively introduced by Donaldson [Don09, Section 4] in the framework of his program of finding numerical approximation to distinguished Kähler metrics on complex projective manifolds. They are defined as

$$(3.5) \quad \begin{aligned} Q_{K,p} &= \frac{\operatorname{Vol}(X)}{n_p} \iota_p T_p : L_1(X) \rightarrow L_\infty(X) , \\ Q_p &= \frac{\operatorname{Vol}(X)}{n_p} T_p \iota_p : \mathcal{L}(\mathcal{H}_p) \rightarrow \mathcal{L}(\mathcal{H}_p) , \end{aligned}$$

where for any $p \in \mathbb{N}^*$, the map $\iota_p : \mathcal{L}(\mathcal{H}_p) \rightarrow L_\infty(X)$ has been defined in [Don09, Section 2.2.1] for all $A \in \mathcal{L}(\mathcal{H}_p)$ and $x \in X$ by the formula

$$(3.6) \quad \iota_p(A)(x) = \sum_{j=1}^{n_p} h^p(As_j(x), s_j(x)) ,$$

where $\{s_j\}_{j=1}^{n_p}$ is an orthonormal basis of \mathcal{H}_p . By definition of the coherent state projector $F_p : X \rightarrow \mathcal{S}(\mathcal{H}_p)$ and in the language of Section 2, equation (3.6) reads

$$(3.7) \quad \iota_p(A)(x) = \frac{1}{n_p} R_p(x) T_p^*(A)(x) .$$

On the other hand, by their definitions (3.5), the non-vanishing parts of the spectra of $Q_{K,p}$ and Q_p are finite and coincide together with their multiplicities. Write the eigenvalues of Q_p as

$$(3.8) \quad \beta_{0,p} \geq \beta_{1,p} \geq \beta_{2,p} \geq \dots \geq \beta_{k,p} \geq \dots ,$$

and set

$$p' := \left(\frac{n_p}{\operatorname{Vol}(X)} \right)^{\frac{1}{d}} .$$

For some Kähler metrics of constant scalar curvature, Donaldson considered the Q_p -operator as a finite-dimensional approximation of the heat operator and predicted

(see [Don09, p. 611]) that as $p \rightarrow +\infty$, the spectrum of Q_p approximate the spectrum of $e^{-\frac{\Delta}{4\pi p'}}$, and an approximation to the eigenfunctions of $e^{-\frac{\Delta}{4\pi p'}}$ can be extracted from the eigenvectors of Q_p . The following result, which follows from Theorem 3.1 using the classical asymptotics of the Rawnsley function as $p \rightarrow +\infty$, confirms Donaldson's prediction for all Kähler metrics of constant scalar curvature. A detailed proof is given in Section 3.5.

THEOREM 3.2. — *Assume that the Kähler metric of X has constant scalar curvature. For every integer $k \in \mathbb{N}$, we have the following asymptotic estimate as $p \rightarrow +\infty$,*

$$(3.9) \quad 1 - \beta_{k,p} = \frac{1}{4\pi p'} \lambda_k + \mathcal{O}(p^{-2}) .$$

Furthermore, for every sequence in $\{A_p\}_{p \in \mathbb{N}^*}$ of normalized eigenvectors of Q_p in $\mathcal{L}(\mathcal{H}_p)$ corresponding to the eigenvalue $\beta_{k,p}$ for all $p \in \mathbb{N}^*$, there is a subsequence of

$$(3.10) \quad \left\{ \frac{\iota_p A_p}{\|\iota_p A_p\|_p} \right\}_{p \in \mathbb{N}^*}$$

converging to an eigenfunction of the Laplace–Beltrami operator corresponding to λ_k in the \mathcal{C}^∞ -sense, where $\|\cdot\|_p$ is the norm on $L_2(X, \alpha_p)$.

We refer to [Fin12b, KMS16] for a related study of the asymptotic behaviour of the spectrum of certain geometric operators arising in Donaldson's program.

Let us introduce the following useful notion [Don09, Fin12a].

DEFINITION 3.3. — *Let (X, ω) be a closed Kähler manifold, and let (L, h) be a Hermitian holomorphic line bundle over X whose Chern connection has curvature $-2\pi i \omega$. Fix a positive integer $p \in \mathbb{N}^*$ so that the Kodaira map $X \rightarrow H^0(X, L^p)$ is an embedding. We say that the data (X, L^p, h^p) is balanced if the corresponding Rawnsley function $R_p : X \rightarrow \mathbb{R}$ is constant.*

Note that for the balanced data (X, L^p, h^p) the Berezin transform \mathcal{B}_p and the $Q_{K,p}$ -operator coincide, as well as \mathcal{E}_p and the Q_p -operator. In that case, the result of Theorem 3.1 is relevant in [Don09, Section 4.3]. We refer the reader to [Don09, Section 4.1] and to [Fin10, Section 1.4.1] for an interpretation of these operators in terms of complex geometry of (X, L^p) . Let us finally mention that the approximation of the heat operator by the Q_K -operator has been explored by Liu and Ma in [LM07], and that the analogue of the refined Karabegov–Schlichenmaier expansion of Proposition 3.8 for the Q_K -operator has been shown by Ma and Marinescu in [MM12, Theorem 6.1]. Some ingredients of their approach are instrumental for us.

It follows from Theorem 3.1 that the spectral gap of the Berezin–Toeplitz POVM equals

$$(3.11) \quad \gamma(W_p) = \frac{\hbar}{4\pi} \lambda_1 + \mathcal{O}(\hbar^2) , \quad \hbar = \frac{1}{p} .$$

In particular, this yields that the eigenvalue 1 of \mathcal{B}_p is simple (i.e., has multiplicity 1) for all sufficiently large p .

Example 3.4. — Take the projective line $X = \mathbb{C}P^1 = S^2$ of area 1. Let $L = \mathcal{O}(1)$ be the holomorphic line bundle over X dual to the tautological one. The quantum Hilbert space \mathcal{H}_p of global holomorphic sections of L^p can be identified with the $(p+1)$ -dimensional space of homogeneous polynomials of degree p of 2 variables. A representation-theoretical argument (see [Zha98, Don09] and Remark 6.7 below) shows that the eigenvalue γ_1 of the Berezin transform equals $\frac{p}{p+2}$. The Kähler metric on X has constant curvature. For such metrics the first eigenvalue λ_1 of the Laplace–Beltrami operator equals $\frac{8\pi}{\text{Area}} = \frac{8}{\pi}$. We get that

$$\gamma = 1 - \gamma_1 = \frac{2}{p+2} = \frac{1}{4\pi p} \lambda_1 + \mathcal{O}(p^{-2}) ,$$

as predicted by (3.11).

The upper bound in (3.11) immediately follows from the Karabegov–Schlichenmaier asymptotic expansion (1.1) of the Berezin transform [KS01]

$$\mathcal{B}_p(f) = f - \frac{1}{4\pi p} \Delta f + \mathcal{O}(p^{-2}) ,$$

for every smooth function f on X , where the remainder $\mathcal{O}(p^{-2})$ depends on f . Indeed, choosing f to be the $L_2(X, \alpha_p)$ -normalized first eigenfunction of Δ , we see that

$$\gamma(W_p) \leq ((1 - \mathcal{B}_p)f, f)_p \leq \frac{1}{4\pi p} \lambda_1 + \mathcal{O}(p^{-2}) ,$$

where $(\cdot, \cdot)_p$ is the scalar product on $L_2(X, \alpha_p)$.

The prototypical example illustrating a link between the Berezin transform and the Laplace–Beltrami operator is the flat space \mathbb{R}^{2n} , where the Berezin transform \mathcal{B}_p simply coincides with the heat operator $e^{\frac{-h\Delta}{4\pi}}$ (see [Ber75]). It would be interesting to explore the following problem motivated by a conversation with J.-M. Bismut. Denote by $\chi(t)$ the indicator function of the interval $[0, 1]$.

PROBLEM 3.5. — Call a non-decreasing sequence $r(p)$ in $p \in \mathbb{N}^*$ admissible if

$$\left\| (\mathcal{B}_p - e^{\frac{-\Delta}{4\pi p}}) \chi \left(\frac{\Delta}{r(p)} \right) \right\| = \mathcal{O}(p^{-2}) ,$$

where the norm stands for the operator norm in L_2 . According to Theorem 3.1, the constant sequence $r(p) = C$ is admissible for all C . Is the sequence $r(p) = p^\tau$ admissible for $\tau > 0$? What is the maximal possible growth rate of an admissible sequence?

Let us finally make a couple of comments on the physical intuition behind the Berezin transform. It has been noted in the introduction that the Berezin transform can be defined as the composition of the quantization and the dequantization. It is instructive to interpret it in terms of the quantization only. Let σ be a classical state, i.e. a Borel probability measure on X , and following [CP18a], define its quantization as

$$\Theta_p(\sigma) = \int_X F(x) d\sigma(x) ,$$

where as earlier $F(x)$ stands for the coherent state projector at $x \in X$. Let further $f \in L_2(X)$ be a classical observable. It was noticed in [CP18a, (11)] that the expectation $((T_p(f), \Theta_p(\sigma)))$ of the value of the quantized observable $T_p(f)$ in the quantized state $\Theta_p(\sigma)$ equals the classical expectation $\int_X \mathcal{B}(f) d\sigma$ of the Berezin transform $\mathcal{B}(f)$ in the classical state σ . Thus in the context of Berezin–Toeplitz quantization, we get another interpretation of the blurring of quantization measured by \mathcal{B} . Furthermore, in view of Theorem 3.1, we know that \mathcal{B} is a Markov operator with strictly positive spectral gap. Thus it has unique stationary measure α_p whose density against the phase volume is given by $\frac{R_p}{n_p}$, as in formula (3.2). Interestingly enough, this provides an interpretation of the Rawnsley function without appealing to a specific choice of coherent states.

3.2. Comments on the proof

The proof of Theorem 3.1 occupies the rest of this section, and we will deduce Theorem 3.2 as a consequence of it in Section 3.5. Our argument has the same structure as the one in a paper by Lebeau and Michel [LM10] on the Markov operator associated to the semiclassical random walk on manifolds. The key intermediate results are as follows:

- (i) An apriori estimate stating that for any eigenfunction f of \mathcal{B}_p whose eigenvalue is sufficiently bounded away from 0, *any* Sobolev norm $\|f\|_{H^q}$ is bounded by $C_q \|f\|_{L_2}$. See Lemma 3.10 below which is a counterpart of [LM10, Lemma 5].
- (ii) The operators $\mathcal{A}_p := p(\mathbb{1} - \mathcal{B}_p)$ and $\frac{\Delta}{4\pi}$ turn out to be $\sim p^{-1}$ -close as operators from L_2 to H^q for the Sobolev space H^q with *some* sufficiently large q , see formula (3.41) below which is a counterpart of [LM10, formula (3.28)], and which can be considered as a refinement of the expansion (1.1) obtained in [KS01].

Combining (i) and (ii) we conclude that, roughly speaking, eigenfunctions of \mathcal{A}_p as in (i) are “approximate” eigenfunctions of the Laplacian, which eventually implies that the spectra of \mathcal{A}_p and Δ are close to one another, which yields the desired result (see the ending of our proof which is parallel to the one in [LM10]).

Proving (i) and (ii) forms the main bulk of the work. In contrast to [LM10], our proof does not involve micro-local analysis. The main ingredients we use is the expansion of the Bergman kernel due to Dai, Liu and Ma [DLM06] (see Theorem 3.7) and a comparison between the Berezin transform and the heat operator motivated by the work of Liu and Ma on Donaldson’s Q_K -operator [LM07] (see Proposition 3.9 below).

Finally, an acknowledgment is in order. After a weaker version of Theorem 3.1 was posted and formula (3.11) was stated as a question, Alix Deleporte kindly shared with us his ideas concerning the proof of (3.11). He sent us notes [Del] containing a number of preliminary steps in the direction of (i) and (ii) above. While the original arguments of Deleporte dealt with the case of real-analytic Kähler manifolds and line bundles and were based on the asymptotic expansion from [RSVN18, Del], he informed us that they also could be adjusted to the \mathcal{C}^∞ -case.

3.3. Preparations

Recall that the measure dv_X associated to the canonical volume form $\frac{\omega^d}{d!}$ is also the Riemannian volume form of X . Let $\langle \cdot, \cdot \rangle_{L_2}$ be the usual L_2 -scalar product on $\mathcal{C}^\infty(X, \mathbb{C})$, and let $\| \cdot \|_{L_2}$ be the associated norm. For all $j \in \mathbb{N}$, let $e_j \in \mathcal{C}^\infty(X, \mathbb{C})$ be the normalized eigenfunction associated with the j^{th} eigenvalue of the Laplace–Beltrami operator, so that $\|e_j\|_{L_2} = 1$ and $\Delta e_j = \lambda_j e_j$ as in (3.3) for all $j \in \mathbb{N}$. Then for any $f \in \mathcal{C}^\infty(X, \mathbb{C})$, we have the following equality in L_2 ,

$$(3.12) \quad f = \sum_{j=0}^{+\infty} \langle f, e_j \rangle_{L_2} e_j.$$

For any $F : \mathbb{R} \rightarrow \mathbb{R}$ bounded, we define the bounded operator $F(\Delta)$ acting on $L_2(X, \mathbb{C})$ by the formula

$$(3.13) \quad F(\Delta)f = \sum_{i=0}^{+\infty} F(\lambda_i) \langle f, e_i \rangle_{L_2} e_i.$$

The bounded operator $e^{-t\Delta}$ thus defined for all $t > 0$ is called the *heat operator*. For any $m \in \mathbb{N}$, let $|\cdot|_{\mathcal{C}^m}$ be a \mathcal{C}^m norm on $\mathcal{C}^\infty(X, \mathbb{C})$. The following result is classical and can be found for example in [Kan77], [BGV92, Theorem 2.29(2.8)].

PROPOSITION 3.6. — *For any $m \in \mathbb{N}$, there exists $C_m > 0$ such that for any $f \in \mathcal{C}^\infty(X, \mathbb{C})$ and all $t > 0$, we have*

$$(3.14) \quad |e^{-t\Delta}f - f + t\Delta f|_{\mathcal{C}^m} \leq C_m t^2 |f|_{\mathcal{C}^{m+4}}.$$

For any $m \in \mathbb{N}^*$, let $\| \cdot \|_{H^m}$ be a Sobolev norm of order m on $\mathcal{C}^\infty(X, \mathbb{C})$. Using the elliptic estimates for the Laplace–Beltrami operator, for m even we define $\| \cdot \|_{H^m}$ by

$$(3.15) \quad \|f\|_{H^m} := \left\| \Delta^{\frac{m}{2}} f \right\|_{L_2} + \|f\|_{L_2}.$$

Note that the Laplacian Δ is symmetric with respect to the corresponding scalar product on H^m . By convention, we set $\|f\|_{H^0} := \|f\|_{L_2}$.

Next, turn to the Berezin transform. Recall that the Hermitian product on L and the Riemannian measure dv_X induce an L_2 -scalar product on sections of L^p for any $p \in \mathbb{N}^*$, and write $L_2(X, L^p)$ for the associated Hilbert space. The central tool for the study of the Berezin transform is the Schwartz kernel $\Pi_p(x, y)$ of the orthogonal projector $\Pi_p : L_2(X, L^p) \rightarrow \mathcal{H}_p$, called the *Bergman kernel*. Recall that for fixed x and y , this is an element of $L_x^p \otimes \bar{L}_y^p$, where L_x^p denotes the fiber of L^p at $x \in X$ and the bar stands for the conjugate line bundle. Since the bundle L comes with a Hermitian metric, we can measure the point-wise norm $|\Pi_p(x, y)|$. By [LF18, Corollary 9.1.4(2)], we have that $|\Pi_p(x, y)| = |\langle \xi_{x,p}, \xi_{y,p} \rangle|$, where $\xi_{x,p}$ is the non-normalized *coherent state* at $x \in X$ defined up to a phase factor (see e.g. [CP18b, LF18] for the definition). The Rawnsley function R_p is given by $R_p(x) = |\xi_{x,p}|^2$, and thus satisfies $R_p(x) = \Pi_p(x, x)$. Since $F_p(x)$ is the projector to $\xi_{x,p}$, we have that

$$|\Pi_p(x, y)|^2 = ((F_p(x), F_p(y))) R_p(x) R_p(y).$$

It follows from (2.6) and (3.2) that

$$(3.16) \quad \begin{aligned} (\mathcal{B}_p f)(x) &= n_p \int_X ((F_p(x), F_p(y))) f(y) d\alpha_p(y) \\ &= \frac{1}{R_p(x)} \int_X |\Pi_p(x, y)|^2 f(y) dv_X(y) , \end{aligned}$$

so that the Schwarz kernel of \mathcal{B}_p with respect to dv_X is given by

$$(3.17) \quad \mathcal{B}_p(x, y) = \frac{|\Pi_p(x, y)|^2}{R_p(x)} .$$

Let $\|\cdot\|_p$ be the norm on $L_2(X, \alpha_p)$. From the classical asymptotic expansion of R_p as $p \rightarrow +\infty$, we get a constant $C > 0$ such that

$$(3.18) \quad \left(\frac{1}{\text{Vol}(X)} - Cp^{-1} \right) \|\cdot\|_{L_2} \leq \|\cdot\|_p \leq \left(\frac{1}{\text{Vol}(X)} + Cp^{-1} \right) \|\cdot\|_{L_2} .$$

3.4. Asymptotic expansion of the Berezin transform

For a comprehensive account on the off-diagonal expansion of the Bergman kernel as well as tools of Berezin–Toeplitz quantization in this context, we refer to [MM07].

We always assume that $p \in \mathbb{N}^*$ is as large as needed. For any $s > 0$, we use the notation $O(p^{-s})$ as $p \rightarrow +\infty$ in the usual sense, uniformly in \mathcal{C}^m -norm for all $m \in \mathbb{N}^*$. The notation $O(p^{-\infty})$ means $O(p^{-s})$ for any $s > 0$.

Let $\varepsilon_0 > 0$ be smaller than the injectivity radius of X . Fix a point $x_0 \in X$, and let $Z = (Z_1, \dots, Z_{2d}) \in \mathbb{R}^{2d}$ with $|Z| < \varepsilon_0$ be geodesic normal coordinates around x_0 , where $|\cdot|$ is the Euclidean norm of \mathbb{R}^{2d} . In these coordinates, the canonical volume form is given by

$$(3.19) \quad dv_X(Z) = \kappa_{x_0}(Z) dZ ,$$

with $\kappa_{x_0}(0) = 1$. For any kernel $K(\cdot, \cdot) \in \mathcal{C}^\infty(X \times X, \mathbb{C})$, we write $K_{x_0}(\cdot, \cdot)$ for its image in these coordinates, and we write $|K_x|_{\mathcal{C}^m(X)}$ for the \mathcal{C}^m -norm of the family of functions K_x with respect to $x \in X$.

Let d^X be the Riemannian distance on X . We will derive Theorem 3.1 as a consequence of the following asymptotic expansion as $p \rightarrow +\infty$ of the Schwartz kernel of the Berezin transform.

THEOREM 3.7. — *For any $m, k \in \mathbb{N}$ and $\varepsilon > 0$, there is $C > 0$ such that for all $p \in \mathbb{N}^*$ and $x, y \in X$ satisfying $d^X(x, y) > \varepsilon$,*

$$(3.20) \quad |\mathcal{B}_p(x, y)|_{\mathcal{C}^m} \leq Cp^{-k} .$$

For any $m, k \in \mathbb{N}$, there is $N \in \mathbb{N}$ and $C > 0$ such that for any $x_0 \in X$, $|Z|, |Z'| < \varepsilon_0$ and for all $p \in \mathbb{N}^*$, we have

$$(3.21) \quad \left| p^{-d} B_{p,x_0}(Z, Z') - \sum_{r=0}^{k-1} p^{-\frac{r}{2}} J_{r,x_0}(\sqrt{p}Z, \sqrt{p}Z') \exp(-\pi p|Z - Z'|^2) \kappa_{x_0}^{-1}(Z') \right|_{\mathcal{C}^m(X)} \\ \leq C p^{-\frac{k}{2}} (1 + \sqrt{p}|Z| + \sqrt{p}|Z'|)^N \exp\left(-\sqrt{p} \left| \frac{Z - Z'}{C} \right| \right) + O(p^{-\infty}),$$

where $\{J_{r,x_0}(Z, Z')\}_{r \in \mathbb{N}}$ is a family of polynomials in $Z, Z' \in \mathbb{R}^{2n}$ of the same parity as r , depending smoothly on $x_0 \in X$. Furthermore, for any $Z, Z' \in \mathbb{R}^{2n}$ we have

$$(3.22) \quad J_{0,x_0}(Z, Z') = 1 \quad \text{and} \quad J_{1,x_0}(Z, Z') = 0.$$

This readily follows from formula (3.17) expressing the Schwarz kernel of the Berezin transform via the Bergman kernel Π_p and the analogous result of Dai, Liu and Ma in [DLM06, Theorem 4.18'] for the Bergman kernel.

For any $x \in X$, let $B^X(x, \varepsilon_0)$ be the geodesic ball of radius $\varepsilon_0 > 0$ around x , and write $B(0, \varepsilon_0) \subset \mathbb{R}^{2d}$ for the Euclidean ball of radius ε_0 around 0. The following proposition is a refinement of the Karabegov–Schlichenmaier expansion [KS01, (1.2)] of the Berezin transform, where we make explicit the remainder term.

PROPOSITION 3.8. — *For any $m \in \mathbb{N}$, there exists $C_m > 0$ such that for any $f \in \mathcal{C}^\infty(X, \mathbb{C})$ and all $p \in \mathbb{N}^*$, we have*

$$(3.23) \quad \left| \mathcal{B}_p f - f + \frac{\Delta}{4\pi p} f \right|_{\mathcal{C}^m} \leq \frac{C_m}{p^2} |f|_{\mathcal{C}^{m+4}}.$$

Proof. — For any $x \in X$, write f_x for the image of f restricted to $B^X(x, \varepsilon_0)$ in normal coordinates around x . From (3.20), we know that for any $\varepsilon > 0$ and $x \in X$,

$$(3.24) \quad (\mathcal{B}_p f)(x) = \int_X \mathcal{B}_p(x, y) f(y) dv_X(y) \\ = \int_{B^X(x, \varepsilon_0)} \mathcal{B}_p(x, y) f(y) dv_X(y) + O(p^{-\infty}) |f|_{\mathcal{C}^0} \\ = \int_{B(0, \varepsilon_0)} B_{p,x}(0, Z) f_x(Z) \kappa_x(Z) dZ + O(p^{-\infty}) |f|_{\mathcal{C}^0}.$$

For any $k \in \mathbb{N}^*$ and $m \in \mathbb{N}$, we will use the following Taylor expansion of f_x up to order $k - 1$, for all $p \in \mathbb{N}^*$ and $|Z| < \varepsilon_0$,

$$(3.25) \quad f_x(Z) = \sum_{0 \leq |\alpha| \leq k-1} \frac{\partial^{|\alpha|} f_x}{\partial Z^\alpha} \frac{Z^\alpha}{\alpha!} + O_m(|Z|^k) |f|_{\mathcal{C}^{m+k}} \\ = \sum_{0 \leq |\alpha| \leq k-1} p^{-\frac{|\alpha|}{2}} \frac{\partial^{|\alpha|} f_x}{\partial Z^\alpha} \frac{(\sqrt{p}Z)^\alpha}{\alpha!} + p^{-\frac{k}{2}} O_m(|\sqrt{p}Z|^k) |f|_{\mathcal{C}^{m+k}},$$

where O_m means that the expansion is uniform in $x \in X$ as well as all its derivatives up to order $m \in \mathbb{N}$, and does not depend on f .

We will compute the asymptotic expansion as $p \rightarrow +\infty$ of (3.24) using the Taylor expansion (3.25) of f and the asymptotic expansion (3.21) of the Berezin transform up to order 3. First, using the fact that $\mathcal{B}_p 1 = 1$ for all $p \in \mathbb{N}^*$, we know that the polynomials $J_{r,x}(Z, Z')$ of the asymptotic expansion (3.21) of the Berezin transform satisfy

$$(3.26) \quad \int_{\mathbb{R}^{2n}} J_{r,x}(0, Z) \exp(-\pi p|Z|^2) dZ = 0 ,$$

for all $x \in X$ and $r \in \mathbb{N}^*$. On another hand, recall from (3.22) that $J_{0,x} \equiv 1$ and $J_{1,x} \equiv 0$ for all $x \in X$. Using the parity of Gaussian functions, a change of variable $Z \mapsto \frac{Z}{\sqrt{p}}$ and the Taylor expansion (3.25) for $k = 4$, we get that

$$(3.27) \quad \begin{aligned} p^d \int_{B(0, \varepsilon_0)} \exp(-\pi p|Z|^2) f_x(Z) dZ \\ = f(x) + p^{-1} \sum_{j=1}^{2n} \frac{\partial^2 f_x}{\partial Z_j^2}(0) \int_{\mathbb{R}^{2n}} \frac{Z_j^2}{2} \exp(-\pi|Z|^2) dZ + |f|_{\mathcal{C}^{m+4}} O_m(p^{-2}) \\ = f(x) - p^{-1} \frac{\Delta}{4\pi} f(x) + |f|_{\mathcal{C}^{m+4}} O_m(p^{-2}) . \end{aligned}$$

Recall that $J_{r,x}(0, Z) \in \mathbb{C}[Z]$ is a polynomial in $Z \in \mathbb{R}^{2n}$ of the same parity than $r \in \mathbb{N}$, so that using (3.25), (3.26) and the parity of Gaussian functions, we get in the same way

$$(3.28) \quad \begin{aligned} p^d \int_{B(0, \varepsilon_0)} J_{2,x}(0, \sqrt{p}Z) \exp(-\pi p|Z|^2) f_x(Z) dZ \\ = f(x) \int_{\mathbb{R}^{2n}} J_{2,x}(0, Z) \exp(-\pi|Z|^2) dZ + O_m(p^{-1}) |f|_{\mathcal{C}^{m+2}} \\ = O_m(p^{-1}) |f|_{\mathcal{C}^{m+2}} , p^d \int_{B(0, \varepsilon_0)} J_{3,x}(0, \sqrt{p}Z) \exp(-\pi p|Z|^2) f_x(Z) dZ \\ = O_m(p^{-\frac{1}{2}}) |f|_{\mathcal{C}^{m+1}} . \end{aligned}$$

Finally, again using a change of variable $Z \mapsto \frac{Z}{\sqrt{p}}$, we get for any $N \in \mathbb{N}^*$ and $p \in \mathbb{N}^*$,

$$(3.29) \quad p^d \int_{B(0, \varepsilon_0)} (1 + |\sqrt{p}Z|)^N \exp\left(\frac{-\sqrt{p}|Z|}{C}\right) f_x(Z) dZ = O_m(1) |f|_{\mathcal{C}^m} .$$

This completes the proof of (3.23). \square

In view of Propositions 3.6 and 3.8, it is natural to compare the Berezin transform with the heat operator by setting $t = (4\pi p)^{-1}$. This leads to the following result, which is essentially a refinement of [LM07, Theorem 0.1].

PROPOSITION 3.9. — *For any $m \in \mathbb{N}$, there exists $C_m > 0$ such that for any $f \in \mathcal{C}^\infty(X, \mathbb{C})$ and all $p \in \mathbb{N}^*$, we have*

$$(3.30) \quad \left\| \left(e^{-\frac{\Delta}{4\pi p}} - \mathcal{B}_p \right) f \right\|_{H^m} \leq \frac{C_m}{p} \|f\|_{H^m} .$$

Proof. — Set $S_p := e^{\frac{\Delta}{4\pi p}} - \mathcal{B}_p$, which acts on $L_2(X, \mathbb{C})$ for all $p \in \mathbb{N}^*$ and admits a smooth Schwartz kernel $S_p(\cdot, \cdot)$ with respect to dv_X . Comparing Theorem 3.7 with the classical asymptotic expansion of the heat kernel, as given for example in [BGV92, Theorem 2.29], [Kan77], we see that

$$(3.31) \quad S_p(x, y) = O(p^{-\infty}),$$

for all $x, y \in X$ satisfying $d^X(x, y) > \varepsilon_0$, and using the formula (3.22) for the first two coefficients, we get for any $m \in \mathbb{N}$ a constant $C > 0$ and $N \in \mathbb{N}$ such that

$$(3.32) \quad |S_{p, x_0}(Z, Z')|_{\mathcal{C}^m(X)} \leq Cp^{-1}(1 + \sqrt{p}|Z| + \sqrt{p}|Z'|)^N \exp\left(\frac{-\sqrt{p}|Z - Z'|}{C}\right) + O(p^{-\infty}).$$

Let us first show (3.30) for $m = 0$. For any $f \in \mathcal{C}^\infty(X, \mathbb{C})$ and any $\varepsilon > 0$, by Cauchy–Schwarz inequality and (3.31) for S_p , we get the following version of the Schur test for all $p \in \mathbb{N}^*$,

$$(3.33) \quad \begin{aligned} \|S_p f\|_{L_2}^2 &\leq \int_X \left(\int_X |S_p(x, y)| dv_X(y) \right) \left(\int_X |S_p(x, y)| |f(y)|^2 dv_X(y) \right) dv_X(x) \\ &\leq \sup_{x \in X} \left(\int_X |S_p(x, y)| dv_X(y) \right) \sup_{y \in X} \left(\int_X |S_p(x, y)| dv_X(x) \right) \|f\|_{L_2}^2 \\ &\leq \sup_{x \in X} \left(\int_{B(x, \varepsilon_0)} |S_p(x, y)| dv_X(y) \right) \sup_{y \in X} \left(\int_{B(x, \varepsilon_0)} |S_p(x, y)| dv_X(x) \right) \|f\|_{L_2}^2 \\ &\quad + O(p^{-\infty}) \|f\|_{L_2}^2. \end{aligned}$$

Then (3.30) for $m = 0$ follows from (3.32) with $Z = 0$ or $Z' = 0$ respectively, as in (3.29).

To deal with the case of arbitrary $m \in \mathbb{N}^*$, let us assume by induction that (3.30) is satisfied for $m - 1$. Considering the estimates (3.31) and (3.32) with corresponding $m \in \mathbb{N}^*$, note that for any differential operator D_x of order m in $x \in X$, there exists a differential operator $D'_{x,y}$ in $x, y \in X$ of total order m but of order at most $m - 1$ in $x \in X$, such that the operator $S_p^{(m)}$ defined through its kernel for all $x, y \in X$ by

$$(3.34) \quad S_p^{(m)}(x, y) := D_x S_p(x, y) + D'_{x,y} S_p(x, y)$$

also satisfies (3.31) and (3.32). Then for all $x \in X$ and $p \in \mathbb{N}^*$, we get

$$(3.35) \quad \begin{aligned} \int_X D_x S_p(x, y) f(y) dv_X(y) &= - \int_X (D'_{x,y} S_p(x, y)) f(y) dv_X(y) + (S_p^{(m)} f)(x) \\ &= \int_X D'_x S_p(x, y) (D''_y f(y)) dv_X(y) + (S_p^{(m)} f)(x), \end{aligned}$$

where D'_x and D''_y are differential operators, respectively in x and in y , obtained from $D'_{x,y}$ using a partition of unity and integration by parts in local charts, so that in particular D'_x is of order $m - 1$ in $x \in X$. Then using the induction hypothesis, the inequality (3.30) for m follows from the same inequality for $m - 1$ replacing f by any number of derivatives of f , and from the estimates (3.32) and (3.33) for $S_p^{(m)}$ in the same way than before. \square

3.5. Spectrum

Recall that $\|\cdot\|_p$ denotes the norm on $L_2(X, \alpha_p)$. In this section, we consider a sequence $\{f_p\}_{p \in \mathbb{N}^*}$, with $f_p \in \mathcal{C}^\infty(X, \mathbb{C})$ such that

$$(3.36) \quad \|f_p\|_p = 1, \quad \mathcal{B}_p f_p = \mu_p f_p,$$

for some $\mu_p \in \text{Spec}(\mathcal{B}_p)$ for all $p \in \mathbb{N}^*$. The following estimate is crucial for the proof of Theorem 3.1.

LEMMA 3.10. — *Assume that the sequence $\{p(1 - \mu_p)\}_{p \in \mathbb{N}^*}$ is bounded by some constant $L > 0$. Then for all $m \in \mathbb{N}$, there exists $C_{L,m} > 0$ such that for all $p \in \mathbb{N}^*$, we have*

$$(3.37) \quad \|f_p\|_{H^{2m}} \leq C_{L,m}.$$

Proof. — Note that (3.37) is automatically verified for $m = 0$ by (3.18) and (3.36). By induction on $m \in \mathbb{N}$, let us assume that (3.37) is satisfied for $m - 1$. Let us write

$$(3.38) \quad \begin{aligned} p \left(e^{-\frac{\Delta}{4\pi p}} - \mathcal{B}_p \right) f_p &= p(1 - \mu_p) f_p - p \left(\mathbb{1} - e^{-\frac{\Delta}{4\pi p}} \right) f_p \\ &= p(1 - \mu_p) f_p - \Delta F \left(\frac{\Delta}{p} \right) f_p, \end{aligned}$$

where the bounded operator $F(\Delta/p)$ acting on $L_2(X, \mathbb{C})$ is defined as in (3.13) for the continuous function $F : \mathbb{R} \rightarrow \mathbb{R}$ given for any $s \in \mathbb{R}^*$ by $F(s) = 4\pi(1 - e^{-s/4\pi})/s$. As $|p(1 - \mu_p)| < L$ for all $p \in \mathbb{N}^*$, by Proposition 3.9 and formula (3.15) for $\|\cdot\|_{H^{2m}}$, this gives a constant $C_m > 0$ such that

$$(3.39) \quad \left\| F \left(\frac{\Delta}{p} \right) f_p \right\|_{H^{2m}} \leq C_m \|f_p\|_{H^{2m-2}}.$$

On the other hand, note that by hypothesis, we have $\mu_p \rightarrow 1$ as $p \rightarrow +\infty$. Using Proposition 3.9 again, we then get $\varepsilon_m > 0$ and $p_m \in \mathbb{N}^*$ such that for all $p > p_m$,

$$(3.40) \quad \begin{aligned} \left\| F \left(\frac{\Delta}{p} \right) f_p \right\|_{H^{2m}} &\geq \left\| F \left(\frac{\Delta}{p} \right) f_p + \left(\mathcal{B}_p - e^{-\frac{\Delta}{4\pi p}} \right) f_p \right\|_{H^{2m}} - \left\| \left(\mathcal{B}_p - e^{-\frac{\Delta}{4\pi p}} \right) f_p \right\|_{H^{2m}} \\ &\geq \inf_{s>0} \left\{ F(s) + \mu_p - e^{-\frac{s}{4\pi}} \right\} \|f_p\|_{H^{2m}} - C_m p^{-1} \|f_p\|_{H^{2m}} \\ &\geq \varepsilon_m \|f_p\|_{H^{2m}}. \end{aligned}$$

This together with (3.39) gives (3.37). \square

Proof of Theorem 3.1. — For any $f \in \mathcal{C}^\infty(X, \mathbb{C})$, by Proposition 3.8, we get that

$$(3.41) \quad \left\| p(\mathbb{1} - \mathcal{B}_p) f - \frac{\Delta}{4\pi} f \right\|_{L_2} \leq C p^{-1} |f|_{\mathcal{C}^4} \leq C p^{-1} \|f\|_{H^q},$$

with q even and large enough. The inequality on the right follows from Sobolev embedding theorem, and the same is true in $L_2(X, \alpha_p)$ -norm by (3.18). Let now

$j \in \mathbb{N}$ be fixed. If $e_j \in \mathcal{C}^\infty(X, \mathbb{C})$ satisfies $\Delta e_j = \lambda_j e_j$ and $\|e_j\|_{L_2} = 1$, then by (3.41) we get $C_j > 0$ not depending on $p \in \mathbb{N}^*$ such that

$$(3.42) \quad \left\| p(\mathbb{1} - \mathcal{B}_p)e_j - \frac{\lambda_j}{4\pi} e_j \right\|_p \leq C_j p^{-1}.$$

Thus if $m_j \in \mathbb{N}$ is the multiplicity of λ_j as an eigenvalue of Δ , the estimate (3.42) for all eigenfunctions of Δ associated with λ_j gives a constant $C > 0$ such that

$$(3.43) \quad \# \left(\text{Spec} \left(p(\mathbb{1} - \mathcal{B}_p) \right) \cap \left[\frac{\lambda_j}{4\pi} - Cp^{-1}, \frac{\lambda_j}{4\pi} + Cp^{-1} \right] \right) \geq m_j.$$

This immediately follows from the variational principle for the operator

$$p(\mathbb{1} - \mathcal{B}_p) - \frac{\lambda_j}{4\pi} \mathbb{1}$$

acting on $L_2(X, \alpha_p)$.

Consider now for every $p \in \mathbb{N}^*$ a normalized eigenfunction $f_p \in \mathcal{C}^\infty(X, \mathbb{C})$ of \mathcal{B}_p as in (3.36) such that the associated sequence $\{p(1 - \mu_p)\}_{p \in \mathbb{N}^*}$ of eigenvalues of $p(\mathbb{1} - \mathcal{B}_p)$ is bounded. Combining Lemma 3.10 with the right inequality in (3.41), we get $C > 0$ such that

$$(3.44) \quad \|p(1 - \mu_p)f_p - \Delta f_p\|_{L_2} \leq Cp^{-1}.$$

In particular, we get that

$$(3.45) \quad \text{dist}(p(1 - \mu_p), \text{Spec } \Delta) \leq Cp^{-1}.$$

Finally, let us show that there exists $p_0 \in \mathbb{N}^*$ such that (3.43) is in fact an equality for $p > p_0$. To this end, let $l \in \mathbb{N}^*$ with $l \geq m_j$ be such that for all $p \in \mathbb{N}^*$, there exists an orthonormal family $f_{k,p}$, $1 \leq k \leq l$, of eigenfunctions of \mathcal{B}_p in $L_2(X, d\alpha_p)$ with associated eigenvalues $\mu_{k,p} \in \mathbb{R}$, $1 \leq k \leq l$, satisfying

$$(3.46) \quad p(1 - \mu_{k,p}) \in \left[\lambda_j - Cp^{-1}, \lambda_j + Cp^{-1} \right], \text{ for all } 1 \leq k \leq l.$$

As the inclusion of the Sobolev space H^q in H^{q-1} is compact, by Lemma 3.10 there exists a subsequence of $\{f_{k,p}\}_{p \in \mathbb{N}^*}$ converging to a function f_k in H^{q-1} -norm, for all $1 \leq k \leq l$. In particular, taking $q > 2$, the family f_k , $1 \leq k \leq l$, is orthonormal in $L_2(X, \mathbb{C})$ and satisfies $\Delta f_k = \lambda_j f_k$ for all $1 \leq k \leq l$ by (3.44). By definition of the multiplicity $m_j \in \mathbb{N}$ of λ_j , this forces $l = m_j$.

Let us sum up our findings. First, the equality

$$\# \left(\text{Spec} \left(p(\mathbb{1} - \mathcal{B}_p) \right) \cap \left[\frac{\lambda_j}{4\pi} - Cp^{-1}, \frac{\lambda_j}{4\pi} + Cp^{-1} \right] \right) = m_j,$$

where m_j is the multiplicity of λ_j as the eigenvalue of Δ , together with (3.45) readily yields the first statement of the theorem:

$$1 - \gamma_{k,p} = \frac{1}{4\pi p} \lambda_k + \mathcal{O}(p^{-2}).$$

Second, observe that we got a subsequence of $f_{k,p}$, $p \in \mathbb{N}^*$ converging to f_k in the Sobolev H^{q-1} sense, where q even can be chosen arbitrarily large. By the Sobolev

embedding theorem, this yields a subsequence which \mathcal{C}^l -converges to f_k with arbitrary l . Iterating this argument for this subsequence we get that there exists a sequence $p_l \rightarrow +\infty$ such that

$$|f_{k,p_l} - f_k|_{\mathcal{C}^l} \leq \frac{1}{l},$$

which means that f_{k,p_l} converges to f_k in the \mathcal{C}^∞ -sense. This completes the proof. \square

Proof of Theorem 3.2. — For any $p \in \mathbb{N}^*$, using equation (3.7) for ι_p and combining the definition (3.5) of the $Q_{K,p}$ -operator with formula (3.16) for the Berezin transform $\mathcal{B}_p = \frac{1}{n_p} T_p^* T_p$ acting on $f \in \mathcal{C}^\infty(X, \mathbb{C})$, we get

$$(3.47) \quad (Q_{K,p}f)(x) = \frac{\text{Vol}(X)}{n_p} R_p(x) \mathcal{B}_p(f)(x) = \frac{\text{Vol}(X)}{n_p} \int_X |\Pi_p(x, y)|^2 f(y) dv_X(y).$$

We will show that when the scalar curvature is constant, the analogue of Theorem 3.1 holds for this operator. As $\frac{p'}{p} = 1 + \mathcal{O}(p^{-1})$ by the Riemann–Roch theorem, this will imply Theorem 3.2 via the morphism ι_p which relates Q_p with $Q_{K,p}$, see (3.5).

Recall that $R_p : X \rightarrow \mathbb{R}$ denotes the Rawnsley function, and that $n_p = \dim_{\mathbb{C}} \mathcal{H}_p$. By the classical asymptotic expansion of the Bergman kernel, which can be found for example in [MM07, Section 4.1.1], we know that when the scalar curvature is constant, we have

$$(3.48) \quad \frac{\text{Vol}(X)}{n_p} R_p = 1 + \mathcal{O}(p^{-2}).$$

As this expansion holds in \mathcal{C}^m -norm for all $m \in \mathbb{N}^*$ and by the definition \mathcal{B}_p and $Q_{K,p}$ in formulas (3.16) and (3.47) respectively, we get a constant $C_m > 0$ for any $m \in \mathbb{N}^*$ such that

$$(3.49) \quad \|Q_{K,p} - \mathcal{B}_p\|_{H^m} \leq C_m p^{-2}.$$

It is then easy to see that Lemma 3.10 holds for any sequence $\{f_p\}_{p \in \mathbb{N}^*}$ with $f_p \in \mathcal{C}^\infty(X, \mathbb{C})$ such that

$$(3.50) \quad \|f_p\|_{L_2} = 1, \quad Q_{K,p}f_p = \mu_p f_p,$$

with $\{p(1 - \mu_p)\}_{p \in \mathbb{N}^*}$ bounded, simply using the estimate (3.49) to replace \mathcal{B}_p by $Q_{K,p}$ in (3.38) and (3.40). We can then follow the proof of Theorem 3.1 above to get the same result for $Q_{K,p}$, using the estimate (3.49) to replace \mathcal{B}_p by $Q_{K,p}$ in (3.41) and (3.42), and using (3.18) to replace $\|\cdot\|_p$ by $\|\cdot\|_{L_2}$ in (3.42). Finally, the form (3.10) for the normalized sequence of eigenfunction of $Q_{K,p}$ follows from the fact that $\iota_p Q_p = Q_{K,p} \iota_p$ by definition (3.5) of Q_p and $Q_{K,p}$. This completes the proof of Theorem 3.2. \square

Remark 3.11. — Theorem 3.1 can be extended to the case of a general closed symplectic manifold (X, ω) of real dimension $2d$, and (L, h, ∇) a Hermitian line bundle with Hermitian connection ∇ of curvature $-2\pi i\omega$. In fact, one can in general consider the following *renormalized Bochner Laplacian* acting on $\mathcal{C}^\infty(X, L^p)$ for any $p \in \mathbb{N}^*$, first introduced by Guillemin and Uribe [GU88],

$$(3.51) \quad \Delta_p := \Delta^{L^p} - 2\pi d p,$$

where Δ^{L^p} stands for the usual Bochner Laplacian on L^p . By [GU88, Theorem 2.a], the spectrum of Δ_p is contained in $I \cup (C_1 p - C_2, +\infty)$ for all $p \in \mathbb{N}^*$, for some $C_1, C_2 > 0$ and some interval $I \subset \mathbb{R}$ containing 0. We can then consider Π_p as the associated spectral projection corresponding to I and set $\mathcal{H}_p = \text{Im}(\Pi_p)$. Using the work [MM08] of Ma and Marinescu on the kernel of Π_p , we can then consider the Berezin–Toeplitz POVM of Section 3.1. By [LMM17, (2.31), (3.2)], the Berezin transform admits an asymptotic expansion similar to Theorem 3.7, except for the formula (3.22), where we only have $J_{1,x_0}(0, Z') = 0$ for all $Z' \in \mathbb{R}^{2d}$ as a consequence of [ILMM17, Lemma 6.1, Lemma 6.2] (see also [MM08, (2.32)]). Then Propositions 3.8 and 3.9 hold, and it is straightforward to adapt the rest of the proof of Theorem 3.1 in Section 3.5. Note that the corresponding estimates in Proposition 3.8 and 3.9 can be seen as refinements of [LMM17].

Remark 3.12. — On the other hand, Theorem 3.1 can be extended to the case of *weighted Berezin transforms*, introduced by Engliš in [Eng00] in the case of pseudoconvex domains. This corresponds to the case where one replaces the canonical volume form $\frac{\omega^d}{d!}$ by a general smooth volume form ν in the setting of Section 3.1. In fact, let us consider the Hilbert space $\mathcal{H}_{\nu,p}$ of global holomorphic sections of L^p together with the L_2 -inner product with respect to the measure $d\nu$ instead of the Liouville measure dv_X . Then using the trick of Ma and Marinescu in [MM07, Section 4.1.9], we can define for any $p \in \mathbb{N}^*$ large enough the $\mathcal{L}(\mathcal{H}_{\nu,p})$ -valued POVM

$$(3.52) \quad dW_{\nu,p} = n_p F_{\nu,p} d\alpha_{\nu,p},$$

where $F_{\nu,p} : X \rightarrow \mathcal{S}(\mathcal{H}_{\nu,p})$ is the map sending $x \in X$ to the orthogonal projector with kernel the space of sections vanishing at $x \in X$ and $\alpha_{p,\nu}$ is given by

$$(3.53) \quad d\alpha_{\nu,p}(x) = \frac{R_{\nu,p}(x)}{n_p} d\nu(x),$$

where $R_{\nu,p} : X \rightarrow \mathbb{R}$ is the weighted Rawnsley function, given by the value on the diagonal of the Schwarz kernel with respect to ν of the orthogonal projector operator $\Pi_{\nu,p} : L^2(X, L^p, d\nu) \rightarrow \mathcal{H}_{\nu,p}$. Using [MM07, Section 4.19] again as well as the general version of the expansion Theorem 3.7 given in [DLM06, Theorem 4.18'], the proof of Theorem 3.1 above extends verbatim to this case, to get the estimate

$$(3.54) \quad 1 - \gamma_{\nu,k,p} = \frac{1}{4\pi p} \lambda_k + \mathcal{O}(p^{-2})$$

as $p \rightarrow \infty$, where $\gamma_{\nu,k,p}$ is the k^{th} eigenvalue of the Berezin transform of $W_{\nu,p}$ and λ_k is the k^{th} eigenvalue of the Laplace–Beltrami operator associated with the Kähler metric, for all $k \in \mathbb{N}$. Note in particular that the first term of the right hand side of equation (3.54) does not depend on the choice of the smooth volume form ν . It would be interesting to understand the general mechanism behind this fact, in the spirit of Theorem 5.4.(ii), showing that the spectral gap of $W_{\nu,p}$ is constant up to $O(\frac{1}{p^2})$ under deformations of ν .

4. Berezin transform and Donaldson's iterations

In [Don09] Donaldson, as a part of his program of developing approximate methods for detecting canonical metrics on Kähler manifolds, discovered a remarkable class of dynamical systems on the space of all Hermitian inner products on a given complex vector space. We shall show in this section that the linearization of such a system at a fixed point can be identified with the quantum channel introduced in (2.2) above and prove that under certain natural assumptions, it is injective and has strictly positive spectral gap. Using earlier results by Donaldson, we will then deduce the main result of this section, Theorem 4.4, stating that the iterations of this system converge exponentially fast to the fixed point.

For a complex n -dimensional vector space \mathcal{V} , denote by $\mathcal{P}rod(\mathcal{V})$ the space of Hermitian inner products on \mathcal{V} . Given such a $q \in \mathcal{P}rod(\mathcal{V})$, let $\mathcal{H} := (\mathcal{V}, q)$ be the corresponding Hilbert space, and define a map

$$(4.1) \quad \Phi_q : \mathbb{P}(\mathcal{V}^*) \longrightarrow \mathcal{L}(\mathcal{H})$$

sending a hyperplane $H \subset \mathcal{V}$, naturally seen as an element of $\mathbb{P}(\mathcal{V}^*)$ via the kernel of linear forms, to the unique orthogonal projector $\Phi_q(H) \in \mathcal{L}(\mathcal{H})$ with respect to q satisfying $\text{Ker } \Phi_q(H) = H$.

Let ν be a Borel measure on $\mathbb{P}(\mathcal{V}^*)$, so that $|\nu| := \nu(\mathbb{P}(\mathcal{V}^*)) < \infty$. Following Donaldson [Don09, p. 581], we say that $q \in \mathcal{P}rod(\mathcal{V})$ is ν -balanced if the operator-valued measure

$$(4.2) \quad dW_q(z) := n \Phi_q(z) \frac{d\nu(z)}{|\nu|} ,$$

defines an $\mathcal{L}(\mathcal{H})$ -valued POVM on $\mathbb{P}(\mathcal{V}^*)$ as in (2.1). This translates into the condition

$$(4.3) \quad n \int_{\mathbb{P}(\mathcal{V}^*)} \Phi_q(z) \frac{d\nu(z)}{|\nu|} = \mathbb{1} .$$

Example 4.1. — Consider a Hilbert space $\mathcal{H} = (\mathcal{V}, q)$ with $\dim_{\mathbb{C}} \mathcal{H} = n$, and let W be a pure $\mathcal{L}(\mathcal{H})$ -valued POVM, defined as in formula (2.1). Let us identify the measure α on Ω with a measure on $\mathbb{P}(\mathcal{V}^*)$ via push-forward by the associated map

$$(4.4) \quad F : \Omega \longrightarrow \mathbb{P}(\mathcal{V}^*) ,$$

where $\mathbb{P}(\mathcal{V}^*)$, seen as the set of hyperplanes in \mathcal{V} , is identified with the set of rank one projectors in $\mathcal{S}(\mathcal{H}) \subset \mathcal{L}(\mathcal{H})$ via their kernel as above. It is then an immediate consequence of the definitions that $q \in \mathcal{P}rod(\mathcal{V})$ is α -balanced.

Example 4.2. — As a particular case of Example 4.1, consider the Berezin–Toeplitz POVM W_p on a closed quantizable Kähler manifold X associated to a Hermitian holomorphic line bundle L for $p \in \mathbb{N}^*$ large enough, as in Section 3.1. The associated Hilbert space is $\mathcal{H}_p = (H^0(X, L^p), q_p)$, where $H^0(X, L^p)$ is the space of holomorphic sections of L^p and $q_p \in \mathcal{P}rod(H^0(X, L^p))$ is the L_2 -Hermitian product induced by the Kähler metric. In this case, the map (4.4) is given by the Kodaira embedding

$$(4.5) \quad F_p : X \longrightarrow \mathbb{P}(H^0(X, L^p)^*) ,$$

and we get as a special case of the previous example that $q_p \in \mathcal{P}rod(H^0(X, L^p))$ is α_p -balanced. Then following e.g. [Fin12a, Proposition 8.3] and by formula (3.2) for α_p , the data (X, L^p, h^p) is balanced in the sense of Definition 3.3 if and only if the product q_p is dv_X -balanced.

Example 4.3. — Let X be a complex manifold together with a holomorphic line bundle L over X such that the Kodaira map F_p given by (4.5) is an embedding for $p \in \mathbb{N}^*$ sufficiently large, and let ν be a smooth volume form over X . Then L^p over X is naturally identified with the pullback by F_p of the dual of the tautological line bundle over $\mathbb{P}(H^0(X, L^p)^*)$, and given a Hermitian inner product $q_p \in \mathcal{P}rod(H^0(X, L^p))$ on $H^0(X, L^p)$, we write h_p for the Hermitian metric induced on L^p by the corresponding Fubini–Study metric. Then by e.g. [Don09, p. 581], the product $q_p \in \mathcal{P}rod(H^0(X, L^p))$ is ν -balanced if and only if it coincides up to constant with the L_2 -inner product on $H^0(X, L^p)$ induced by h_p and the measure $d\nu$. On the other hand, following [MM07, Section 4.19, Section 5.1.4], the last assertion of Example 4.2 holds in the same way when one replaces dv_X by $d\nu$, so that the weighted Rawnsley function $R_{\nu,p} : X \rightarrow \mathbb{R}$ of Remark 3.12 is constant if and only if q_p is ν -balanced. This shows that if q_p is ν -balanced, the induced POVM (4.2) coincides with the Berezin–Toeplitz POVM (3.52) weighted by ν of Remark 3.12.

Donaldson proved [Don09, p. 582] (see also an extensive discussion below) that for every $p \in \mathbb{N}^*$ large enough, there always exists a unique ν -balanced Hermitian inner product $q_p \in \mathcal{P}rod(H^0(X, L^p))$. For $p \in \mathbb{N}^*$ large enough, consider the symplectic form ω_p on X obtained by the pull-back under the Kodaira map F_p of the Fubini–Study form on $\mathbb{P}(H^0(X, L^p)^*)$ corresponding to q_p . Equivalently, $-2i\pi\omega_p$ is the Chern curvature of h_p . By [Don09, p. 584] (see also [Kel09]), the sequence $\{\frac{1}{p}\omega_p\}_p$ converges as $p \rightarrow \infty$ to the unique Kähler form ω_∞ in $c_1(L)$ solving the *Calabi problem* $\omega^d = c\nu$, for some $c > 0$. This illustrates the role of ν -balanced products as finite-dimensional approximations of the solution of the Calabi problem.

Under some natural assumptions on the measure ν , the existence of ν -balanced Hermitian inner products was established by Bourguignon, Li and Yau [BLY94], where they use such products to give an upper bound for the first eigenvalue of the Laplacian of complex manifolds embedded in the projective space. This generalizes the seminal work of Hersch [Her70], where he shows that the first eigenvalue of any metric over S^2 is smaller than the one of the round metric, using the notion of balanced product in its simplest form.

Following instead Donaldson in [Don09], let us associate to a measure ν on $\mathbb{P}(\mathcal{V}^*)$ the dynamical system $\mathcal{T}_\nu : \mathcal{P}rod(\mathcal{V}) \rightarrow \mathcal{P}rod(\mathcal{V})$ defined for all $q \in \mathcal{P}rod(\mathcal{V})$ by

$$(4.6) \quad \mathcal{T}_\nu(q) := n \int_{\mathbb{P}(\mathcal{V}^*)} q(\Phi_q(z) \cdot, \cdot) \frac{d\nu(z)}{|\nu|}.$$

Using condition (4.3), we then see that $q \in \mathcal{P}rod(\mathcal{V})$ is ν -balanced if and only if it is a fixed point of \mathcal{T}_ν . Under mild conditions on the measure ν , Donaldson proved that for every initial condition $q_0 \in \mathcal{P}rod(\mathcal{V})$, the iterations $\mathcal{T}_\nu^r(q_0)$ converge to such a fixed point as $r \rightarrow +\infty$, and that this fixed point is unique up to the action of \mathbb{R}_+ on $\mathcal{P}rod(\mathcal{V})$ by scalar multiplication.

The main result of this section is the *exponential convergence* of Donaldson's iteration process to the ν -balanced product, for all initial conditions.

THEOREM 4.4. — *Suppose that the measure ν on $\mathbb{P}(\mathcal{V}^*)$ is supported on a complex subvariety $Y \subset \mathbb{P}(\mathcal{V}^*)$, with ν absolutely continuous on every irreducible component of Y . Assume that*

(i) *for any projective subspace Σ of $\mathbb{P}(\mathcal{V}^*)$, we have*

$$(4.7) \quad \frac{\nu(\Sigma)}{\dim \Sigma + 1} < \frac{|\nu|}{n} ;$$

(ii) *at least one irreducible component of Y is not contained in any proper projective subspace of $\mathbb{P}(\mathcal{V}^*)$.*

Then for any $q_0 \in \mathcal{P}rod(\mathcal{V})$, there exists a ν -balanced product $q \in \mathcal{P}rod(\mathcal{V})$ and constants $C > 0$, $\beta \in (0, 1)$, such that for all $r \in \mathbb{N}$, we have

$$(4.8) \quad \text{dist}(\mathcal{T}_\nu^r(q_0), q) \leq C\beta^r .$$

Note that if Y is irreducible, assumptions (i) and (ii) are satisfied as soon as Y is not contained in a proper projective subspace of $\mathbb{P}(\mathcal{V}^*)$. Thus Theorem 4.4 applies in particular to the important case of Example 4.3, where ν is induced by a smooth volume form over a complex manifold Y embedded in a projective space via Kodaira embedding. Conversely, if the whole variety Y (in contrast with its irreducible components) lies in a proper projective subspace of $\mathbb{P}(\mathcal{V}^*)$, then there exists $u \in \mathcal{V}$ such that $\Phi_q(z)u = 0$ for all $z \in Y$, contradicting condition (4.3), so that there does not exist any ν -balanced Hermitian product.

The proof of Theorem 4.4 will rely on the Propositions 4.6, 4.7 and 4.8 below. In particular, Proposition 4.6 generalizes the result of Donaldson in [Don09, p. 581], which essentially states that the iterations $\mathcal{T}_\nu^r(q_0)$ converge to a ν -balanced product as $r \rightarrow +\infty$ for all $q_0 \in \mathcal{P}rod(\mathcal{V})$ if either Y is a complex variety which is not contained in any proper projective subspace, or Y is a finite collection of points satisfying (i). Specifically, Donaldson's assumption 2 in [Don09, p. 581] is precisely assumption (i) in the case $\dim Y = 0$, and Donaldson's assumption 1 in [Don09, p. 581] is satisfied in the case Y is a complex variety which is not contained in any proper projective subspace, but do not imply assumption (ii) in general. The proof of Proposition 4.6 follows closely the lines of [Don09, p. 581].

On the other hand, the role of Propositions 4.7 and 4.8 in the proof of Theorem 4.4 is based on the key observation, which is a reformulation of [Don09, p. 609], that the linearization of \mathcal{T}_ν at a fixed point $q \in \mathcal{P}rod(\mathcal{V})$ coincides with the quantum channel (2.2) of the POVM (4.2) associated with q .

To see this, let us first choose a base point $q_0 \in \mathcal{P}rod(\mathcal{V})$ and identify (\mathcal{V}, q_0) with $(\mathbb{C}^n, \langle \cdot, \cdot \rangle)$, where $\langle z, w \rangle = \sum_j z_j \bar{w}_j$. Writing $\mathcal{L}(\mathbb{C}^n)_+$ for the set of positive Hermitian $n \times n$ matrices, this identifies $G \in \mathcal{L}(\mathbb{C}^n)_+$ with $q(\cdot, \cdot) := \langle G\cdot, \cdot \rangle \in \mathcal{P}rod(\mathcal{V})$. Next, identify $[z] \in \mathbb{CP}^{n-1}$ with the hyperplane

$$\left\{ w : \langle z, w \rangle = q(G^{-1}z, w) = 0 \right\} .$$

From the definition (4.1) of Φ_q we have

$$\Phi_q([z])\xi = \frac{\langle G\xi, G^{-1}z \rangle}{\langle GG^{-1}z, G^{-1}z \rangle} G^{-1}z = G^{-1} \left(\frac{\langle \xi, z \rangle}{|z|^2} z \right) \cdot \frac{|z|^2}{\langle G^{-1}z, z \rangle} = G^{-1}\Pi_z\xi \cdot \frac{|z|^2}{\langle G^{-1}z, z \rangle},$$

where Π_z denotes the orthogonal projector with respect to $\langle \cdot, \cdot \rangle$ to the line generated by $z \in \mathbb{C}^n \setminus \{0\}$. Thus,

$$q(\Phi_q(z)\xi, \xi) = \langle \Pi_z\xi, \xi \rangle \cdot \frac{|z|^2}{\langle G^{-1}z, z \rangle}.$$

Therefore, we can reformulate the definition (4.6) of \mathcal{T}_ν in coordinates by the formula

$$(4.9) \quad \mathcal{T}_\nu(G) = n \int_{\mathbb{C}P^{n-1}} \Pi_z \frac{|z|^2}{\langle G^{-1}z, z \rangle} \frac{d\nu(z)}{|\nu|}.$$

Recall that the tangent space of $\mathcal{P}rod(\mathcal{V})$ at any $q \in \mathcal{P}rod(\mathcal{V})$ is canonically identified with the space of Hermitian operators $\mathcal{L}(\mathcal{H})$ of $\mathcal{H} := (\mathcal{V}, q)$. Then if $q \in \mathcal{P}rod(\mathcal{V})$ is ν -balanced, so that $\mathcal{T}_\nu(q) = q$, the differential $D_q\mathcal{T}_\nu$ of \mathcal{T}_ν at q acts on $\mathcal{L}(\mathcal{H})$.

LEMMA 4.5. — *For any ν -balanced Hermitian product $q \in \mathcal{P}rod(\mathcal{V})$, the differential of \mathcal{T}_ν at q satisfies $D_q\mathcal{T}_\nu = \mathcal{E}_q$, where \mathcal{E}_q is the quantum channel (2.2) of the associated POVM W_q defined by (4.2).*

Proof. — Let $q \in \mathcal{P}rod(\mathcal{V})$ be a ν -balanced Hermitian product, and let us identify (\mathcal{V}, q) with $(\mathbb{C}^n, \langle \cdot, \cdot \rangle)$ as above. Let $G(t) \in \mathcal{L}(\mathbb{C}^n)_+$ be a path such that $G(0) = \mathbb{1}$. Abbreviating $\dot{G} := \dot{G}(0)$ and using formula (4.9), we get

$$\left. \frac{d}{dt} \right|_{t=0} \mathcal{T}_\nu(G(t)) = n \int_{\mathbb{C}P^n} \Pi_z \frac{\langle \dot{G}z, z \rangle}{|z|^2} \frac{d\nu(z)}{|\nu|}.$$

Recall that $((\cdot, \cdot))$ denotes the natural scalar product on $\mathcal{L}(\mathcal{H})$. Then noticing that $\langle \dot{G}z, z \rangle / |z|^2 = ((\dot{G}, \Pi_z))$, we get

$$\left. \frac{d}{dt} \right|_{t=0} \mathcal{T}_\nu(G(t)) = n \int_{\mathbb{C}P^n} \Pi_z ((\dot{G}, \Pi_z)) \frac{d\nu(z)}{|\nu|},$$

which is precisely formula (2.2) for the quantum channel associated to W_q defined by (4.2), as we have $\Phi_q(z) = \Pi_z$ for all $z \in \mathbb{C}^n \setminus \{0\}$ in the identification of (\mathcal{V}, q) with $(\mathbb{C}^n, \langle \cdot, \cdot \rangle)$. This concludes the proof. \square

Recall that the quantum channel $\mathcal{E}_q : \mathcal{L}(\mathcal{H}) \rightarrow \mathcal{L}(\mathcal{H})$ satisfies $\mathcal{E}_q(\mathbb{1}) = \mathbb{1}$, and that its *spectral gap* is the quantity $\gamma = 1 - \lambda_1$, where

$$(4.10) \quad 1 = \lambda_0 \geq \lambda_1 \geq \lambda_2 \geq \cdots \geq 0$$

is the decreasing sequence of eigenvalues of \mathcal{E}_q . Then Proposition 4.7 establishes the positivity of the spectral gap of $\mathcal{E}_q = D_q\mathcal{T}_\nu$ under assumption (i), showing that Donaldson's prediction in [Don09, Section 4.1] on the largest eigenvalue of the linearization of \mathcal{T}_ν at a ν -balanced product in fact holds for general projective smooth manifolds Y , as assumption (i) is automatically satisfied as soon as Y is not contained in any proper projective subspace.

Proposition 4.8 shows the invertibility of $\mathcal{E}_q = D_q \mathcal{T}_\nu$ under assumption (ii). This is a key assumption in the classical Grobman–Hartman theorem, which we use in Theorem 4.4 to show that the iterations of the dynamical system \mathcal{T}_ν converge exponentially fast to a fixed point. As assumption (ii) is automatically satisfied for a projective smooth manifold Y not contained in any proper projective subspace, this strengthens Donaldson’s prediction in [Don09, Section 4.1] on the asymptotic rate of convergence of the dynamical system \mathcal{T}_ν . Namely, with only the positivity of the spectral gap, we expect that the rate of convergence is exponentially fast for almost all initial conditions, while Theorem 4.4 shows that it actually holds for all initial conditions.

PROPOSITION 4.6. — *Assume that assumption (i) of Theorem 4.4 holds. Then for any $q_0 \in \mathcal{Prod}(\mathcal{V})$, the iterations $\mathcal{T}_\nu^r(q_0)$ converge to a fixed point as $r \rightarrow +\infty$, unique up to the action of \mathbb{R}_+ by scalar multiplication.*

Proof. — Fix $q_0 \in \mathcal{Prod}(\mathcal{V})$, and identify (\mathcal{V}, q_0) with $(\mathbb{C}^n, \langle \cdot, \cdot \rangle)$, where $\langle \cdot, \cdot \rangle$ denotes the canonical Hermitian product of \mathbb{C}^n . Recall that $\mathcal{L}(\mathbb{C}^n)_+$ denotes the set of positive Hermitian $n \times n$ matrices. Following [Don09, p. 582], for any $[z] \in \mathbb{C}P^{n-1}$, let $z \in \mathbb{C}^n$ be a lift of norm 1, and for any $G \in \mathcal{L}(\mathbb{C}^n)_+$, set

$$(4.11) \quad \psi_{[z]}(G) := \log \langle G^{-1}z, z \rangle + \frac{1}{n} \log \det G.$$

This quantity does not depend on the choice of a lift of $[z] \in \mathbb{C}P^{n-1}$ of norm 1, and the second term makes it invariant under multiplication of G by a positive scalar. Given a Borel measure ν on $\mathbb{C}P^{n-1}$, we then define a functional on $\mathcal{L}(\mathbb{C}^n)_+$ by the formula

$$(4.12) \quad \Psi_\nu(G) = \int_{\mathbb{C}P^{n-1}} \psi_{[z]}(G) d\nu([z]),$$

for any $G \in \mathcal{L}(\mathbb{C}^n)_+$. Using formula (4.9), we see that $G \in \mathcal{L}(\mathbb{C}^n)_+$ is a critical point of Ψ_ν if and only if it is a fixed point of \mathcal{T}_ν . Thus to show the existence and uniqueness of such a fixed point up to the action of \mathbb{R}_+ , we can restrict Ψ_ν to the space $\mathcal{L}(\mathbb{C}^n)_+^1$ of positive Hermitian matrices of determinant 1, and it suffices to show that Ψ_ν is strictly convex and proper along any geodesic of $\mathcal{L}(\mathbb{C}^n)_+^1$ for its natural Riemannian metric as a symmetric space. In fact, any strictly convex and proper function over \mathbb{R} has a unique absolute minimum, which is also its unique critical point. Now as two points can always be joined by a geodesic, we conclude in that case that a fixed point of \mathcal{T}_ν on $\mathcal{L}(\mathbb{C}^n)_+^1$ coincide with a minimum of Ψ_ν , which exists and is unique.

Recall that the structure of symmetric space on $\mathcal{L}(\mathbb{C}^n)_+^1$ is given by the map

$$(4.13) \quad \begin{aligned} \mathrm{SL}_n(\mathbb{C}) &\longrightarrow \mathcal{L}(\mathbb{C}^n)_+^1 \\ G &\longmapsto \sqrt{G^*G}, \end{aligned}$$

which realizes $\mathcal{L}(\mathbb{C}^n)_+^1$ as the quotient of the special linear group $\mathrm{SL}_n(\mathbb{C})$ by the special unitary group $\mathrm{SU}(n)$. The usual scalar product $((\cdot, \cdot))$ on the space of $n \times n$ matrices induces a Riemannian metric on $\mathcal{L}(\mathbb{C}^n)_+^1$ through the identification of its tangent space at any point with the space of traceless matrices. By general theory of

symmetric spaces, geodesics are simply the images of 1-parameter groups of $\mathrm{SL}_n(\mathbb{C})$ through the above map, so that up to the action of $\mathrm{SU}(n)$ by conjugation, they are of the form $G_t \in \mathcal{L}(\mathbb{C}^n)_+^1$, with

$$(4.14) \quad G_t = \mathrm{diag}(e^{\lambda_1 t}, e^{\lambda_2 t}, \dots, e^{\lambda_n t}),$$

for all $t \in \mathbb{R}$, where $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$ satisfy $\sum_{j=1}^n \lambda_j = 0$. Now if ν satisfies assumption (i) of Theorem 4.4, its pullback by the action of a unitary matrix also satisfies this assumption, and thus we are reduced to show strict convexity and properness of

$$(4.15) \quad t \mapsto \Psi_\nu(G_t) = \int_{\mathbb{C}P^{n-1}} \log \left(\sum_{j=1}^n e^{\lambda_j t} |z_j|^2 \right) d\nu([z]), \quad t \in \mathbb{R}.$$

Now convexity follows from a direct computation, with strict convexity as long as the total mass of ν is not contained in any projective subspace of $\mathbb{C}P^{n-1}$, which is a straightforward consequence of assumption (i).

Let us now show properness, i.e. that $\Psi_\nu(G_t) \rightarrow +\infty$ when $t \rightarrow \pm\infty$. By considering the geodesic going to the opposite direction, it suffices to show it when $t \rightarrow +\infty$. Consider an irreducible component $Z \subset Y$, and let $k \leq n$ be the largest integer such that Z is contained in the projective subspace

$$(4.16) \quad \Sigma_k := \{[0 : \dots : 0 : z_k : \dots : z_n] \in \mathbb{C}P^{n-1}\} \subset \mathbb{C}P^{n-1}$$

As ν is absolutely continuous over the smooth part of Z , this means in particular that the function $\log |z_k|^2$ restricted to Z is integrable with respect to ν . We thus get a constant $C_Z > 0$ such that

$$(4.17) \quad \int_Z \log \left(\sum_{j=1}^n e^{\lambda_j t} |z_j|^2 \right) d\nu([z]) \geq \int_Z \log (e^{\lambda_k t} |z_k|^2) d\nu([z]) \\ \geq \lambda_k t \nu(Z) - C_Z.$$

For any $k \leq n$, write $\nu_k > 0$ for the total mass of the irreducible components of Y for which k is the largest integer such that they are not contained in Σ_k as above. We then get a constant $C_Y > 0$ such that

$$(4.18) \quad \Psi_\nu(G_t) \geq t \sum_{j=1}^n \lambda_j \nu_j - C_Y.$$

We are thus reduced to show that $\sum_{j=1}^n \lambda_j \nu_j > 0$. Notice now that assumption (i) implies

$$(4.19) \quad \sum_{j=k}^n \nu_j < \frac{n-k}{n} \sum_{j=1}^n \nu_j, \quad \text{for all } 1 \leq k \leq n.$$

Using $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$ and $\sum_{j=1}^n \lambda_j = 0$, we then get

$$\begin{aligned}
 \sum_{j=1}^n \lambda_j \nu_j &= \lambda_0 \sum_{j=1}^n \nu_j + \sum_{k=1}^n (\lambda_k - \lambda_{k-1}) \sum_{j=k}^n \nu_j \\
 (4.20) \quad &> \left(\lambda_0 + \sum_{k=1}^n \frac{n-k}{n} (\lambda_k - \lambda_{k-1}) \right) \sum_{j=1}^n \nu_j \\
 &> \left(\frac{1}{n} \sum_{k=1}^n \lambda_k \right) \sum_{j=1}^n \nu_j = 0.
 \end{aligned}$$

This implies properness.

Let us now show the convergence of iterations of \mathcal{T}_ν to a fixed point. We will first show that \mathcal{T}_ν decreases Ψ_ν , so that iterations have an accumulation point by properness, and we will then show that this accumulation point is in fact a fixed point. First note that for any $G \in \mathcal{L}(\mathbb{C}^n)_+$, using the fact that projectors are of trace 1, formula (4.3), together with (4.9), gives $\text{tr}[\mathcal{T}_\nu(G)G^{-1}] = n$. Using the strict concavity of the logarithm, we thus get

$$\begin{aligned}
 (4.21) \quad \frac{1}{n} \log \det(\mathcal{T}_\nu(G)) - \frac{1}{n} \log \det(G) &= \frac{1}{n} \log \det(\mathcal{T}_\nu(G)G^{-1}) \\
 &\leq \log \left(\frac{\text{tr}[\mathcal{T}_\nu(G)G^{-1}]}{n} \right) = 0,
 \end{aligned}$$

with equality if and only if $\mathcal{T}_\nu(G)G^{-1} = \mathbb{1}$. Thus to show that $\Psi_\nu(\mathcal{T}_\nu(G)) \leq \Psi_\nu(G)$, by definition (4.12) of Ψ_ν , we only need to show that \mathcal{T}_ν decreases the integral against ν of the first term of formula (4.11). Again by concavity of the logarithm, we get

$$\begin{aligned}
 (4.22) \quad \int_{\mathbb{C}P^{n-1}} \log \langle \mathcal{T}_\nu(G)^{-1}z, z \rangle d\nu([z]) &- \int_{\mathbb{C}P^{n-1}} \log \langle G^{-1}z, z \rangle d\nu([z]) \\
 &\leq \log \left(\int_{\mathbb{C}P^{n-1}} \frac{\langle \mathcal{T}_\nu(G)^{-1}z, z \rangle}{\langle G^{-1}z, z \rangle} d\nu([z]) \right) \\
 &\leq \log \left(\frac{1}{n} \text{tr}[\mathcal{T}_\nu(G)\mathcal{T}_\nu(G)^{-1}] \right) = 0,
 \end{aligned}$$

where we used formula (4.9) for $\mathcal{T}_\nu(G)$ together with the fact that $\langle Az, z \rangle = |z|^2 \text{Tr}[\Pi_z A]$ for all $z \in \mathbb{C}^n \setminus \{0\}$ and $A \in \text{End}(\mathbb{C}^n)$. Equations (4.21) and (4.22), together with the definition of Ψ_ν given by formulas (4.11) and (4.12), show that $\Psi_\nu(\mathcal{T}_\nu(G)) \leq \Psi_\nu(G)$ for all $G \in \mathcal{L}(\mathbb{C}^n)_+$.

To conclude, note first that properness over $\mathcal{L}(\mathbb{C}^n)_+^1$ and invariance under the action of \mathbb{R}_+ implies that Ψ_ν is bounded from below over the whole $\mathcal{L}(\mathbb{C}^n)_+$. Thus for any $G_0 \in \mathcal{L}(\mathbb{C}^n)_+$, we get that the decreasing sequence $\{\Psi_\nu(\mathcal{T}_\nu^r(G_0))\}_{r \in \mathbb{N}}$ converges to its lower bound. As both terms in the definition of Ψ_ν are decreasing under iterations of \mathcal{T}_ν by (4.21) and (4.22), we then deduce that $\{\log \det(\mathcal{T}_\nu^r(G_0))\}_{r \in \mathbb{N}}$, thus also $\{\det(\mathcal{T}_\nu^r(G_0))\}_{r \in \mathbb{N}}$, are bounded in \mathbb{R} , and that

$$(4.23) \quad \frac{1}{n} \log \det(\mathcal{T}_\nu^{r+1}(G_0)\mathcal{T}_\nu^r(G_0)^{-1}) \longrightarrow 0, \quad \text{as } r \rightarrow +\infty.$$

Now from properness of Ψ_ν over $\mathcal{L}(\mathbb{C}^n)_+^1$ and boundedness in \mathbb{R} of the sequences $\{\Psi_\nu(\mathcal{T}_\nu^r(G_0))\}_{r \in \mathbb{N}}$ and $\{\det(\mathcal{T}_\nu^r(G_0))\}_{r \in \mathbb{N}}$, we get that the sequence $\{\mathcal{T}_\nu^r(G_0)\}_{r \in \mathbb{N}}$ admits an accumulation point $G_\infty \in \mathcal{L}(\mathbb{C}^n)_+$. On the other hand, by strict concavity of the logarithm, formula (4.23) and the equality case in formula (4.21) imply

$$(4.24) \quad \mathcal{T}_\nu^{r+1}(G_0) \mathcal{T}_\nu^r(G_0)^{-1} \longrightarrow \mathbb{1}, \quad \text{as } r \rightarrow +\infty.$$

We thus get that $G_\infty \in \mathcal{L}(\mathbb{C}^n)_+$ is the unique accumulation point, and satisfies $\mathcal{T}_\nu(G_\infty) = G_\infty$. This concludes the proof of Proposition 4.6. \square

In the following Proposition 4.7, we use the result that a fixed point of \mathcal{T}_ν exists as soon as ν satisfies assumption (i), which was proved in the previous Proposition 4.6.

PROPOSITION 4.7. — *Assume that assumption (i) holds. Then for any ν -balanced product $q \in \mathcal{Prod}(\mathcal{V})$, the associated quantum channel \mathcal{E}_q as in Lemma 4.5 has positive spectral gap.*

Proof. — Let $q \in \mathcal{Prod}(\mathcal{V})$ be a ν -balanced product, and identify (\mathcal{V}, q) with $(\mathbb{C}^n, \langle \cdot, \cdot \rangle)$, so that $\Phi_q(z) = \Pi_z$ for all $z \in \mathbb{C}^n \setminus \{0\}$ in the definition (4.2) of W_q , where Π_z is the orthogonal projector on $[z]$ with respect to $\langle \cdot, \cdot \rangle$. Assume that ν satisfies assumption (i) of Theorem 4.4, and normalize it by setting $\alpha := \nu/|\nu|$. For any $z \in \mathbb{C}^n \setminus \{0\}$, we denote by $[z]$ its class in $\mathbb{C}P^{n-1}$. For any $z, w \in \mathbb{C}^n \setminus \{0\}$, we write

$$(4.25) \quad \mathcal{B}_q([z], [w]) = n \frac{|\langle z, w \rangle|^2}{|z|^2 |w|^2}$$

for the Schwartz kernel with respect to α of the Berezin transform (2.3) on $L_2(\mathbb{C}P^{n-1}, \nu)$ associated with W_q . Let Y_1, \dots, Y_k be the irreducible components of Y . Since $(z, w) \mapsto \langle z, w \rangle$ is holomorphic in z and anti-holomorphic in w , for every $i, j \leq k$, we get that

- (a) either $\mathcal{B}_q([z], [w]) = 0$ for all $([z], [w]) \in Y_i \times Y_j$,
- (b) or $\mathcal{B}_q([z], [w]) \neq 0$ for almost all $([z], [w]) \in Y_i \times Y_j$.

Consider a graph Γ with vertices $1, \dots, k$, where i, j are connected by an edge whenever (b) occurs of $Y_i \times Y_j$. In particular, each i is connected by an edge to itself.

Recall that

$$\int \mathcal{B}_q(x, y) d\alpha(y) = \int \mathcal{B}_q(x, y) d\alpha(x) = 1.$$

Using the Schur test as in formula (3.33) above, we apply Cauchy–Schwarz inequality on the formula

$$(4.26) \quad \int \mathcal{B}_q(x, y) \phi(y) d\alpha(y) = \int \mathcal{B}_q(x, y)^{\frac{1}{2}} \mathcal{B}_q(x, y)^{\frac{1}{2}} \phi(y) d\alpha(y),$$

to get for any $\phi \in L^2(\mathbb{C}P^{n-1}, \nu)$,

$$(4.27) \quad \begin{aligned} \|\mathcal{B}_q \phi\|_{L_2}^2 &= \int \left(\int \mathcal{B}_q(x, y) \phi(y) d\alpha(y) \right)^2 d\alpha(x) \\ &\leq \int \left(\int \mathcal{B}_q(x, y) d\alpha(y) \cdot \int \mathcal{B}_q(x, y) \phi^2(y) d\alpha(y) \right) d\alpha(x) = \|\phi\|_{L_2}. \end{aligned}$$

In particular, the equality $\mathcal{B}_q \phi = \phi$ can hold only if the inequality above is an equality, and by the equality case of Cauchy–Schwarz inequality, this implies that for α -almost all x , there exists $c \neq 0$ such that $c \mathcal{B}_q(x, y)^{1/2} = \mathcal{B}_q(x, y)^{1/2} \phi(y)$ for

α -almost all X . In terms of the graph defined in the previous step, this yields that ϕ is constant on every subset of the form $\bigcup_{j \in \text{star}(i)} Y_j$, where $i = 1, \dots, k$. Thus if ϕ is a non constant function satisfying $\mathcal{B}_q \phi = \phi$, it follows that Γ is disconnected. Denote by Γ_i , $i = 1, \dots, k$ the connected components, and put $Z_i = \bigcup_{j \in \Gamma_i} Y_j$.

Assuming that there exists a non-constant ϕ satisfying $\mathcal{B}_q \phi = \phi$ as above, we will show that assumption (i) can not hold. Recall that we work with the POVM $dW_q(x) = n \Pi_x d\alpha(x)$, where Π_x is the orthogonal projector to the line $x \in \mathbb{C}P^{n-1}$ with respect to $\langle \cdot, \cdot \rangle$. With this notation, $\mathcal{B}_q(x, y) = 0$ yields $\Pi_x \Pi_y = 0$. Write $P = W_q(Z_1)$ and $P' = W_q(Z_2 \cup \dots \cup Z_k)$. It follows that $P + P' = \mathbb{1}$ and $PP' = 0$. Thus P is an orthogonal projector whose image is a proper projective subspace Σ of $\mathbb{C}P^{n-1}$ of dimension $m - 1$, with

$$m = \text{tr } [P] = \text{tr } [W_q(Z_1)] = n \alpha(Z_1) = n \frac{\nu(Z_1)}{|\nu|}.$$

Observe also that if $Pz = 0$, we get

$$\int_{Z_1} \langle \Pi_x z, z \rangle d\nu(x) = 0,$$

and hence $\langle \Pi_x z, z \rangle = 0$ for ν -almost all x . Since ν is absolutely continuous on each irreducible component of X , it follows that x is orthogonal to z for all $x \in Z_1$, and hence $Z_1 \subset \Sigma$. We conclude that

$$\frac{\nu(\Sigma)}{m} \geq \frac{\nu(Z_1)}{m} = \frac{|\nu|}{n},$$

so that assumption (i) does not hold. \square

The proof of this last Proposition is a variation on the theme of [BMS94, Proposition 4.1].

PROPOSITION 4.8. — *Assume that assumption (ii) holds. Then for any ν -balanced product $q \in \mathcal{P}rod(\mathcal{V})$, the associated quantum channel \mathcal{E}_q as in Lemma 4.5 is invertible.*

Proof. — Let $q \in \mathcal{P}rod(\mathcal{V})$ be ν -balanced, and identify (\mathcal{V}, q) with $(\mathbb{C}^n, \langle \cdot, \cdot \rangle)$. For any $z \in \mathbb{C}^n \setminus \{0\}$, we denote by $[z]$ its class in $\mathbb{C}P^{n-1}$.

Denote by \tilde{Y} the cone of Y in \mathbb{C}^n . Assume on the contrary that an Hermitian matrix $A \neq 0$ lies in the kernel of \mathcal{E}_q , and set

$$F_A([z], [w]) := \frac{\langle Az, w \rangle}{|z| \cdot |w|}.$$

Since $\mathcal{E}_q = n^{-1}TT^*$ and $T^*(A)([z]) = F_A([z], [z])$ by the results of Section 2, we have $F_A([z], [z]) = 0$ for all $[z] \in Y$. Noticing that the function $(z, w) \mapsto \langle Az, w \rangle$ is holomorphic in z and anti-holomorphic in w and that it vanishes on the diagonal of $\tilde{Y} \times \tilde{Y}$, we conclude that F vanishes on $Z \times Z$ for every irreducible component Z of Y .

Pick any irreducible component Z . If it fully lies in $\text{Ker } A$, we have that Z is contained in a proper projective subspace. Otherwise, pick $[u] \in Z$ so that $Au \neq 0$. We thus proved that any other $[z] \in Z$ satisfies a linear equation $\langle z, Au \rangle = 0$,

meaning that Z lies in a proper projective subspace. This is in contradiction with assumption (ii). \square

Using the Propositions above, we are then ready to prove Theorem 4.4.

Proof of Theorem 4.4. — Suppose that ν satisfies the assumptions (i) and (ii), and fix $q_0 \in \mathcal{P}rod(\mathcal{V})$. By Proposition 4.6, the iterations $\mathcal{T}^r(q_0)$ converge to a fixed point $q \in \mathcal{P}rod(\mathcal{V})$ as $r \rightarrow +\infty$, so that we can use it to identify (\mathcal{V}, q) with $(\mathbb{C}^n, \langle \cdot, \cdot \rangle)$. Identify diffeomorphically $\mathcal{L}(\mathbb{C}^n)_+$ with $\mathcal{L}(\mathbb{C}^n)_+^1 \times \mathbb{R}_+$ via the map

$$\Theta : G \longmapsto (\mathcal{D}(G), \det(G)), \quad \text{where } \mathcal{D}(G) := \frac{G}{\det(G)^{\frac{1}{n}}}.$$

Then for every $r \in \mathbb{N}$,

$$(4.28) \quad \Theta \mathcal{T}_\nu^r \Theta^{-1}(G, g) = \left(\mathcal{D}(\mathcal{T}_\nu^r(G)), g \cdot \det \mathcal{T}_\nu^r(G) \right).$$

Recall that by Lemma 4.5, the differential of \mathcal{T}_ν at q coincides with the quantum channel \mathcal{E}_q , and recall that $q \in \mathcal{P}rod(\mathcal{V})$ is sent to the identity $\mathbb{1} \in \mathcal{L}(\mathbb{C}^n)_+$ in the identification of (\mathcal{V}, q) with $(\mathbb{C}^n, \langle \cdot, \cdot \rangle)$. Since $\mathcal{L}(\mathbb{C}^n)_+^1$ is a slice of the \mathbb{R}_+ -action and \mathcal{T}_ν is \mathbb{R}_+ -equivariant, the differential of $\mathcal{D} \circ \mathcal{T}_\nu$ equals to the restriction of \mathcal{E}_q to the tangent space $T_{\mathbb{1}}\mathcal{L}(\mathbb{C}^n)_+^1$, which consists of all trace 0 Hermitian matrices. Then by Propositions 4.7 and 4.8, the spectrum of this differential is contained in $(0, 1)$, so that $\mathcal{D} \circ \mathcal{T}_\nu$ is a local diffeomorphism of $\mathcal{L}(\mathbb{C}^n)_+^1$ in a neighborhood its hyperbolic fixed point $\mathbb{1}$, and conjugate through a local homeomorphism to its linearization at $\mathbb{1}$ by the classical Hartman–Grobman theorem. In particular, taking $\beta \in (0, 1)$ as the largest eigenvalue of \mathcal{E} in $(0, 1)$, we get a constant $C > 0$ such that

$$(4.29) \quad \text{dist}\left(\mathcal{D}(\mathcal{T}_\nu^r(G_0)), \mathbb{1}\right) \leq C\beta^r, \quad \text{for all } r \in \mathbb{N},$$

where $G_0 \in \mathcal{L}(\mathbb{C}^n)_+$ denotes the image of $q_0 \in \mathcal{P}rod(\mathcal{V})$ in the identification of (\mathcal{V}, q) with $(\mathbb{C}^n, \langle \cdot, \cdot \rangle)$. By (4.28), in order to complete the proof of the exponential convergence of the orbit of G_0 to $\mathbb{1}$, we need to show that for r large enough

$$(4.30) \quad \left| \det \mathcal{T}_\nu^r(G_0) - 1 \right| < C\beta^r.$$

To this end recall that the functional Ψ_ν of the proof of Proposition 4.6 is decreasing under iterations of \mathcal{T}_ν and invariant with respect to the action of \mathbb{R}_+ by multiplication. By (4.29) and the differentiability of Ψ_ν at $\mathbb{1}$, there exists a constant $C > 0$ such that

$$(4.31) \quad 0 \leq \Psi_\nu(\mathcal{T}_\nu^r(G_0)) - \Psi_\nu(\mathbb{1}) \leq C\beta^r.$$

Now as both (4.21) and (4.22) are non-positive and as $\mathcal{T}_\nu^r(G_0) \rightarrow \mathbb{1}$ as $r \rightarrow +\infty$, recalling the definition (4.11)-(4.12) of Ψ_ν we deduce in particular that

$$(4.32) \quad 0 \leq \log \det(\mathcal{T}_\nu^r(G_0)) \leq C\beta^r.$$

Since for x close to 1, we have $2|\log x| \geq |1 - x|$, this yields (4.30). The proof of Theorem 4.4 is complete. \square

Remark 4.9. — Consider the setting of Example 4.3 above for all $p \in \mathbb{N}^*$ large enough, where $q_p \in \mathcal{P}rod(H^0(X, L^p))$ is the unique ν -balanced product and h_p the induced Fubini–Study metric on L^p over X , with Chern curvature $-2i\pi\omega_p$. Recall that in that case, the induced POVM (4.2) coincides with the weighted Berezin–Toeplitz POVM (3.52) of Remark 3.12. Then using a refined version of Theorem 3.1 and Lemma 4.5, one can show that the exponential convergence rate $\beta_p \geq 0$ of Donaldson’s iterations in Theorem 4.4 satisfies as $p \rightarrow \infty$ the estimate

$$(4.33) \quad \beta_p = \frac{\lambda_1(\omega_\infty)}{4\pi p} + o(p^{-1}) ,$$

where $\lambda_1(\omega_\infty)$ is the first eigenvalue of the Laplace–Beltrami operator associated with the metric induced by the unique Kähler form ω_∞ in $c_1(L)$ solving the Calabi problem $\omega^d = c\nu$ for some $c > 0$. This follows from the estimate (3.54) on the spectral gap of the weighted Berezin transform, together with the uniformity on the metric in the estimates of [DLM06, Theorem 4.18’] and the fact that the sequence $\{\frac{1}{p}\omega_p\}$ converges to ω_∞ as $p \rightarrow \infty$. This complements a result of Keller in [Kel09, Proposition 4.7].

5. POVMs and geometry of measures

Assume that we are given an $\mathcal{L}(\mathcal{H})$ -valued POVM on Ω satisfying equation (2.1), i.e., of the form $dW = n F d\alpha$ for some $F : \Omega \rightarrow \mathcal{S}(\mathcal{H})$. In this section we discuss spectral properties of the Berezin transform associated to W in terms of the geometry of the measure

$$(5.1) \quad \sigma_W := F_*\alpha$$

on $\mathcal{S}(\mathcal{H})$, focusing on its multi-scale features, and on stability of the spectral gap under perturbations of the measure. Recall that for pure POVMs we have encountered measure (5.1) in Example 4.1.

Write $\mathcal{V} \subset \mathcal{L}(\mathcal{H})$ for the affine subspace consisting of all trace 1 operators, dist for the distance on \mathcal{V} associated to the scalar product $((A, B)) = \text{tr}(AB)$ on $\mathcal{L}(\mathcal{H})$. Given a compactly supported probability measure σ on \mathcal{V} , introduce the following objects:

- the center of mass $C(\sigma) = \int_{\mathcal{V}} v d\sigma(v)$;
- the mean squared distance from the origin,

$$I(\sigma) = \int_{\mathcal{V}} \text{dist}(C, v)^2 d\sigma(v) ;$$

- the mean squared distance to the best fitting line

$$J(\sigma) = \inf_{\ell} \int_{\mathcal{V}} \text{dist}(v, \ell)^2 d\sigma(v) ,$$

where the infimum is taken over all affine lines $\ell \subset \mathcal{V}$.

The infimum in the definition of J is attained at the (not necessarily unique) *best fitting line* which is known to pass through the center of mass C (Pearson, 1901; see [Far99, p. 188] for a historical account).⁽⁴⁾

Observe that the center of mass $C(\sigma_W)$ for the measure σ_W given by (5.1) coincides with the maximally mixed state $\frac{1}{n}\mathbb{1}$.

THEOREM 5.1. — *The spectral gap $\gamma(W)$ depends only on the push-forward measure σ_W on $\mathcal{S}(\mathcal{H})$:*

$$\gamma(W) = 1 - n(I(\sigma_W) - J(\sigma_W)) .$$

Proof. — Let $\ell \subset \mathcal{V}$ be any line passing through the center of mass $\frac{1}{n}\mathbb{1}$ generated by a trace zero unit vector $A \in \mathcal{L}(\mathcal{H})$. For a point $B \in \mathcal{V}$ we have

$$\text{dist}(B, \ell)^2 = \left(\left(B - \frac{1}{n}\mathbb{1}, B - \frac{1}{n}\mathbb{1} \right) \right) - \left(\left(B - \frac{1}{n}\mathbb{1}, A \right) \right)^2 .$$

Integrating over σ_W and taking infimum over ℓ we get that

$$(5.2) \quad J(\sigma_W) = I(\sigma_W) - K ,$$

with

$$(5.3) \quad K = \sup_{\substack{\text{tr}(A)=0 \\ \text{tr}(A^2)=1}} \int_{\mathcal{V}} ((B, A))^2 dF_*\alpha(B) .$$

The latter integral can be rewritten as

$$(5.4) \quad \int_{\Omega} ((F(s), A))^2 d\alpha(s) = n^{-1}((\mathcal{E}(A), A)) ,$$

so by definition $K = n^{-1}\gamma_1 = n^{-1}(1 - \gamma(W))$. Substituting this into (5.2), we deduce the Theorem 5.1. \square

Remark 5.2. — Observe that the supremum in (5.3) is attained at a unit vector A generating the best fitting line. By (5.4), A is an eigenvector of \mathcal{E} with the eigenvalue γ_1 .

Example 5.3. — For a pure POVM W , i.e. when F is a one-to-one map from Ω to the set of rank-one projectors,

$$\text{dist}(C, F(s))^2 = \text{tr} \left[\left(\frac{1}{n}\mathbb{1} - F(s) \right)^2 \right] = 1 - \frac{1}{n}$$

for all $s \in \Omega$, and hence $I(\sigma_W) = 1 - \frac{1}{n}$. Thus, by Theorem 5.1,

$$(5.5) \quad J(\sigma_W) = \frac{n - 2 + \gamma}{n} .$$

For instance, consider the (pure!) Berezin–Toeplitz POVM W_p from Example 4.2. Let us use formula (5.5) in order to calculate J . Recall that by the Riemann–Roch theorem (see [Fin12a, Propositions 2.25 and 4.21])

$$n_p = Vp^d + Up^{d-1} + \mathcal{O}(p^{d-2}) ,$$

⁽⁴⁾The problem of finding J and the corresponding minimizer ℓ appears in the literature under several different names including “total least squares” and “orthogonal regression”.

where

$$V = \text{Vol}(X) = \frac{[\omega]^d}{d!}, \quad U = c_1(X) \cup \frac{[\omega]^{d-1}}{(d-1)!}.$$

It follows from formula (3.11) for γ_p that

$$J(\sigma_{W_p}) = 1 - \frac{2}{V}p^{-d} + \frac{8\pi U + V\lambda_1}{4\pi V^2}p^{-d-1} + \mathcal{O}(p^{-d-2}).$$

For instance, for the dual to the tautological bundle over $\mathbb{C}P^1$ in Example 3.4 $n = p + 1$ and $\gamma = \frac{2}{p+2}$ so by (5.5) $J = 1 - \frac{2}{p+2}$.

Furthermore, we explore robustness of the gap $\gamma(W)$, as a function of the measure σ_W , with respect to perturbations in the Wasserstein distances on the space of Borel probability measures on $\mathcal{S}(\mathcal{H})$. They are defined as follows. For compactly supported Borel probability measures σ_1, σ_2 on a metric space (X, d) the L_2 -Wasserstein distance is given by

$$\delta_2(\sigma_1, \sigma_2) := \inf_{\nu} \left(\int_{X \times X} \text{dist}(x_1, x_2)^2 d\nu(x_1, x_2) \right)^{\frac{1}{2}},$$

and the L_∞ -Wasserstein distance by

$$\delta_\infty(\sigma_1, \sigma_2) := \inf_{\nu} \sup_{(x_1, x_2) \in \text{supp}(\nu)} \text{dist}(x_1, x_2),$$

where in both cases the infimum is taken over all Borel probability measures ν on $X \times X$ with marginals σ_1 and σ_2 .

THEOREM 5.4. — *Let σ_V and σ_W be measures on $\mathcal{S}(\mathcal{H})$ associated to POVMs V and W respectively.*

- (i) $|\gamma(V) - \gamma(W)| \leq c(n)\delta_2(\sigma_V, \sigma_W)$, where $c(n)$ depends on the dimension $n = \dim \mathcal{H}$;
- (ii) *If in addition V and W are pure POVMs, there exists a universal constant c such that*

$$(5.6) \quad |\gamma(V) - \gamma(W)| \leq c\delta_\infty(\sigma_V, \sigma_W).$$

Note that this result enables us to compare spectral gaps of POVMs defined on different sets (but having values in the same Hilbert space). This idea goes back to [OC09]⁽⁵⁾. Let us emphasize that the estimate in (ii) is *dimension-free*. This is important, for instance, for comparison of spectral gaps corresponding to different Berezin–Toeplitz quantization schemes.

Theorem 5.4(i) immediately follows from the fact that $C(\sigma)$, $I(\sigma)$ and $J(\sigma)$ are Lipschitz in σ with respect to L_2 -Wasserstein distance. The details will appear in MSc thesis by V. Kaminker.

For the proof of part (ii), we need the following auxiliary statement. In what follows we write $\|A\|_2$ for the Hilbert–Schmidt norm $(\text{tr}(AA^*))^{\frac{1}{2}}$.

⁽⁵⁾In [OC09] the authors consider the L_1 -version of this distance, and call it the Kantorovich distance.

LEMMA 5.5. — *Let P, Q be rank 1 orthogonal projectors. Then for every $A \in \mathcal{L}(\mathcal{H})$,*

$$|\operatorname{tr}(A(P - Q))| \leq \sqrt{2} \|P - Q\|_2 \left(\operatorname{tr}(A^2(P + Q)) \right)^{\frac{1}{2}}.$$

Proof. — Suppose that P and Q are orthogonal projectors to unit vectors ξ and η , respectively. By tuning the phase of ξ , we can assume that $\langle \xi, \eta \rangle \geq 0$. We have

$$\begin{aligned} |\operatorname{tr}(A(P - Q))| &= |\langle A\xi, \xi \rangle - \langle A\eta, \eta \rangle| \\ &= |\langle \xi - \eta, A\xi \rangle + \langle A\eta, \xi - \eta \rangle| \leq |\xi - \eta| (|A\xi| + |A\eta|) \\ &= |\xi - \eta| \left(\left\langle A^2\xi, \xi \right\rangle^{\frac{1}{2}} + \left\langle A^2\eta, \eta \right\rangle^{\frac{1}{2}} \right) \leq \sqrt{2} |\xi - \eta| \left(\left\langle A^2\xi, \xi \right\rangle + \left\langle A^2\eta, \eta \right\rangle \right)^{\frac{1}{2}} \\ &= \sqrt{2} |\xi - \eta| \left(\operatorname{tr}(A^2P) + \operatorname{tr}(A^2Q) \right)^{\frac{1}{2}}. \end{aligned}$$

But since $0 \leq \langle \xi, \eta \rangle \leq 1$,

$$\begin{aligned} |\xi - \eta| &= (2 - 2\langle \xi, \eta \rangle)^{\frac{1}{2}} \leq (2 - 2\langle \xi, \eta \rangle^2)^{\frac{1}{2}} \\ &= \left(\operatorname{tr}(P - Q)^2 \right)^{\frac{1}{2}} = \|P - Q\|_2. \end{aligned}$$

This completes the proof of Lemma 5.5. \square

Proof. — Proof of Theorem 5.4 (ii): Denote by \mathcal{P} the space of all rank 1 orthogonal projectors on \mathcal{H} . We can assume without loss of generality that pure POVMs V and W are defined on subsets Ω_1 and Ω_2 of \mathcal{P} , respectively, and that the maps $F_i : \Omega_i \rightarrow \mathcal{P}$ are the inclusions. Thus representation (2.1) in this case can be simplified as

$$dV(s) = n s d\alpha_1(s), \quad dW(t) = n t d\alpha_2(t),$$

where $\sigma_V = \alpha_1$ and $\sigma_W = \alpha_2$ are Borel probability measures supported in Ω_1 and Ω_2 , respectively. Let us emphasize that here and below s, t stand for rank 1 orthogonal projectors. Pick any measure ν on $\mathcal{P} \times \mathcal{P}$ with marginals α_1 and α_2 and write

$$\Delta := \max_{(s,t) \in \operatorname{supp}(\nu)} \|s - t\|_2.$$

We use the fact that the operators $\mathcal{E}_1, \mathcal{E}_2 : \mathcal{L}(\mathcal{H}) \rightarrow \mathcal{L}(\mathcal{H})$ given by formula (2.2) have the same spectrum as the Berezin transform. For $A \in \mathcal{L}(H)$ with $\operatorname{tr}(A^2) = 1$ put

$$D := \left| ((\mathcal{E}_1 A, A)) - ((\mathcal{E}_2 A, A)) \right|.$$

One readily rewrites

$$\begin{aligned} D &= n \left| \int_{\Omega_1} ((F_1(s), A))^2 d\alpha_1(s) - \int_{\Omega_2} ((F_2(t), A))^2 d\alpha_2(t) \right| \\ &\leq n \int_{\Omega_1 \times \Omega_2} \left| \operatorname{tr}((s - t)A) \right| \left| \operatorname{tr}((s + t)A) \right| d\nu. \end{aligned}$$

By Lemma 5.5,

$$\left| \operatorname{tr}((s - t)A) \right| \leq \sqrt{2} \|s - t\|_2 \left(\operatorname{tr}(A^2(s + t)) \right)^{\frac{1}{2}}.$$

By Cauchy–Schwarz, writing

$$(s+t)A = (s+t)^{\frac{1}{2}} \left((s+t)^{\frac{1}{2}} A \right) ,$$

we get

$$\left| \operatorname{tr}((s+t)A) \right| \leq (\operatorname{tr}(s+t))^{\frac{1}{2}} \left(\operatorname{tr}(A^2(s+t)) \right)^{\frac{1}{2}} = \sqrt{2} \left(\operatorname{tr}(A^2(s+t)) \right)^{\frac{1}{2}} .$$

It follows that

$$D \leq 2n \max_{(s,t) \in \operatorname{supp}(\nu)} \|s-t\|_2 \int \operatorname{tr} \left(A^2(s+t) \right) d\nu .$$

The integral on the right can be rewritten as

$$\operatorname{tr} \left(A^2 \int_{\Omega_1} s d\alpha_1(s) \right) + \operatorname{tr} \left(A^2 \int_{\Omega_2} t d\alpha_2(t) \right) = \frac{2}{n} ,$$

since

$$\int_{\Omega_1} s d\alpha_1(s) = \int_{\Omega_2} t d\alpha_2(t) = \frac{1}{n} \mathbb{1}$$

and $\operatorname{tr}(A^2) = 1$. It follows that $D \leq 4\Delta$. Choosing ν so that Δ becomes arbitrary close to $\delta := \delta_\infty(\alpha_1, \alpha_2)$, and taking A with

$$(5.7) \quad \operatorname{tr}(A) = 0 , \quad \operatorname{tr}(A^2) = 1$$

to be an eigenvector of \mathcal{E}_1 with the first eigenvalue $\gamma_1(\mathcal{E}_1)$, we get that

$$|\gamma_1(\mathcal{E}_1) - ((\mathcal{E}_2 A, A))| \leq 4\delta .$$

But due to the variational characterization of the first eigenvalue, $\gamma_1(\mathcal{E}_2) = \max((\mathcal{E}_2 A, A))$, where the maximum is taken over all A satisfying (5.7). It follows that $\gamma_1(\mathcal{E}_1) - \gamma_1(\mathcal{E}_2) \leq 4\delta$. By symmetry, $\gamma_1(\mathcal{E}_2) - \gamma_1(\mathcal{E}_1) \leq 4\delta$, which yields the theorem with $c = 4$. \square

Our next result provides a geometric characterization of the eigenfunction of the operator \mathcal{B} with the eigenvalue γ_1 . Let $A \in \mathcal{L}(\mathcal{H})$ be the trace zero unit vector generating the best fitting line corresponding to W . In view of Theorem 5.1,

$$\gamma_1 = 1 - \gamma(W) = n(I - J) ,$$

with $I = I(\sigma_W)$ and $J = J(\sigma_W)$.

THEOREM 5.6. — *The function*

$$(5.8) \quad \psi_1 : \Omega \rightarrow \mathbb{R}, \quad s \mapsto \frac{((F(s), A))}{\sqrt{I - J}}$$

is an eigenfunction of the operator \mathcal{B} with the eigenvalue γ_1 . Furthermore, $\|\psi_1\| = 1$.

In other words, up to a multiplicative constant, the first eigenfunction sends $s \in \Omega$ to the projection of the density $F(s)$ to the best fitting line.

Proof. — By Remark 5.2 above, the operator A generating the best fitting line is an eigenvector of the quantum channel \mathcal{E} : $\mathcal{E}A = \gamma_1 A$. Since $\mathcal{E} = n^{-1}TT^*$ and $\mathcal{B} = n^{-1}T^*T$, we have $\mathcal{B}(T^*A) = \gamma_1 T^*A$ and $(T^*A, T^*A) = n\gamma_1$. Furthermore, $T^*A(s) = n((F(s), A))$ and $n\gamma_1 = n^2(I - J)$. Choosing $\psi_1 = \frac{T^*A}{\|T^*A\|}$, we get (5.8). \square

Next, we discuss the *diffusion distance* on Ω associated to the Markov operator \mathcal{B} (see [CL06]). This distance, which originated in geometric analysis of data sets, depends on a positive parameter τ playing the role of the time in the corresponding random process. Take any orthonormal eigenbasis $\{\psi_k\}$ corresponding to eigenvalues $1 = \gamma_0 \geq \gamma_1 \geq \gamma_2 \dots$ of \mathcal{B} such that ψ_0 is constant. The diffusion distance D_τ is defined by

$$(5.9) \quad D_\tau(s, t) = \left(\sum_{k \geq 1} \gamma_k^{2\tau} (\psi_k(s) - \psi_k(t))^2 \right)^{\frac{1}{2}} \quad \forall s, t \in \Omega.$$

If $\gamma_1 < 1$, i.e., the spectral gap is positive, this expression decays exponentially.

Suppose now that $\gamma_2 < \gamma_1$. In this case the asymptotic behavior of $D_\tau(s, t)$ as $\tau \rightarrow \infty$ is given by

$$(5.10) \quad D_\tau(s, t) = \gamma_1^\tau \frac{|((F(s) - F(t), A))|}{(I - J)^{\frac{1}{2}}} (1 + o(1)), \text{ if } ((F(s), A)) \neq ((F(t), A)),$$

and $D_\tau(s, t) = \mathcal{O}(\gamma_2^\tau)$ otherwise. The difference in these asymptotic formulas highlights the multi-scale behaviour of the metric space (Ω, D_τ) . In the first approximation, this space consists of the level sets of the function $s \mapsto ((F(s), A))$ situated at the distance $\sim \gamma_1^\tau$ from one another, while each fiber has the diameter $\lesssim \gamma_2^\tau$. Viewing POVMs as data clouds in \mathcal{S} opens up a prospect of using various tools of geometric data analysis for studying POVMs. The above result on the diffusion distance associated to a POVM can be considered as a step in this direction.

6. Case study: representations of finite groups

In this section we will be interested in finite POVMs associated to irreducible representations of finite groups. We start with some preliminaries from Woldron's book [Wal18]. Let G be a finite set.

DEFINITION 6.1. — *A finite collection $\{f_s\}_{s \in G}$ of non-zero vectors in a finite-dimensional Hilbert space \mathcal{H} is said to be a tight frame if there exists a number $A > 0$, called the frame bound, such that*

$$(6.1) \quad A\|f\|^2 = \sum_{s \in G} |\langle f, f_s \rangle|^2, \forall f \in \mathcal{H}.$$

Denote by P_s the orthogonal projector to f_s . One readily checks that for such a frame, the operators

$$(6.2) \quad W_s := \frac{\|f_s\|^2}{A} P_s, \quad s \in G,$$

form a $\mathcal{L}(\mathcal{H})$ -valued POVM on G .

Suppose from now on that G is a finite group, and we are given its non-trivial irreducible unitary representation ρ on a d_ρ -dimensional Hilbert space V .⁽⁶⁾ One can show [Wal18] that the vectors $\{f_s := \frac{1}{\sqrt{d_\rho}} \rho(s)\}_{s \in G}$ form a tight frame in the operator

⁽⁶⁾ All the representations considered below are assumed to be unitary.

space $\mathcal{H} := \text{End}(V)$ equipped with the Hermitian product $((C, D)) = \text{tr}(CD^*)$ with the frame bound $A = |G|/d_\rho^2$. Write $n = d_\rho^2 = \dim \mathcal{H}$. By (6.2), the corresponding POVM $W = \{W_s\}$, $s \in G$ is given by $W_s = nP_s\alpha_s$ with $\alpha_s = \frac{1}{|G|}$. Interestingly enough, the spectrum of the corresponding Berezin transform can be calculated via the characters of irreducible unitary representations of G .

Denote by $\chi_\rho : G \rightarrow \mathbb{C}$, $\chi_\rho(s) := \text{tr}(\rho(s))$ the character of the representation ρ . Consider a basis in $L_2(G)$ consisting of the indicator functions of the elements of G . It readily follows from the definition that the Berezin transform \mathcal{B} corresponding to the POVM W is given by a matrix

$$\mathcal{B}_{ts} = n \text{tr}(P_t P_s) \alpha_s = \frac{1}{|G|} u(st^{-1}) ,$$

where $u(s) := |\chi_\rho(s)|^2$. The eigenvalues of this matrix and their multiplicities are given by the following proposition, see [Dia88, Chapter 3E].

PROPOSITION 6.2. — *The eigenvalues of \mathcal{B} are given by*

$$\lambda_\varphi := \frac{1}{d_\varphi |G|} \sum_{s \in G} u(s) \chi_\varphi(s) ,$$

where φ runs over irreducible representations of G , and the contribution of each φ into the multiplicity of λ_φ is d_φ^2 .

Let us emphasize that it could happen that $\lambda_\varphi = \lambda_\psi$ for different representations φ and ψ . Note also that by Lemma 6.5(i) below, $\lambda_\varphi = 1$ when φ is the trivial one-dimensional representation.

Remark 6.3. — We claim that the gap $\gamma(W)$ is rational. Indeed, for a unitary representation ψ by complex unitary matrices denote $\psi'(s) = \overline{\psi(s)}$, where the bar stands for the complex conjugation. Note that u is the character of the (in general, reducible) representation $\theta := \rho \otimes \rho'$. By the Schur orthogonality relations,

$$\frac{1}{|G|} \sum_{s \in G} u(s) \chi_\varphi(s)$$

equals the multiplicity of φ in the decomposition of θ into irreducible representations, and hence is an integer. The claim follows from Proposition 6.2.

The main result of this section is the following algebraic criterion of the positivity of the spectral gap of W . Following [Isa06, Chapter 12] we define the *vanishing-off subgroup* $\mathcal{V}(\rho)$ to be the smallest subgroup of G such that χ_ρ vanishes on $G \setminus \mathcal{V}(\rho)$:

$$\mathcal{V}(\rho) = \langle s \in G \mid \chi_\rho(s) \neq 0 \rangle .$$

Since the character χ_ρ is conjugation invariant, $\mathcal{V}(\rho)$ is normal.

THEOREM 6.4. — *The following are equivalent:*

- (i) $\mathcal{V}(\rho) \neq G$;
- (ii) $\gamma(W) = 0$, i.e., there exists a non-trivial irreducible unitary representation φ of G with $\lambda_\varphi = 1$.

In the next Lemma 6.5, we collect some standard facts from the representation theory (see e.g. [Dia88, Chapter 2]) which will be used in the proof of Theorem 6.4. We write Irrep for the set of all unitary irreducible representations of G up to an isomorphism.

LEMMA 6.5. —

- (i)
$$\frac{1}{|G|} \sum_{s \in G} |\chi_\rho(s)|^2 = 1, \quad \forall \rho \in \text{Irrep};$$
- (ii)
$$\sum_{\varphi \in \text{Irrep}} d_\varphi^2 = |G|.$$

Proof. — *Proof of Theorem 6.4:* We begin by proving (ii) \Rightarrow (i): Assume that there exists a non-trivial irreducible representation φ with $\lambda_\varphi = 1$. By using Lemma 6.5 and the explicit formula for the eigenvalues from Proposition 6.2 we see that

$$\begin{aligned} 1 &\stackrel{6.2}{=} \frac{1}{d_\varphi |G|} \sum_{s \in G} |\chi_\rho(s)|^2 \chi_\varphi(s) = \\ &= \frac{1}{|G|} \sum_{s \in G} |\chi_\rho(s)|^2 - \frac{1}{d_\varphi |G|} \sum_{s \in G} |\chi_\rho(s)|^2 (d_\varphi - \chi_\varphi(s)) \stackrel{6.5}{=} \\ &= 1 - \frac{1}{d_\varphi |G|} \sum_{s \in G} |\chi_\rho(s)|^2 (d_\varphi - \chi_\varphi(s)) \end{aligned}$$

By taking the real part of both sides we get

$$\sum_{s \in G} |\chi_\rho(s)|^2 \text{Re}(d_\varphi - \chi_\varphi(s)) = 0$$

Note that since $\varphi(s)$ is unitary, all its eigenvalues are of the form $e^{i\theta}$ so $|\chi_\varphi(s)| = |\text{tr}(\varphi(s))| \leq d_\varphi$ and $\chi_\varphi(s) = d_\varphi$ iff $\varphi(s)$ is the identity. Hence $\text{Re}(d_\varphi - \chi_\varphi(s))$ must be non-negative. Since $|\chi_\rho(s)|^2$ is also non-negative, $\chi_\varphi(s) = d_\varphi$ for every $s \in G$ with $\chi_\rho(s) \neq 0$. As we have seen above $\chi_\varphi(s) = d_\varphi$ if and only if $\varphi(s) = \mathbb{1}$. It follows that the vanishing off subgroup $\mathcal{V}(\rho)$ is contained in the normal subgroup

$$\text{Ker}(\varphi) := \{s \mid \varphi(s) = \mathbb{1}\}.$$

Since φ is irreducible and non-trivial, the latter subgroup $\neq G$, and hence $\mathcal{V}(\rho) \neq G$, as required.

Next, we prove (i) \Rightarrow (ii): Assume $\mathcal{V}(\rho) \neq G$. Consider the quotient $H := G/\mathcal{V}(\rho)$, which is a non-trivial group, and let $\pi : G \rightarrow H$ be the natural projection. Take any non-trivial irreducible representation ψ of H . Then $\varphi := \psi \circ \pi$ is an irreducible representation of G . We claim that $\lambda_\varphi = 1$. Indeed, for $s \in \mathcal{V}(\rho)$ we have $\varphi(s) = \mathbb{1}$ and hence $\chi_\varphi(s) = d_\varphi$, and for $s \notin \mathcal{V}(\rho)$ holds $\chi_\rho(s) = 0$. It follows that

$$\lambda_\varphi = \sum_{s \in \mathcal{V}(\rho)} \frac{1}{d_\varphi |G|} |\chi_\rho(s)|^2 d_\varphi = \frac{1}{|G|} \sum_{s \in G} |\chi_\rho(s)|^2 \stackrel{6.5}{=} 1.$$

This proves the claim and hence completes the proof of the Theorem 6.4. \square

COROLLARY 6.6. — *If G is a simple group, then the gap of W is positive.*

Proof. — Indeed, otherwise by Theorem 6.4 and the simplicity of G , $\mathcal{V}(\rho) = \{1\}$, which means that $\chi_\rho(s) = 0$ for every $s \neq 1$. Then the first statement of Lemma 6.5 yields $|G| = d_\rho^2$, while the second statement guarantees that $|G| \geq 1 + d_\rho^2$, since ρ is a non-trivial representation. We get a contradiction. \square

Let us point out that there exist non-simple groups G admitting an irreducible representation ρ with $\mathcal{V}(\rho) = G$. Indeed, consider the irreducible representation $\rho : \mathbb{Z}_m \rightarrow U(\mathbb{C})$, $\rho(s) = e^{2\pi i s/m}$ of the abelian cyclic group \mathbb{Z}_m . Observe that $\mathcal{V}(\rho) = \mathbb{Z}_m$, while \mathbb{Z}_m is simple if and only if m is prime.

Let us describe the diffusion distance D_τ (see (5.9)) corresponding to the POVM W associated to a finite group G and a non-trivial irreducible representation ρ . Recall [Dia88] that for an irreducible representation $\varphi : G \rightarrow \mathbb{U}(n)$, the orthonormal basis of eigenfunctions corresponding to the eigenvalue λ_φ presented in Proposition 6.2 is given by the matrix coefficients of φ multiplied by $\sqrt{d_\varphi}$. Assume that the gap of G is strictly positive, and denote by $\beta_1 > \dots > \beta_k$ all *pair-wise distinct* eigenvalues of \mathcal{B} lying in the open interval $(0, 1)$. Denote

$$R_j := \{\varphi \in \text{Irrep} \mid \lambda_\varphi = \beta_j\}.$$

Then (5.9) yields the following expression for the diffusion distance:

$$(6.3) \quad D_\tau(s, t) = \left(\sum_{j=1}^k \beta_j^{2\tau} \sum_{\varphi \in R_j} d_\varphi \|\varphi(s) - \varphi(t)\|_2^2 \right)^{\frac{1}{2}},$$

where $\|\cdot\|_2$ stands for the Hilbert–Schmidt norm $\|C\|_2 = (\text{tr}(CC^*))^{\frac{1}{2}}$. Note that this expression can be rewritten in terms of the character χ_φ since

$$\|\varphi(s) - \varphi(t)\|_2^2 = 2 \left(d_\varphi - \text{Re } \chi_\varphi(st^{-1}) \right).$$

Define a normal subgroup $\Gamma_j := \bigcap_{\varphi \in R_j} \text{Ker}(\varphi)$, $j = 1, \dots, k$ and a normal series $K_0 \supset K_1 \supset \dots$ with $K_0 = G$, $K_{k+1} = \{1\}$ and

$$K_m := \bigcap_{j=1}^m \Gamma_j, \quad m = 1, \dots, k.$$

It follows from (6.3) that for $\tau \rightarrow +\infty$

$$(6.4) \quad D_\tau(s, t) \sim \beta_{p+1}^\tau \quad \text{for } st^{-1} \in K_p \setminus K_{p+1}.$$

In fact we have a sequence of nested partitions Δ_p of G formed by the cosets of K_p . For every pair of distinct points $s, t \in G$ choose maximal p so that s and t lie in the same element of Δ_p . Then asymptotical formula (6.4) holds, which manifests the multi-scale nature of the diffusion distance.

Let us illustrate this in the case when $G = S_4$ is the symmetric group, and ρ a 3-dimensional irreducible representation. The direct calculation with the character table of S_4 shows that the first non-trivial eigenvalue $\frac{1}{2}$ corresponds to the unique 2-dimensional irreducible representation whose kernel coincides with the normal subgroup K of order 4 of S_4 called the Klein four-group. Thus $D_\tau(s, t) \sim (\frac{1}{2})^\tau$ if s, t

belong to different cosets of K in S_4 , and one can calculate that $D_\tau(s, t) \sim (\frac{1}{3})^\tau$ if s, t are distinct and belong to the same coset.

Remark 6.7. — A modification of the construction presented in this section is related to Berezin–Toeplitz quantization. The modification goes in two directions. First, we deal with unitary representations ρ of compact Lie groups G instead of finite groups, and second, our POVMs are related to the G –orbits in a representation space \mathcal{H} as opposed to the image of ρ in the endomorphisms of \mathcal{H} . Let us very briefly illustrate this in the following simplest case. Consider the irreducible unitary representation ρ_j of the group $G = SU(2)$ in an $n = 2j + 1$ -dimensional Hilbert space \mathcal{H} , $j \in \frac{1}{2}\mathbb{N}$. Fix a maximal torus $K = S^1 \subset G$, and let $w \in \mathcal{H}$ be the maximal weight vector of K , that is $\rho_j(t)w = e^{4\pi i j t}w$ for all $t \in K$. Consider an $\mathcal{L}(\mathcal{H})$ -valued POVM W on $\Omega = G/K = \mathbb{C}P^1$ of the form $dW([g]) = nP_{[g]}d\alpha([g])$, where $[g]$ stands for the class of $g \in G$ in Ω , α is the G -invariant measure on Ω and $P_{[g]}$ is the rank one projector to gw . Note that W is nothing else but the Berezin–Toeplitz POVM W_p from Example 3.4 with $p = 2j$. We refer to [CR12, Chapter 7] for the representation theoretic approach to coherent states and quantization. By using theory of Gelfand pairs (cf. [Dia88, Chapter 3.F]) one can check that the eigenvalues of the Berezin transform are of the form $\lambda_\varphi = (u, \chi_\varphi)_{L_2}$, where φ runs over all irreducible unitary representations of G , χ_φ stands for the character of φ and $u(g) = n|\langle \rho(g)w, w \rangle|^2$. The multiplicity of λ_φ equals d_φ , where d_φ is the dimension of φ . In order to calculate λ_φ , recall that

$$(6.5) \quad \rho_j \otimes \rho_j = \bigoplus_{k=0}^{2j} \rho_k.$$

Writing v for the vector of weight $-j$ of ρ_j , we have

$$u(g) = n \langle (\rho_j \otimes \rho_j)(g)\xi, \xi \rangle, \text{ where } \xi = w \otimes v.$$

In order to complete this calculation, one has to decompose ξ in the sense of (6.5). This can be done with the help of explicit expressions for the Clebsch–Gordan coefficients, and it eventually yields eigenvalues of the Berezin transform, including $\gamma_1 = \frac{j}{j+1}$ (cf. Example 3.4), in agreement with calculations by Zhang [Zha98] and Donaldson [Don09, p. 613]. The details will appear in MSc thesis by D. Shmoish.

7. Two concepts of quantum noise

In the present section we provide two different (and essentially tautological) interpretations of the spectral gap in the context of quantum noise. In quantum measurement theory, there are two concepts of quantum noise: the increment of variance for unbiased approximate measurements as formalized by the noise operator, see below, and a non-unitary evolution of a quantum system described by a quantum channel (a.k.a. a quantum operation, see, e.g. [NC00, Chapter 8]). Such a non-unitary evolution can be caused, for instance, by the quantum state reduction in the process of repeated quantum measurements. Interestingly enough, for pure POVMs, the

spectral gap $\gamma(W)$ brings together these two seemingly remote concepts: it measures the minimal magnitude of noise production in the context of the noise operator, and it equals the spectral gap of the Markov chain modeling repeated quantum measurements.

Given an observable $A \in \mathcal{L}(\mathcal{H})$, write $A = \sum \lambda_i P_i$ for its spectral decomposition, where P_i 's are pair-wise distinct orthogonal projectors. According to the statistical postulate of quantum mechanics, in a state ρ the observable A attains value λ_i with probability $((P_i, \rho))$. It follows that the expectation of A in ρ equals $\mathbb{E}(A, \rho) = ((A, \rho))$ and the variance is given by $\mathbb{V}ar(A, \rho) = ((A^2, \rho)) - \mathbb{E}(A, \rho)^2$. In quantum measurement theory [BLPY16], a POVM W represents a measuring device coupled with the system, while Ω is interpreted as the space of device readings. When the system is in a state $\rho \in \mathcal{S}(\mathcal{H})$, the probability of finding the device in a subset $X \in \mathcal{C}$ equals $\mu_\rho(X) := ((W(X), \rho))$. An experimentalist performs a measurement whose outcome, at every state ρ , is distributed in Ω according to the measure μ_ρ . Given a function $\phi \in L_2(\Omega, \alpha)$ (experimentalist's choice), this procedure yields an *unbiased approximate measurement* of the quantum observable $A := T(\phi)$. The expectation of A in every state ρ equals $((A, \rho))$ and thus coincides with the one of the measurement procedure given by $\int_\Omega \phi d\mu_\rho$ (hence *unbiased*), in spite of the fact that actual probability distributions determined by the observable A (see above) and the random variable (ϕ, μ_ρ) could be quite different (hence *approximate*). In particular, in general, the variance increases under an unbiased approximate measurement:

$$(7.1) \quad \mathbb{V}ar(\phi, \mu_\rho) = \mathbb{V}ar(A, \rho) + ((\Delta_W(\phi), \rho)) ,$$

where $\Delta_W(\phi) := T(\phi^2) - T(\phi)^2$ is the *noise operator*. This operator, which is known to be positive, measures the increment of the variance. We wish to explore the relative magnitude of this increment for the “maximally mixed” state $\theta_0 = \frac{1}{n} \mathbb{1}$. To this end introduce the *minimal noise* of the POVM W as

$$\mathcal{N}_{min}(W) := \inf_{\phi} \frac{((\Delta_W(\phi), \theta_0))}{\mathbb{V}ar(\phi, \mu_{\theta_0})} ,$$

where the infimum is taken over all non-constant functions $\phi \in L_2(\Omega, \alpha)$. It turns out that the minimal noise coincides with the spectral gap:

$$(7.2) \quad \mathcal{N}_{min}(W) = \gamma(W) .$$

Indeed, since $\text{tr}(T(\phi^2)) = n(\phi, \phi)$, we readily get that

$$((\Delta_W(\phi), \theta_0)) = ((\mathbb{1} - \mathcal{B})\phi, \phi) ,$$

where $\mathcal{B} = n^{-1}T^*T$ is the Markov operator given by (2.3), while

$$\mathbb{V}ar(\phi, \mu_{\theta_0}) = (\phi, \phi) - (\phi, 1)^2 .$$

Formula (7.2) follows from the variational principle.

Suppose now that $\Omega \subset \mathcal{S}(\mathcal{H})$ is a finite set consisting of rank one projectors $\{P_1, \dots, P_N\}$ and that W is a pure POVM of the form $W(P_i) := n\alpha_i P_i$, where α is a probability measure on Ω . Given a system in the original state ρ , the result of the measurement equals P_j with probability $p = n\alpha_j((P_j, \rho))$. Recall the quantum state reduction (a.k.a. the wave function collapse) axiom for so called *Lüders* repeated

quantum measurements: if the result of the measurement equals P_j , the system moves from the original state ρ to the new (reduced) state

$$\rho' = \frac{1}{p} W(P_j)^{\frac{1}{2}} \rho W(P_j)^{\frac{1}{2}} = P_j .$$

It follows that if the original state ρ is chosen from Ω , the repeated quantum measurements are described by the Markov chain with transition probabilities $n\alpha_j((P_i, P_j))$. The corresponding Markov operator equals \mathcal{B} , and *the spectral gap of the Markov chain coincides with the spectral gap $\gamma(W)$ of the POVM W* . Furthermore, given an original state $\rho \in \Omega$, the expected value of the reduced state equals $\mathcal{E}(\rho)$. It follows that if $\gamma(W) > 0$, $\mathcal{E}^k(\rho)$, $k \rightarrow \infty$ converge to the maximally mixed quantum state $\frac{1}{n}\mathbb{1}$ at the exponential rate $\sim (1 - \gamma(W))^k$. In other words, *for pure POVMs the spectral gap controls the convergence rate to the maximally mixed state under repeated quantum measurements*.

Acknowledgments

We are grateful to D. Aharonov, S. Artstein-Avidan, J.-M. Bismut, L. Charles, G. Kalai, G. Kindler, M. Krivelevich, S. Nonnenmacher, Y. Ostrover, A. Reznikov, A. Veselov and S. Weinberger for useful discussions on various aspects of this work. We thank O. Oreshkov for bringing our attention to the paper [OC09] and illuminating comments, as well as I. Polterovich for very helpful remarks. We thank A. Deleporte for sharing with us his ideas on the proof of Theorem 3.1, and J. Fine for providing us highly useful insights and references on Donaldson's program. Finally, we thank the anonymous referee for numerous useful comments.

BIBLIOGRAPHY

- [AS17] Guillaume Aubrun and Stanislaw J. Szarek, *Alice and Bob meet Banach. The interface of asymptotic geometric analysis and quantum information theory*, Mathematical Surveys and Monographs, vol. 223, American Mathematical Society, 2017. ↑1346
- [Ber75] Feliks Alexandrovitch Berezin, *Quantization*, Math. USSR, Izv. **8** (1975), no. 5, 1116–1175. ↑1352
- [BGL13] Dominique Bakry, Ivan Gentil, and Michel Ledoux, *Analysis and geometry of Markov diffusion operators*, Grundlehren der Mathematischen Wissenschaften, vol. 348, Springer, 2013. ↑1348
- [BGV92] Nicole Berline, Evra Getzler, and Michèle Vergne, *Heat kernels and Dirac operators*, Grundlehren der Mathematischen Wissenschaften, vol. 298, Springer, 1992. ↑1354, 1358
- [BLPY16] Paul Busch, Pekka J. Lahti, Juha-Pekka Pellonpää, and Kari Ylinen, *Quantum measurement*, Theoretical and Mathematical Physics, Springer, 2016. ↑1345, 1383
- [BLY94] Jean-Pierre Bourguignon, Peter Li, and Shing-Tung Yau, *Upper bound for the first eigenvalue of algebraic submanifolds*, Comment. Math. Helv. **69** (1994), no. 2, 199–207. ↑1364
- [BMS94] Martin Bordemann, Eckard Meinrenken, and Martin Schlichenmaier, *Toeplitz quantization of Kähler manifolds and $\mathrm{gl}(N)$, $N \rightarrow \infty$ limits*, Commun. Math. Phys. **165** (1994), no. 2, 281–296. ↑1349, 1371

- [CDS07] Giulio Chiribella, Giacomo Mauro D’Ariano, and Dirk Schlingemann, *How continuous quantum measurements in finite dimensions are actually discrete*, Phys. Rev. Lett. **98** (2007), no. 19, 190043 (4 pages). ↑1346
- [CL06] Ronald Raphaël Coifman and Stéphane Lafon, *Diffusion maps*, Appl. Comput. Harmon. Anal. **21** (2006), no. 1, 5–30. ↑1378
- [CP18a] Laurent Charles and Leonid Polterovich, *Quantum speed limit versus classical displacement energy*, Ann. Henri Poincaré **19** (2018), no. 4, 1215–1257. ↑1352, 1353
- [CP18b] ———, *Sharp correspondence principle and quantum measurements*, St. Petersburg. Math. J. **29** (2018), no. 1, 177–207. ↑1345, 1354
- [CR12] Monique Combescure and Didier Robert, *Coherent states and applications in mathematical physics*, Theoretical and Mathematical Physics, Springer, 2012. ↑1382
- [Del] Alix Deleporte, *private correspondence*. ↑1353
- [Dia88] Persi Diaconis, *Group Representations in Probability and Statistics*, Institute of Mathematical Statistics Lecture Notes – Monograph Series, vol. 11, Institute of Mathematical Statistics, 1988. ↑1379, 1380, 1381, 1382
- [DLM06] Xianzhe Dai, Kefeng Liu, and Xiaonan Ma, *On the asymptotic expansion of Bergman kernel*, J. Differ. Geom. **72** (2006), no. 1, 1–41. ↑1345, 1353, 1356, 1362, 1373
- [Don09] Simon Kirwan Donaldson, *Some numerical results in complex differential geometry*, Pure Appl. Math. Q. **5** (2009), no. 2, 571–618, Special Issue: In honor of Friedrich Hirzebruch. Part 1. ↑1345, 1350, 1351, 1352, 1363, 1364, 1365, 1366, 1367, 1382
- [EF02] Yonina Chana Eldar and G. David Jun. Forney, *Optimal tight frames and quantum measurement*, IEEE Trans. Inf. Theory **48** (2002), no. 3, 599–610. ↑1346
- [Eng00] Miroslav Engliš, *A Forelli–Rudin construction and asymptotics of weighted Bergman kernels*, J. Funct. Anal. **177** (2000), no. 2, 257–281. ↑1362
- [Far99] Richard William Farebrother, *Fitting Linear Relationships: A History of the Calculus of Observations 1750–1900*, Springer Series in Statistics, Springer, 1999. ↑1374
- [Fin10] Joel Fine, *Calabi flow and projective embeddings*, J. Differ. Geom. **84** (2010), no. 3, 489–523, with an appendix by Kefeng Liu and Xiaonan Ma. ↑1351
- [Fin12a] ———, *Notes on canonical Kähler metrics and quantisation*, 2012, preprint, available at http://homepages.vub.ac.be/~joelfine/preprints/2012-07_Cologne.pdf. ↑1351, 1364, 1374
- [Fin12b] ———, *Quantization and the Hessian of Mabuchi energy*, Duke Math. J. **161** (2012), no. 14, 2753–2798. ↑1345, 1351
- [FM05] Hermann Flaschka and John Millson, *Bending flows for sums of rank one matrices*, Can. J. Math. **57** (2005), no. 1, 114–158. ↑1346, 1347
- [GM00] Apostolos A. Giannopoulos and Vitali D. Milman, *Extremal problems and isotropic positions of convex bodies*, Isr. J. Math. **117** (2000), 29–60. ↑1346
- [GU88] Victor W. Guillemin and Alejandro Uribe, *The Laplace operator on the n -th tensor power of a line bundle: eigenvalues which are uniformly bounded in n* , Asymptotic Anal. **1** (1988), no. 2, 105–113. ↑1361, 1362
- [Hay06] Masahito Hayashi, *Quantum Information. An introduction*, Springer, 2006. ↑1347
- [Her70] Joseph Hersch, *Quatre propriétés isopérimétriques de membranes sphériques homogènes*, C. R. Acad. Sci., Paris, Sér. A **270** (1970), 1645–1648. ↑1364
- [ILMM17] Louis Ioos, Wen Lu, Xiaonan Ma, and George Marinescu, *Berezin–Toeplitz quantization for eigenstates of the Bochner–Laplacian on symplectic manifolds*, J. Geom. Anal. **30** (2017), no. 3, 2615–2646. ↑1362
- [Isa06] I. Martin Isaacs, *Character Theory of Finite Groups*, AMS Chelsea Publishing, 2006, corrected reprint of the 1976 original. ↑1379

- [Kan77] Yakar Kannai, *Off diagonal short time asymptotics for fundamental solution of diffusion equation*, Commun. Partial Differ. Equations **2** (1977), no. 8, 781–830. ↑1354, 1358
- [Kel09] Julien Keller, *Ricci iterations on Kähler classes*, J. Inst. Math. Jussieu **8** (2009), no. 4, 743–768. ↑1364, 1373
- [KMS16] Julien Keller, Julien Meyer, and Reza Seyyedali, *Quantization of the Laplacian operator on vector bundles. I*, Math. Ann. **366** (2016), no. 3-4, 865–907. ↑1351
- [KS01] Alexander V. Karabegov and Martin Schlichenmaier, *Identification of Berezin–Toeplitz deformation quantization*, J. Reine Angew. Math. **540** (2001), 49–76. ↑1344, 1345, 1352, 1353, 1356
- [Lan98] Nicolaas P. Landsman, *Mathematical topics between classical and quantum mechanics*, Springer Monographs in Mathematics, Springer, 1998. ↑1345
- [LF18] Yohann Le Floch, *A Brief Introduction to Berezin–Toeplitz Operators on Compact Kähler Manifolds*, CRM Short Courses, Springer, 2018. ↑1349, 1354
- [LM07] Kefeng Liu and Xiaonan Ma, *A remark on: “Some numerical results in complex differential geometry”*, Math. Res. Lett. **14** (2007), no. 2, 165–171. ↑1345, 1351, 1353, 1357
- [LM10] Gilles Lebeau and Laurent Michel, *Semi-classical analysis of a random walk on a manifold*, Ann. Probab. **38** (2010), no. 1, 277–315. ↑1345, 1348, 1353
- [LMM17] Wen Lu, Xiaonan Ma, and George Marinescu, *Donaldson’s Q -operators for symplectic manifolds*, Sci. China, Math. **60** (2017), no. 6, 1047–1056. ↑1345, 1362
- [MM07] Xiaonan Ma and George Marinescu, *Holomorphic Morse Inequalities and Bergman Kernels*, Progress in Mathematics, vol. 254, Birkhäuser, 2007. ↑1355, 1361, 1362, 1364
- [MM08] ———, *Generalized Bergman kernels on symplectic manifolds*, Adv. Math. **217** (2008), no. 4, 1756–1815. ↑1345, 1362
- [MM12] ———, *Berezin–Toeplitz quantization on Kähler manifolds*, J. Reine Angew. Math. **662** (2012), 1–56. ↑1351
- [NC00] Michael A. Nielsen and Isaac L. Chuang, *Quantum Computation and Quantum Information*, Cambridge University Press, 2000. ↑1382
- [OC09] Ognian Oreshkov and John Calsamiglia, *Distinguishability measures between ensembles of quantum states*, Phys. Rev. A **79** (2009), no. 3, 032336. ↑1346, 1375, 1384
- [RSVN18] Ophélie Rouby, Johannes Sjostrand, and San Vu Ngoc, *Analytic Bergman operators in the semiclassical limit*, <https://arxiv.org/abs/1808.00199>, 2018. ↑1353
- [Rud13] Daniel Rudolf, *An introduction to Markov chain Monte Carlo on general state spaces*, 2013, preprint, available at <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.461.7352&rep=rep1&type=pdf>. ↑1348
- [Sch10] Martin Schlichenmaier, *Berezin–Toeplitz quantization for compact Kähler manifolds. A review of results*, Adv. Math. Phys. **2010** (2010), 927280 (38 pages). ↑1349
- [Wal18] Shayne F. D. Waldron, *An Introduction to Finite Tight Frames*, Applied and Numerical Harmonic Analysis, Birkhäuser; Springer, 2018. ↑1378
- [Zha98] Genkai Zhang, *Berezin transform on compact Hermitian symmetric spaces*, Manuscr. Math. **97** (1998), no. 3, 371–388. ↑1352, 1382

Manuscript received on 24th February 2019,
 revised on 23rd September 2019,
 accepted on 19th February 2020.

Recommended by Editor N. Anantharaman.
 Published under license CC BY 4.0.



This journal is a member of Centre Mersenne.



Louis IOOS

louisioos@mail.tau.ac.il
School of Mathematical Sciences,
Tel Aviv University,
Tel Aviv, 69978, (Israel)

Victoria KAMINKER

vika999@gmail.com
School of Mathematical Sciences,
Tel Aviv University,
Tel Aviv, 69978, (Israel)

Leonid POLTEROVICH

polterov@tauex.tau.ac.il
School of Mathematical Sciences,
Tel Aviv University,
Tel Aviv, 69978, (Israel)

Dor SHMOISH

dorshmoish@mail.tau.ac.il
School of Mathematical Sciences,
Tel Aviv University,
Tel Aviv, 69978, (Israel)