



ANNALES
HENRI LEBESGUE

ALBERTO CATTANEO

IVAN CONTRERAS

SPLIT CANONICAL RELATIONS

RELATIONS CANONIQUES SCINDÉES

ABSTRACT. — A Lagrangian subspace L of a weak symplectic vector space is called *split Lagrangian* if it has an isotropic (hence Lagrangian) complement. When the symplectic structure is strong, it is sufficient for L to have a closed complement, which can then be moved to become isotropic. The purpose of this note is to develop the theory of compositions and reductions of split canonical relations for symplectic vector spaces. We give conditions on a coisotropic subspace C of a weak symplectic space V which imply that the induced canonical relation L_C from V to C/C^ω is split, and, from these, we find sufficient conditions for split canonical relations to compose well. We prove that the canonical relations arising in the Poisson sigma model from the Lagrangian field theoretical approach are split, giving a description of symplectic groupoids integrating Poisson manifolds in terms of split canonical relations.

RÉSUMÉ. — Un sous-espace Lagrangien L d'un espace vectoriel symplectique faible est appelé *Lagrangien scindé* s'il a un complément isotrope (donc Lagrangien). Lorsque la structure symplectique est forte, il suffit que L ait un complément fermé, qui peut ensuite être déplacé pour devenir isotrope. Le but de cette note est de développer la théorie des compositions et des réductions des relations canoniques scindées pour les espaces vectoriels symplectiques. Nous donnons des conditions sur un sous-espace coisotrope C d'un espace symplectique faible V qui impliquent que la relation canonique induite L_C de V à C/C^ω est scindée, et en déduisons des conditions suffisantes pour que les relations canoniques scindées soient composables. Nous prouvons que les relations canoniques résultant du modèle sigma de Poisson dans l'approche lagrangienne de la théorie des champs sont scindées, donnant une description des groupoïdes symplectiques intégrant les variétés de Poisson en termes de relations canoniques scindées.

Keywords: Canonical relations, symplectic reduction, Poisson sigma model, symplectic groupoids.
2020 *Mathematics Subject Classification:* 53D22, 17B63, 53D17.

DOI: <https://doi.org/10.5802/ahl.69>

(*) This research was (partly) supported by the NCCR SwissMAP, funded by the Swiss National Science Foundation, and by the COST Action MP1405 QSPACE, supported by COST (European Cooperation in Science and Technology). I.C. was partially supported by SNF Grant P300P2-154552. A. S. C. acknowledges partial support of SNF Grant No. 200020_172498/1.

1. Introduction

The functorial description of classical and quantum field theories requires enhancement of the usual symplectic category, allowing infinite dimensional objects (weak symplectic manifolds), and allowing general canonical relations, not just symplectomorphisms, as morphisms. As is explained in [Wei10], the construction of such an extended symplectic category encounters several obstacles, even in finite dimensions, and many alternatives have been proposed over time. These constructions face even more technical difficulties when infinite dimensional spaces are taken into account.

One of the problems arises from the nature of the objects of this extended category: the manifolds which model the space of fields for classical field theories are in general Fréchet manifolds, and conventional tools such as the inverse function theorem, Frobenius theorem, etc. fail to hold in general in the Fréchet category.

Another inconvenience, which is a crucial impediment to establishing functorial quantization, is the fact that the composition of canonical relations is not well defined in general. In the finite dimensional case they compose well when certain transversality conditions are satisfied; in infinite dimensions, we can find examples of canonical relations whose transversal composition is isotropic but not coisotropic (see e.g. [Con13]).

The objective of this paper is to analyze the special case in which we allow the objects to be (infinite dimensional) weak symplectic vector spaces and the morphisms to be *split canonical relations*, which are isotropic subspaces with isotropic complements. This notion of Lagrangian subspace was first considered in the context of infinite dimensional symplectic linear spaces in [Wei71], and it happens to coincide with the conventional notion (maximal isotropic) in the finite dimensional case (see e.g. [Arn67, Wei71]). For the infinite dimensional case, we determine sufficient conditions for split canonical relations to compose well and to have well defined reductions.

Split Lagrangian subspaces appear naturally in the Hamiltonian formulation for Lagrangian field theories with boundary, following the BV-BFV formalism, as in [CMR12]. This type of field theory naturally produces an isotropic submanifold L of the symplectic space of boundary fields $\mathcal{F}_{\partial M}$. A *good* choice of a boundary conditions for this theory consists of the choice of a submanifold L' on which the Noether 1-form vanishes and which intersects L in isolated points (the solutions to the Euler–Lagrange system of equations).

For this choice of boundary conditions, we want to insure that the variational problem has no boundary contributions. This condition implies first of all that L' must be isotropic. Globally, this condition is even stronger, as it requires the vanishing of the class of the Noether 1-form (changing it by exact terms is allowed as it corresponds to change the action functional by a boundary term). Locally, at a point of $L \cap L'$, the tangent spaces to L and to L' must then be complementary isotropic subspaces.

The last part of the paper (Section 6) is devoted to proving that the evolution relations arising from the Poisson sigma model (PSM), a two dimensional topological theory, are split canonical relations obeying a *neat intersection* condition which allows compositions and reductions. These considerations help us to provide an alternative and shorter proof that such evolution relations are Lagrangian for the

PSM case [CC15, Con13] and that the reduced version of these relations determines a *split* version of the conventional symplectic groupoids for integrable Poisson manifolds. We conjecture that the split condition for canonical relations is satisfied in a larger class of classical field theories with boundary. In particular, we intend to study the case in which the symplectic space of fields happens to be reflexive.

The study of split Lagrangian spaces can be naturally extended to the framework of Banach manifolds, for which the existence of split Lagrangian submanifolds implies the existence of complementary isotropic smooth distributions, which play an important role in the symplectic formulation of field theories with boundary. Also, we intend to extend the construction of split coisotropic subspaces to Banach manifolds, and adapt the existence of c -triples (Definition 3.2) to the existence and local uniqueness of coisotropic embeddings of finite dimensional presymplectic manifolds, due to Gotay [Got82].

Acknowledgements

The authors thank Alan Weinstein for his useful insights at different stages of this project, in particular for the idea of the proof of Theorem 2.10 and the finite dimensional case (Section 3.2). The authors thank the anonymous referee for the useful comments and suggestions.

2. Definitions and basic properties

In this article, we consider topological vector spaces over \mathbb{R} . Thus, the dual space V^* of V is the space of *continuous* linear functionals of V . The topology for Banach and Hilbert spaces is the metric topology, and the topology for Fréchet spaces is the initial topology, i.e. the coarsest topology for which all the seminorms are continuous⁽¹⁾.

DEFINITION 2.1. — *A closed subspace W of a vector space V is called split if it has a closed complement K in V .*

DEFINITION 2.2. — *A (possibly infinite dimensional) vector space V is called weak symplectic if it is equipped with a skew symmetric bilinear form ω such that the induced linear map $\omega^\sharp: V \rightarrow V^*$ is injective. It is symplectic if ω^\sharp is an isomorphism.*

Note that if the vector space V is symplectic, then $(V^*)^* \cong V^* \cong V$, thus V is reflexive. This does not necessarily hold for weak symplectic spaces (Proposition 2.3).

W^ω denotes the symplectic orthogonal subspace of a subspace W in a weak symplectic vector space. It is necessarily closed, since it is determined by the condition $\omega(W, \bullet) = 0$, and ω is continuous. We leave to the reader the proof of the following useful properties of symplectic orthogonality.

PROPOSITION 2.3. — *Let V be a weak symplectic space and W, Z subspaces of V . Then:*

⁽¹⁾With this topology, the Fréchet spaces are particular instances of locally convex spaces.

- (1) $W \subseteq Z \implies Z^\omega \subset W^\omega$.
- (2) $(W + Z)^\omega = W^\omega \cap Z^\omega$.
- (3) $(W \cap Z)^\omega \supseteq W^\omega + Z^\omega$.
- (4) $W \subseteq W^{\omega\omega}$.
- (5) $W^\omega = W^{\omega\omega\omega}$.

In the symplectic (e.g. finite dimensional weak symplectic) case, we have a slightly stronger result.

PROPOSITION 2.4. — *When V is symplectic and W is a closed subspace, the Inequality (4) above becomes an equality.*

Proof. — Suppose that the Inequality (4) is strict. Then there is an element v of $V \setminus W$ for which the linear functional $\omega(v, \cdot)$ annihilates W^ω . On the other hand, since W is closed, by the Hahn–Banach theorem (which holds in any Fréchet space [Rud91]), there is a linear functional which takes the value 1 on v and annihilates W . Since ω is symplectic, this linear functional is realized by some $z \in V$; i.e. $z \in W^\omega$ and $\omega(z, v) = 1$. But this contradicts the assumption that $\omega(v, \cdot)$ annihilates W^ω . \square

DEFINITION 2.5. — *A subspace L of a weak symplectic vector space V is called isotropic if $L \subseteq L^\omega$, coisotropic if $L^\omega \subseteq L$ and Lagrangian if $L = L^\omega$.*

Remark 2.6. — Note that a subspace L is Lagrangian if and only if it is maximal isotropic, and there are many of these subspaces, by Zorn’s lemma. A Lagrangian subspace is closed because it is the symplectic orthogonal space of itself.

PROPOSITION 2.7. — *If $V = L \oplus L'$, where L and L' are isotropic, then L and L' are each Lagrangian. (In particular, they are closed.)*

Proof. — If L were properly contained in an isotropic subspace L_1 , then L_1 would have a nontrivial intersection with L' , and any element of this intersection would be symplectically orthogonal to both L and L' , in contradiction to the injectivity of ω^\sharp . This shows that L is maximal isotropic, i.e. Lagrangian. Of course, the same argument applies to L' . \square

Remark 2.8. — We will call a subspace L *split Lagrangian* if it is Lagrangian and has a Lagrangian complement. For example, the spaces L and L' in Proposition 2.7 are both split Lagrangian. We will also refer to (L, L') as a splitting l-pair.

Theorem 2.10 below shows that, when the ambient space is symplectic, and not just weakly symplectic, any closed complement to a Lagrangian subspace can be moved to become Lagrangian itself. Thus, in the symplectic case, the two possible interpretations of the term “split Lagrangian subspace” are equivalent; i.e. a subspace which is split and Lagrangian is split Lagrangian. We do not know whether this remains true in the general weakly symplectic case.

LEMMA 2.9. — *If $W \oplus Z$ is a splitting of a symplectic space V by closed subspaces, then $W^\omega \oplus Z^\omega$ is also a splitting of V by closed subspaces.*

Proof. — The splitting $V = W \oplus Z$ leads to a splitting $V^* = W^0 \oplus Z^0$ of V^* into the sum of the annihilators of W and Z respectively. The inverse of the isomorphism ω^\sharp then takes W^0 to W^ω and Z^0 to Z^ω , giving the required splitting of V . \square

THEOREM 2.10. — *If V is symplectic, any Lagrangian subspace $L \subseteq V$ which is split as a subspace has a Lagrangian complement L^c in V and is hence (L, L^c) is a splitting l -pair.*

Proof. — Let K be a closed complement to L in V . By Lemma 2.9, K^ω is also a closed complement to L .

We define the *average* of K and K^ω to be

$$L^c := \left\{ \frac{1}{2} (k + k') : k \in K, k' \in K^\omega, k - k' \in L \right\}.$$

By Proposition 2.7, we just have to prove that L^c is an isotropic complement to L in V .




Figure 2.1. The average construction for a split weak symplectic vector space V .

We first prove that it is a complement. If v in L^c belongs to L as well, then we must have $k + k'$ in the definition above both belonging to L . But $k - k'$ belongs to L as well, so k and k' belong to L . But, since K and K' are complements to L , k and k' must be zero. Thus, $L^c \cap L = \{0\}$. On other hand, any element v of V may be expressed as either $l + k$ or $l' + k'$, where l and l' belong to L , $k \in K$, and $k' \in K'$. Then $k - k' = l' - l$ belongs to L , and so $l_c = \frac{1}{2}k + \frac{1}{2}k'$ belongs to L^c . Now v is the average $\frac{1}{2}(l + k + l' + k')$ of its two expressions, which is the sum of l_c and an element of L . Thus $L^c \oplus L = V$.

We will now verify that L^c is isotropic. Let $l_1^c := \frac{1}{2}k_1 + \frac{1}{2}k_1'$ and $l_2^c := \frac{1}{2}k_2 + \frac{1}{2}k_2'$ be elements in L^c . Then,

$$\begin{aligned}\omega(l_1^c, l_2^c) &= \omega\left(k_1 + \frac{1}{2}(k_1' - k_1), k_2 + \frac{1}{2}(k_2' - k_2)\right) \\ &= \omega(k_1, k_2) + \frac{1}{2}\omega(k_1, (k_2' - k_2)) - \frac{1}{2}\omega(k_2, (k_1' - k_1)) \\ &= \frac{1}{2}\omega(k_1, k_2') - \frac{1}{2}\omega(k_2, k_1') \\ &= 0 - 0 = 0.\end{aligned}$$

□

Remark 2.11. — In [KS82], the authors produce an example of a symplectic Banach space V in which no subspace is split Lagrangian. By Theorem 2.10, these subspaces do not admit any closed complements at all.

DEFINITION 2.12. — A weak canonical relation L between two weakly symplectic vector spaces V and W , denoted by $L: V \rightrightarrows W$, is a Lagrangian subspace of $\bar{V} \oplus W$. A split weak canonical relation $L: V \rightrightarrows W$, is a split Lagrangian subspace of $\bar{V} \oplus W$.

Example 2.13. — Each Lagrangian subspace $L \subseteq V$ may be identified with the canonical relation $\{0\} \times L \subseteq \{0\} \oplus V$ from the zero dimensional vector space $\{0\}$ to V .

Example 2.14. — It is easy to check that, for any weakly symplectic vector space V , the diagonal relation

$$\text{graph}(\text{Id}_V) = \Delta_V: V \rightrightarrows V$$

is isotropic and has the anti-diagonal $-\Delta_V$ as an isotropic complement. Thus, Δ_V is a split canonical relation (as is $-\Delta_V$).

DEFINITION 2.15. — Let C be a coisotropic subspace of V and $\underline{C} := C/C^\omega$ its symplectic reduction (C/C^ω is a weakly symplectic vector space). The reduction relation, denoted by $L_C: V \rightrightarrows \underline{C}$, is the graph

$$L_C := \{(x, [x]) \in V \oplus \underline{C} : x \in C\}$$

of the quotient map from C onto \underline{C} , considered as having source object V .

PROPOSITION 2.16. — L_C is a weak canonical relation.

Proof. — The fact that L_C is isotropic easily follows from

$$\underline{\omega}([x], [y]) = \omega(x, y).$$

Furthermore, consider

$$L_C^\omega = \{(x, y) \in V \oplus \underline{C} : \omega(x, x') - \underline{\omega}(y, [x']) = 0, \forall (x', [x']) \in L_C\}.$$

Then, when $(x, y) \in L_C^\omega$,

$$\omega(x, x') - \underline{\omega}(y, [x]) = \underline{\omega}([x], [x']) - \underline{\omega}(y, [x']) = \underline{\omega}([x] - y, [x']) = 0, \forall [x'] \in \underline{C},$$

and since \underline{C} is symplectic, it implies that $[x] = y$ and hence $(x, y) \in L_C$. □

Remark 2.17. — As we will see later, L_C is not in general split Lagrangian: a sufficient condition for this is to have suitable complements of C and C^ω in V and C respectively.

3. Composition and Reduction

The set-theoretic composition $L_2 \circ L_1$ of canonical relations $L_1: V \rightrightarrows W$ and $L_2: W \rightrightarrows Z$ is a linear subspace of $\overline{V} \oplus Z$. This composition is strongly transversal (i.e. the projection map from $L_C \times_V L$ onto $V \times Z$ is a closed embedding) exactly when L is transversal to C , where $C = V \oplus \Delta_W \oplus Z$.

In finite dimensions, this composition is always Lagrangian, and hence it is a canonical relation $V \rightrightarrows Z$. In infinite dimensions it may be merely an isotropic relation.

In Section 4, we will give two examples where the composition of canonical relations is not Lagrangian. In the first, the composition is closed; in the second, the weak symplectic structures are symplectic.

3.1. Equivalence between reduction and composition; split coisotropic subspaces

There is a natural correspondence (see e.g. [Wei10]) between the reduction and composition of canonical relations, that can be spelled out as follows. For a coisotropic subspace C of a symplectic vector space V , and a Lagrangian subspace L of V , the reduced Lagrangian subspace $\underline{L} \subseteq \underline{C}$, defined as:

$$\underline{L} := (L \cap C)/(L \cap C^\omega),$$

can be obtained as the following composition of canonical relations:

$$\underline{L} = L_C \circ L.$$

On the other hand, the composition of linear canonical relations is itself a particular case of reduction. It can be easily checked that the space $V \times \Delta_W \times Z$ is a coisotropic subspace of $\overline{V} \times W \times \overline{W} \times Z$; and that

$$(\overline{V} \times \Delta_W \times Z)^\omega = \{0_{\overline{V}}\} \times \Delta_W \times \{0_Z\},$$

therefore the reduced space $\overline{V} \times \Delta_W \times Z$ is naturally isomorphic to $\overline{V} \times Z$. Under this natural isomorphism, the composed relation $g \circ f$, where $f: V \rightrightarrows W$ and $g: W \rightrightarrows Z$ are canonical relations, is the reduction of the product $f \times g$, and the composition is transversal if and only if the reduction is.

The following Theorem 3.1 gives sufficient conditions for the reduction relation L_C to be split Lagrangian.

THEOREM 3.1. — *Let C be a coisotropic subspace of V . If there is a complement C^c to C in V and a complement C' to C^ω in C such that*

- (1) C^c is isotropic, and
- (2) C' and C^c are symplectically orthogonal,

then L_C is split Lagrangian.

Proof. — Given a choice of C^c and C' , we have the decomposition

$$V = C^c \oplus C^\omega \oplus C'.$$

If we identify the complement C' in the usual way with the quotient $\underline{C} = C/C^\omega$, we can write the reduction relation as

$$L_C = \left\{ (x_1^c, x_2^\omega, x_3', [x_4]) \in V \oplus \underline{C} : x_1^c = 0, x_2^\omega \in C^\omega, x_3' = [x_4] \in C' \right\}.$$

Consider now the following relation:

$$L_C^- := \left\{ (x_1^c, x_2^\omega, x_3', [x_4]) \in V \oplus \underline{C} : x_1^c \in C^c, x_2^\omega = 0, x_3' = -[x_4] \in C' \right\}.$$

Observe that $L_C + L_C^- = V \oplus \underline{C}$ and that $L_C \cap L_C^- = \{(0, 0, 0, 0)\}$. By Proposition 2.16, L_C is isotropic. Now

$$\begin{aligned} (L_C^-)^\omega &= \left\{ (y_1^c, y_2^\omega, y_3', [y_4]) \in V \oplus \overline{C} : \omega(y_1^c + y_2^\omega + y_3', x_1^c + x_3') + \underline{\omega}[y_4], x_3' \right. \\ &= 0 : \forall (x_1^c, 0, x_3', -x_3') \in L_C^- \left. \right\}. \end{aligned}$$

If $(y_1^c, y_2^\omega, y_3', [y_4]) \in L_C^-$, then $y_2^\omega = 0$ and $y_3' = -[y_4]$, hence

$$\omega(y_1^c + y_2^\omega + y_3', x_1^c + x_3') + \underline{\omega}([y_4], x_3') = \omega(y_1^c, x_1^c) + \omega(y_1^c, x_3') + \omega(y_3', x_1^c).$$

The first term on the right hand side vanishes since C^c is isotropic, by assumption (1). The second and third terms also vanish, by (2). Thus L_C^- is isotropic. \square

DEFINITION 3.2. — *A coisotropic subspace C is split coisotropic if it satisfies the assumptions of Theorem 3.1 for complements C^c and C' . The triple (C, C^c, C') will be called a splitting c-triple.*

We will describe an equivalent formulation of split coisotropic subspaces, by replacing Condition (2) with a suitable choice of a split Lagrangian subspace.

3.2. The finite dimensional case

In the case in which V is a finite dimensional symplectic space, the space of split Lagrangians is the same as the full Lagrangian Grassmannian. However, not every Lagrangian subspace L in $V \oplus W$ (where V and W are finite dimensional symplectic spaces) is of the form $L = L_C$, i.e. L is the reduction relation for some C which is split coisotropic and W is isomorphic to C/C^ω . In this section, we study the finite dimensional case of splitting c-triples and how they are explicitly related with split reduction relations and splitting l-pairs. We also find an equivalent characterization of a split coisotropic subspace C in terms of split Lagrangian complements of C^ω .

Suppose that V is a $2n$ -dimensional vector space and C is a $n + k$ -dimensional coisotropic subspace. This implies that

$$\dim(\underline{C}) = \dim(C/C^\omega) = (n + k) - (n - k) = 2k.$$

It is easy to observe that $\dim(L_C) = n + k = \dim(L_C^c)$, where L_C^c is a complement to L_C in $V \oplus \underline{C}$.

Now, we observe that the space \mathcal{L} of Lagrangian complements to L_C is open in the Lagrangian Grassmannian $\Lambda(n+k)$; therefore it has dimension $d_1 = (n+k)(n+k+1)/2$.

We will now determine “how many” complements in this space are of the form $L_{\bar{C}}$, using the construction from Theorem 3.1, i.e. to determine the dimension of the space of such special complements.

To address this question, we look first at the manifold of splitting c-pairs for C . Each such pair produces a symplectic decomposition of V :

$$V = C' \oplus (C^\omega \oplus C^c).$$

The space \mathcal{C} of choices of the complement C' is open in the Grassmannian $Gr_{2k}(n+k)$; therefore, it has dimension $d = 2k(n-k)$. Now, given a choice of C' , choosing a compatible Lagrangian complement C^c to C^ω in $(C')^\omega$ is equivalent to choosing an element in an open subset of the Lagrangian Grassmannian $\Lambda(n-k)$. Thus, the space of splitting c-triples (C', C^c) (with C' fixed) has dimension $d_2 = (n-k)(n-k+1)/2$. Now, consider the linear map

$$\begin{aligned} \phi_{C'} : \mathcal{C} &\rightarrow \mathcal{L} \\ C^c &\mapsto C^c \oplus \{0\} \oplus -\Delta_{C'}. \end{aligned}$$

It is well defined, due to Theorem 3.1, and since $-\Delta_V = -\Delta_W$ implies that $V = W$, we conclude that $\phi_{C'}$ is 1-to-1.

Using the affine structure of both spaces (the space of Lagrangian complements to L_C and the space of splitting c-triples (C, C', C^c)), and the injectivity of $\phi_{C'}$, we conclude that the number of conditions which a Lagrangian complement to L_C must satisfy in order to come from a compatible pair (C', C^c) is $d := d_1 - d_2 = (2n+1)k$.

Remark 3.3. — We can consider the extreme cases. When $k = n$, i.e. $C = V$, we have that $L_C = \Delta_C$, and d in this particular case is $(2n+1)n = 2n(2n+1)/2$ which is the full dimension of $\Lambda(2n)$. This implies that there is a unique compatible pair (C', C^c) , namely $(V, \{0\})$, which determines uniquely the anti-diagonal $-\Delta_C$ as the relation $L_{\bar{C}}$ in Theorem 3.1. The other extreme takes place when $k = 0$, i.e. C is Lagrangian, in which case $d = 0$, i.e. there are no conditions for $L_{\bar{C}}$. In this particular case, the splitting c-triples (C, C', C^c) in V are the same as the complements C' of C in V , since the reduced space \underline{C} is $C/C = 0$.

3.3. Split coisotropic subspaces

We also observe that since C is coisotropic, $C^\omega \subseteq C$ is an isotropic subspace in V , hence also in $(C')^\omega$. Therefore, Condition (2) in Definition 3.2 is equivalent to finding an isotropic complement C^c of C^ω in $(C')^\omega$, which implies that C' is a split Lagrangian subspace of $(C')^\omega$. Notice that $(C')^\omega$ is a symplectic subspace of V , with dimension $2n - 2k$. Now, let us consider a Lagrangian complement C^c to C^ω in $(C')^\omega$. We prove the following Proposition 3.4, which implies that C^c is also a complement to C in V , symplectically orthogonal to C' .

PROPOSITION 3.4. — *Any lagrangian complement C^c to C^ω in $(C')^\omega$ is also a complement to C in V .*

Proof. — Let $v \in V$. Then

$$\begin{aligned} v &= v_{C'} + v_{(C')^\omega} \\ &= v_{C'} + v_{C^\omega} + v_{C^c} \\ &= v_C + v_{C^c}, \end{aligned}$$

where v_X denotes the projection of v onto the subspace X . Thus $V = C^c + C$. Now, suppose $v \in C \cap C^c$. We have that

$$v = v_{C'} \oplus v_{C^\omega}$$

since $v \in C$. Now, for every $w \in C$, v_{C^ω} is symplectically orthogonal to w . This implies that $v_{C'}$ is also orthogonal to w , thus $v_{C'} \in C^\omega$, which implies that $v_{C'} \in C' \cap C^\omega = \{0\}$. Therefore $V = C^c \oplus C$, as we wanted. \square

This proposition suggests the following equivalent definition of split coisotropic subspace.

DEFINITION 3.5. — *(Equivalent to Definition 3.2). A coisotropic subspace C of a weakly symplectic vector space V is split coisotropic if there is a complement C' to C^ω in C and a complement L to C^ω in $(C')^\omega$, such that L is isotropic (therefore split Lagrangian).*

3.4. Good compositions

For canonical relations between symplectic manifolds, a clean intersection condition is needed to insure that the composition of two such relations is again a smooth manifold. In finite dimensions, this condition is automatically satisfied for linear canonical relations. In infinite dimensions, even in the linear case, we need to impose a condition to insure the good composition of split canonical relations.

DEFINITION 3.6. — *Let (L, L') be a splitting l -pair of V , and let (C, C^c, C') be a splitting c -triple. We say that (L, L') and C intersect neatly if the following equation holds:*

$$(3.1) \quad C' = L \cap C' + L' \cap C'.$$

Remark 3.7. — From Definition 3.5 it follows that this definition does not depend on the choice of the complement C^c .

PROPOSITION 3.8. — *The neatness condition is always satisfied in finite dimensions.*

Proof. — If L is Lagrangian, we can consider the intersection $K = L \cap C'$. Let K^c be a complement to K in C' . Using the same notation as in Section 3.2, $\dim K^c \leq 2k$. Thus, there is a choice of a Lagrangian complement L^c of L such that $K^c \subseteq L^c$. For such choice of complement, the condition in Equation (3.1) is satisfied. \square

The following Proposition 3.9 gives one sufficient condition for neat intersection in infinite dimensions, namely that the split Lagrangian L is contained in the split coisotropic space C .

PROPOSITION 3.9. — *Let C be a split coisotropic subspace of V . If (L, L^c) is a splitting l -pair such that L is contained in C , then (L, L^c) and C intersect neatly.*

Proof. — If L^c is any isotropic complement to L in V , $L^c \cap C$ is an isotropic complement to L in C , due to the modular law for isotropic spaces [LW15]. On the other hand, $\{0\}$ is a complement to $L \cap C$ in L , and these two complements are clearly symplectically orthogonal. \square

The following Theorem 3.10 guarantees that the reduction of a split Lagrangian with neat intersection gives rise to a split Lagrangian reduced subspace.

THEOREM 3.10. — *Let L be a split Lagrangian subspace of V , intersecting neatly C . Then the reduction \underline{L} is again split Lagrangian.*

Proof. — It is clear that L_C is an isotropic subspace of \underline{C} , since $L \cap C$ is an isotropic subspace of V and thus the induced symplectic structure $\underline{\omega}$ in \underline{C} vanishes on

$$\underline{L} = L \cap C / L \cap C^\omega.$$

Now, since $C' \cong \underline{C}$, the Equation (3.1) implies that the closed subspace $L^C \cap C'$ is a complement to $\underline{L} \cong L \cap C'$, and by a similar argument as before this subspace is isotropic. \square

Now, we are in a position to impose conditions insuring that the composition of split canonical relations subspaces is again a split canonical relation.

PROPOSITION 3.11. — *Let V_1, V_2, V_3 be three weak symplectic vector spaces. Then the subspace*

$$C := V_1 \oplus \Delta_{V_2} \oplus V_3 \subset V_1 \oplus \overline{V_2} \oplus V_2 \oplus \overline{V_3}$$

is a split coisotropic subspace.

Proof. — The symplectic orthogonal $C^\omega = \{0\} \oplus \Delta_{V_2} \oplus \{0\}$ of C is contained in C , which is therefore coisotropic. To show that it is in fact split coisotropic, we introduce the complement C^c of C in $V_1 \oplus \overline{V_2} \oplus V_2 \oplus \overline{V_3}$ defined as:

$$C^c := \{0\} \oplus -\Delta_{V_2} \oplus \{0\}.$$

It is isotropic since $-\Delta_{V_2}$ is isotropic. We also have the complement C' of C^ω in C defined as:

$$C' = V_1 \oplus \{0\} \oplus V_3.$$

C^c and C' are symplectically orthogonal since

$$(\omega_1 \oplus \overline{\omega_2} \oplus \omega_2 \oplus \overline{\omega_3})((0, v_2, -v_2, 0), (v_1, 0, 0, v_3)) = 0 - 0 + 0 - 0 = 0,$$

so all the conditions for a split coisotropic subspace are satisfied. \square

DEFINITION 3.12. — *Two split canonical relations $L_1 : V \rightrightarrows W$ and $L_2 : W \rightrightarrows Z$ are called neatly related if the split Lagrangian subspace $L_1 \oplus L_2$ and the subspace $V \oplus \Lambda_W \oplus Z$, which is split coisotropic by Proposition 3.11, intersect neatly in $V_1 \oplus \overline{V_2} \oplus V_2 \oplus \overline{V_3}$.*

THEOREM 3.13. — *The composition of two neatly related split canonical relations is a split canonical relation.*

Proof. — By Theorem 3.1, L_C is split Lagrangian, where

$$C = V_1 \oplus \Delta_{V_2} \oplus V_3.$$

Now, the composition $L_2 \circ L_1$ is the reduction $\underline{(L_1 \times L_2) \cap C}$ of $(L_1 \times L_2) \cap C$ in $\underline{C} = V_1 \oplus \overline{V_3}$. On the other hand, the product if $L_1 : V_1 \rightrightarrows V_2$ and $L_2 : V_2 \rightrightarrows V_3$ are split, then Now, since $(L_1 \times L_2) \cap C$ is isotropic, it follows that $\underline{(L_1 \times L_2) \cap C}$ is isotropic as well. Therefore, invoking Theorem 3.10 and the fact that we can phrase composition as a reduction (Section 3.1), we see that this implies that the space $\underline{(L_1 \times L_2) \cap C}$ is split Lagrangian in $V_1 \oplus V_3$, as we wanted. \square

4. Examples

4.1. An example in which the composition is closed but not Lagrangian

In our first example,⁽²⁾ the weak symplectic spaces are Fréchet spaces in which we will explicitly construct two canonical relations whose composition is isotropic (and closed) but not Lagrangian. Let $W := \mathcal{C}^\infty([0, 1])$ be the Fréchet space of smooth functions on the closed interval $[0, 1]$.

Then we have :

PROPOSITION 4.1. — *The form ω on $V := W \oplus W$ defined by*

$$(4.1) \quad \omega((f \oplus g), (\tilde{f} \oplus \tilde{g})) = \int_0^1 (f\tilde{g} - f\tilde{g})dt$$

is weak symplectic.

Proof. — The map given by ω is clearly skew symmetric. To check that it is weakly non degenerate, suppose that there is an element $(f \oplus g) \in V$ such that $\omega^\sharp(f \oplus g) = 0$. Set $\tilde{f} = \tilde{g} = h$, with h any function in W . Then we get that

$$\int_0^1 ((f - g)h)dt = 0, \forall h \in W,$$

and by the fundamental lemma of variational calculus we conclude that $f = g$. If we set $\tilde{f} = 0$, then

$$\int_0^1 (f\tilde{g})dt = 0, \forall \tilde{g} \in W,$$

and again by the fundamental lemma, we conclude that $f = g = 0$, hence ω^\sharp is weak symplectic. \square

⁽²⁾This example was mentioned briefly in [Con13]; here we give more details.

Remark 4.2. — Observe that ω is not symplectic: the induced map ω^\sharp is not surjective. The linear dual V^* is isomorphic (as a Fréchet space) to $\mathcal{D}([0, 1]) \oplus \mathcal{D}([0, 1])$, where $\mathcal{D}([0, 1])$ denotes the space of distributions on the unit interval.

The image of ω^\sharp is a proper dense subspace of this dual.

Consider the following (closed) subspace of V of codimension 1:

$$C := \{(f \oplus g) \in V : g(0) = 0\}.$$

PROPOSITION 4.3. — *C is a symplectic and coisotropic subspace of V , thus $\underline{C} = C$.*

Proof. — This is equivalent to proving that $C^\omega = \{(0, 0)\}$. Let $(\tilde{f} \oplus \tilde{g}) \in C^\omega$. If we set $g \equiv 0$, then we get that

$$\int_0^1 f\tilde{g} = 0, \forall f \in W,$$

and by the fundamental lemma, we can conclude that $\tilde{g} \equiv 0$. In a similar way we conclude that

$$\int_0^1 \tilde{f}g = 0, \forall g \in W : g(0) = 0,$$

and, again, by the fundamental lemma, $\tilde{f} \equiv 0$. □

Remark 4.4. — Note that, in this example, the space C is not split coisotropic. The complement to C^ω in C is all of C . It is symplectically orthogonal only to zero, which is not big enough to contain a complement to C in V .

Now let us consider the following subspace of V :

$$(4.2) \quad L = \left\{ (f \oplus g) \in V : \int_0^1 f dt = 0, \frac{dg}{dt} \equiv 0 \right\}.$$

PROPOSITION 4.5. — *L is a Lagrangian subspace of V .*

Proof. — L is clearly isotropic. Now look at

$$L^\omega = \left\{ (\tilde{f} \oplus \tilde{g}) \in : \int_0^1 (f\tilde{g} - \tilde{f}g) dt = 0 \text{ whenever } \int_0^1 f dt = 0 \text{ and } \frac{dg}{dt} \equiv 0 \right\}.$$

If we set $f \equiv 0$ and $g \equiv 1$, then we get $\int_0^1 \tilde{f} dt = 0$.

On the other hand, if we set $g \equiv 0$, we get $\int_0^1 f\tilde{g} dt = 0$ whenever $\int_0^1 f dt = 0$. Now let $F(t)$ be any function with $F(0) = 0$ and $F(1) = 0$, and set $f = \frac{dF}{dt}$, so that $\int_0^1 f dt = 0$. Integrating by parts gives

$$\int_0^1 F \frac{d\tilde{g}}{dt} = F\tilde{g} \Big|_0^1 - \int_0^1 f\tilde{g} dt = 0$$

for any F vanishing at 0 and 1. Again by the fundamental lemma, $\frac{d\tilde{g}}{dt}$ must be identically zero, so we have proved that $\tilde{f} \oplus \tilde{g}$ belongs to L when it is in L^ω ; i.e. L is Lagrangian. □

In fact, we have:

PROPOSITION 4.6. — *L is split Lagrangian.*

Proof. — Let us consider

$$L' := \left\{ (f \oplus g) : \frac{df}{dt} \equiv 0, \int_0^1 g dt = 0 \right\}.$$

Using an argument like that in Proposition 4.5, we can check that L' is Lagrangian. Since

$$(f \oplus g) = \left(f - \int_0^1 f dt \oplus \int_0^1 g dt \right) + \left(\int_0^1 f dt \oplus g - \int_0^1 g dt \right) \in L + L'$$

and

$$L \cap L' = \left\{ f \oplus g : f \equiv 2 \int_0^1 f dt, g \equiv 2 \int_0^1 g dt \right\} = \{(0, 0)\},$$

we can conclude that L is split Lagrangian. \square

Now we consider the composition $L_C \circ L$. It is easy to check that

$$L_C \circ L = \left\{ f \oplus g : \int_0^1 f = 0, g \equiv 0 \right\}.$$

PROPOSITION 4.7. — *$L_C \circ L$ is isotropic but not Lagrangian.*

Proof. — This composition is clearly isotropic, but any element of V of the form $f \oplus k$, where k is a constant, belongs to $(L_C \circ L)^\omega$. As long as f has nonzero integral or k is nonzero, this element is not in $L_C \circ L$ itself. Consequently, the reduced space $(L_C \circ L)^\omega / L_C \circ L$ is 2-dimensional, and the composition is not maximal isotropic. \square

Remark 4.8. — In this example, the reduced Lagrangian subspace is finite dimensional. It is an interesting question whether the failure of the composition to be Lagrangian, can be explained in terms of the finite dimensional reduction.

4.2. An example in which the composition is not closed

In this example, the composition of two Lagrangian relations between Hilbert spaces fails to be Lagrangian because a projection of a closed subspace of a Hilbert space is not necessarily closed.

Let $\mathcal{H} = L_2(S^1)$, the Hilbert space of L_2 -functions of the circle, and let $\mathcal{D} : \mathcal{H} \rightarrow \mathcal{H}$ be the differentiation operator

$$\begin{aligned} \mathcal{D} : \mathcal{H} &\rightarrow \mathcal{H} \\ f &\mapsto \frac{df}{d\theta}. \end{aligned}$$

Observe that \mathcal{D} is a closed operator whose domain is the dense (Sobolev) subspace of \mathcal{H} consisting of functions with distributional derivative in L_2 . Therefore, $\mathcal{G} := \text{graph } \mathcal{D}$ is a closed subspace of $\mathcal{K} := \mathcal{H} \oplus \mathcal{H}$ such that the projection $\text{pr}_1(\mathcal{G})$ to the first component \mathcal{H} is a dense subspace but not closed.

Now, consider $V := \mathcal{K} \oplus \mathcal{K}^* = \mathcal{H} \oplus \mathcal{H} \oplus \mathcal{H}^* \oplus \mathcal{H}^*$. This space is symplectic with the canonical form

$$\begin{aligned} \omega \left((f_1, g_1, \xi^1, \mu^1)(f_2, g_2, \xi^2, \mu^2) \right) \\ = \int_{S^1} \langle f_1, \xi^2 \rangle + \langle g_1, \mu^2 \rangle - \langle f_2, \xi^1 \rangle - \langle g_2, \mu^1 \rangle, f_i, g_i \in \mathcal{H}, \xi^i, \mu^i \in \mathcal{H}^*. \end{aligned}$$

PROPOSITION 4.9. — *The subspace $\mathcal{L} := \mathcal{G} \oplus \mathcal{G}^0$ is a split Lagrangian subspace of V , where \mathcal{G}^0 is the annihilator of \mathcal{G} in the dual of \mathcal{K} .*

Proof. — Note that \mathcal{G}^0 is the graph of the adjoint operator \mathcal{D}^* . Since \mathcal{D} is self adjoint, $\mathcal{G} = \mathcal{G}^0$. We also have that $\mathcal{G} \oplus \mathcal{G}$ is an isotropic subspace of V (see e.g [EM99]), and that the orthogonal complement to \mathcal{G} with respect to the inner product on $\mathcal{H} \oplus \mathcal{H}$ is $\mathcal{J}(\mathcal{G})$, where \mathcal{J} is the symplectic map

$$\begin{aligned} \mathcal{J}: \mathcal{H} \oplus \mathcal{H} &\rightarrow \mathcal{H} \oplus \mathcal{H} \\ (f, g) &\mapsto (-g, f). \end{aligned}$$

It is easy to check that this implies that $\mathcal{J}(\mathcal{G})$ is isotropic and hence $\mathcal{J}(\mathcal{G}) \oplus \mathcal{J}(\mathcal{G})$ is an isotropic complement to $\mathcal{G} \oplus \mathcal{G}^0$. \square

Remark 4.10. — An alternative way to prove Proposition 4.9 is by considering $\mathcal{G} \oplus \mathcal{G}^0$ as the conormal bundle of \mathcal{G} . We can show that if a subspace $W \subset V$ is split in V , then its conormal bundle N^*W is split Lagrangian in $V \oplus V^*$.

Now consider the space

$$C = \mathcal{H} \oplus \mathcal{H} \oplus \{0\} \oplus \mathcal{H}^* \subset V.$$

A direct calculation shows that

$$C^\omega = \mathcal{H} \oplus \{0\} \oplus \{0\} \oplus \{0\} \subset C;$$

hence C is coisotropic and its reduced weak symplectic space is $\underline{C} = \mathcal{H} \oplus \mathcal{H}^*$.

Remark 4.11. — Note that, in this example, the space C is split coisotropic. A complement to C in V is given by the isotropic subspace

$$\{0\} \oplus \{0\} \oplus \mathcal{H}^* \oplus \{0\}.$$

A complement to C^ω in C is given by the symplectic subspace

$$\{0\} \oplus \mathcal{H} \oplus \{0\} \oplus \mathcal{H}^*.$$

These are indeed symplectically orthogonal.

In order to describe the reduction $L_C(L) = \underline{L}$ of L in \underline{C} , we compute the intersection

$$\begin{aligned} \mathcal{L} \cap C &= \mathcal{G} \oplus \mathcal{G}^* \cap \mathcal{H} \oplus \mathcal{H} \oplus \{0\} \oplus \mathcal{H}^* \\ &= \mathcal{G} \oplus \{0\} \oplus \{0\}. \end{aligned}$$

It follows that the projection of $\mathcal{L} \cap C$ onto \underline{C} is $(\text{pr}_1(\mathcal{G}), 0) = (\mathcal{D}\text{om}(\mathcal{D}), 0)$. Since the domain of \mathcal{D} is a not closed subspace of \mathcal{H} , it follows that the reduction of \mathcal{L} is isotropic but not Lagrangian.

4.3. An example of a maximally isotropic subspace that is not split Lagrangian

In [KS82], a strong symplectic reflexive Banach space for which there are no split Lagrangian subspaces is constructed. The example in consideration, denoted by Z_2 , can be seen as a *twisted sum* of two Hilbert spaces isomorphic to l_2 . We will show that Z_2 has a closed maximally isotropic subspace M which is maximally isotropic and then by [KS82] Theorem 10 we can conclude that such space is not split Lagrangian.

The construction of Z_2 goes as follows. Given any real sequence $x = \{x_n\}_{n=1}^\infty$, its even subsequence $y = Ex$ is defined as $y_n = x_{2n}$, for all n . Now, if x is a sequence in l_2 , we define the sequence $\phi(x)$ as

$$\begin{aligned}\phi_{2n}(x) &= x_n \\ \phi_{2n-1}(x) &= x_n \log \frac{\|x\|_2}{|x_n|}, n \geq 1.\end{aligned}$$

Z_2 is then defined as the space of sequences x such that

$$\|x\| := \|Ex\|_2 + \|x - \phi(Ex)\|_2 < \infty.$$

In [KS82] is proven that Z_2 is indeed a reflexive Banach space, equipped with the symplectic form

$$\Omega(x, y) = \sum_{n=1}^{\infty} (x_{2n}y_{2n-1} - x_{2n-1}y_{2n}).$$

Furthermore, it is proven that the standard unit vectors $\{e_n\}_{n=1}^\infty$ in \mathbb{R}^∞ form a basis of Z_2 . Now, the subspace M is defined as the space generated by the odd unit vectors:

$$M = \langle e_1, e_3, \dots, e_{2k+1}, \dots \rangle$$

and it is in fact a closed subspace of Z_2 isomorphic to l_2 .

PROPOSITION 4.12. — *M is maximally isotropic.*

Proof. — We compute directly the symplectic orthogonal complement of M :

$$M^\omega = \left\{ \{z\}_{n=1}^\infty : x_{2n}z_{2n-1} - x_{2n-1}z_{2n} = 0, \forall x \in M \right\}.$$

Since $x_{2n} = 0$ for sequences in M , then it follows that $x_{2n-1}z_{2n} = 0$, for all $n \geq 1$. This implies that $z_{2n} = 0$ and therefore

$$M^\omega = \langle e_1, e_3, \dots, e_{2n-1}, \dots \rangle = M$$

and thus M is maximally isotropic. □

Remark 4.13. — In the introduction of [KS82] it is stated without proof that examples of maximally isotropic non split Lagrangians arise from the conormal bundle of uncomplemented closed subspaces of a reflexive Banach space X . It is still unclear to us why this is the case.

The following Section 5 provides an example of a topological field theory, the Poisson sigma model, in which split Lagrangian and split coisotropic spaces arise naturally, describing the evolution relations and Cauchy data respectively. For such spaces, the conditions of Theorems 3.1 and 3.13 are satisfied; therefore, they obey

the rules of the theory of reduction and composition of split canonical relations. As explained in the Introduction (for more details see e.g. [CMR12]), the evolution relations obtained from the symplectic formulation of such theories are always isotropic, and given *good* boundary conditions, they come equipped with closed isotropic complements. In Section 5 we will explicitly construct such isotropic complements, and we will briefly discuss an application to the connection between symplectic groupoids and Poisson structures.

5. Split Lagrangian subspaces in Lagrangian field theories

5.1. Symplectic formulation

In Lagrangian field theories, from the perspective of $BV - BFV$ theories with boundary [CMR12], the space of boundary fields $\mathcal{F}_{\partial\Sigma}$ is an infinite dimensional weak symplectic manifold. The symplectic form $\omega_{\partial\Sigma}$ is usually an exact 2-form whose primitive 1-form α can be constructed from the boundary contribution of the variational problem for the action \mathcal{S} of the theory, that is a function on the space of bulk fields \mathcal{F}_M . The theory comes equipped with a surjective submersion $\pi: \mathcal{F}_\Sigma \rightarrow \mathcal{F}_{\partial\Sigma}$.

5.1.1. The evolution relation

Given $EL(\Sigma)$, the space of solutions of the Euler–Lagrange equations on M , the projection $L_\Sigma := \pi(EL(\Sigma))$ of such solutions represents the analogue of the Hamiltonian flow for finite dimensional classical systems, although it does not correspond in general to a graph of a symplectomorphism on $\mathcal{F}_{\partial\Sigma}$. Such projections are called *evolution relations*. In [CMR12], it is proven that

THEOREM 5.1. — L_Σ is an isotropic subspace of $\mathcal{F}_{\partial\Sigma}$.

For reasonably *good* Lagrangian field theories (Scalar field theory, abelian BF -theory and Yang–Mills theory for instance), the evolution relations are Lagrangian. There are examples, such as the case of the wave equation (see e.g. [CM14]), in which the evolution relation is not always Lagrangian. In [Con13] it was proven that the evolution relations associated to Poisson sigma models where the world-sheet has zero genus are Lagrangian. We will prove that in fact, such evolution relations are split Lagrangians. We will first prove the condition in the cases where the Poisson structure is trivial or constant, and use a change of coordinates argument, to show that the other cases can be reduced to the former.

5.2. Linear Poisson sigma model

A detailed description of the symplectic formulation of the Poisson sigma model as a Lagrangian field theory with boundary can be found in [CC15, CF01, Con13]. The Poisson sigma model (PSM) corresponds to the following data:

- (1) A compact surface Σ , possibly with boundary, called the *source*.
- (2) A finite dimensional Poisson manifold (M, Π) , called the *target*, where Π is a Poisson bi-vector.

The space of fields for this theory is denoted with Φ_Σ and corresponds to the space of vector bundle morphisms of class \mathcal{C}^{k+1} between $T\Sigma$ and T^*M . This space can be parametrized by a pair (X, η) , where $X \in \mathcal{C}^{k+1}(\Sigma, M)$ and $\eta \in \Gamma^k(\Sigma, T^*\Sigma \otimes X^*T^*M)$, and $k \in \{0, 1, \dots, \infty\}$ denotes the regularity type of the map that we choose to work with. On Φ_Σ , the following first order action is defined:

$$S(X, \eta) := \int_\Sigma \langle \eta, dX \rangle + \frac{1}{2} \langle \eta, (\Pi^\# \circ X)\eta \rangle,$$

where,

$$\begin{aligned} \Pi^\# : T^*M &\rightarrow TM \\ \psi &\mapsto \Pi(\psi, \cdot). \end{aligned}$$

Here, dX and η are regarded as elements in $\Omega^1(\Sigma, X^*(TM))$, $\Omega^1(\Sigma, X^*(T^*M))$, respectively, and $\langle \cdot, \cdot \rangle$ is the pairing between $\Omega^1(\Sigma, X^*(TM))$ and $\Omega^1(\Sigma, X^*(T^*M))$ induced by the natural pairing between T_xM and T_x^*M , for all $x \in M$. We have that

$$\text{EL}(\Sigma) := \{ \text{Solutions of the Euler-Lagrange equations} \} \subset \Phi,$$

where, using integration by parts

$$\delta S = \int_\Sigma \frac{\delta \mathcal{L}}{\delta X} \delta X + \frac{\delta \mathcal{L}}{\delta \eta} \delta \eta + \text{boundary terms}.$$

The partial variations are:

$$(5.1) \quad \frac{\delta \mathcal{L}}{\delta X} = dX + \Pi^\#(X)\eta = 0,$$

$$(5.2) \quad \frac{\delta \mathcal{L}}{\delta \eta} = d\eta + \frac{1}{2} \partial \Pi^\#(X)\eta \wedge \eta = 0.$$

It follows [CC15, CF01, Con13] that the Lagrangian subspace L_Σ is the dynamical relation of PSM and that the space C_Σ defined by

$$(5.3) \quad C_\Sigma = \left\{ (X, \eta) \mid dX = \pi^\#(X)\eta, X: \partial\Sigma \rightarrow M, \eta \in \Gamma(T^*I \otimes X^*(T^*M)) \right\},$$

is the manifold defined by the first class constraints and it is a coisotropic submanifold of the space of boundary fields $\Phi_{\partial\Sigma}$.

This model has significant importance for deformation quantization. Namely, the perturbative expansion of the Feynman path integral for the PSM, in the case that Σ is a disc, gives rise to Kontsevich's star product formula [CF00, Kon03], i.e. the semiclassical expansion of the path integral

$$\int_{X(r)=x} f(X(p))g(X(q)) \exp\left(\frac{i}{\hbar} S(X, \eta)\right) dX d\eta$$

around the critical point $X(u) \equiv x, \eta \equiv 0$, where p, q and r are three distinct points of $\partial\Sigma$, corresponds to the star product $f \star g(x)$. We will consider the special case where M is a finite dimensional vector space.

5.3. Zero Poisson structure

We consider by simplicity the one dimensional linear case, i.e. $M = \mathbb{R}$, equipped with the zero Poisson structure⁽³⁾. We will restrict to the case where the worldsheet is a disk, denoted by D_n , with the boundary S^1 split into $2n$ closed intervals I intersecting at the end points. On alternating intervals we impose boundary conditions in such way that the remaining n intervals are free, on for each one of them we associate the weak symplectic space V of boundary fields:

$$V = T^*(\mathcal{C}^\infty([0, 1], \mathbb{R})),$$

where the weak symplectic structure is given by the one in Example 4.1, Equation (4.1). V is naturally identified with the cotangent bundle of the path space of M , understood as follows: V has two canonical coordinates (X, η) (position and momentum), given by

$$\begin{aligned} X &\in \mathcal{C}^\infty([0, 1], \mathbb{R}), \\ \eta &\in \Gamma^\infty(T^*[0, 1] \oplus X^*(T^*\mathbb{R})) \cong \mathcal{C}^\infty([0, 1], \mathbb{R}^* \cong \mathbb{R}). \end{aligned}$$

The boundary condition for the alternating non free intervals is given by $\eta = 0$.

Remark 5.2. — In this version of the PSM, we consider the space of boundary fields to be paths without constraints on the initial and final points. The case of fields with initial and final constraints, i.e. the paths start and end in coisotropic submanifolds of M (see e.g. [Cat14]), and periodic fields (loops instead of paths), will be consider in a subsequent work.

We first observe that the subspace of solutions of Euler–Lagrange equations is described by

$$(5.4) \quad C_\Sigma = \left\{ (X, \eta) : \frac{dX}{dt} \equiv 0 \right\} \cong \mathbb{R} \oplus \mathcal{C}^\infty([0, 1], \mathbb{R}^*).$$

Its symplectic complement is given by

$$C_\Sigma^\omega = \left\{ (X, \eta) : X = 0, \int_0^1 \eta = 0 \right\}.$$

The symplectic reduction is

$$\underline{C}_\Sigma = C_\Sigma / C_\Sigma^\omega = \mathbb{R} \oplus \mathbb{R}^* \cong T^*(\mathbb{R}).$$

PROPOSITION 5.3. — *The coisotropic subspace C_Σ is split.*

Proof. — Consider the following complement to C_Σ in V :

$$C_\Sigma^C := \mathcal{C}_0^\infty([0, 1], \mathbb{R}) \oplus \{0\},$$

where

$$\mathcal{C}_0^\infty([0, 1], \mathbb{R}) := \left\{ X \in \mathcal{C}^\infty([0, 1], \mathbb{R}) : \int_0^1 X(t)dt = 0 \right\}.$$

⁽³⁾The arguments in the sequel can be directly generalized to the case where $M = \mathbb{R}^n$.




Figure 5.1. The boundary fields for PSM with worldsheet D_3 . The blue segments represent the free intervals for which we associate the weak symplectic space V . For the dashed segments we associate fields with boundary condition $\eta = 0$.

It is a complement since the two subspaces are disjoint and any pair (X, η) in V can be written as

$$(X, \eta) = \left(\int_0^1 X(t) dt, \eta \right) + \left(X - \int_0^1 X(t) dt, 0 \right)$$

with the first component in C_Σ , the second component in C_Σ^C . For two elements $(X, 0)$ and $(X', 0)$ in C_Σ^C , we get that

$$\omega((X, 0), (X', 0)) = \int_0^1 (0 \cdot X(t) - 0 \cdot X'(t)) dt = 0,$$

Therefore C_Σ^C is an isotropic complement to C_Σ .

Now consider the following complement to C_Σ^ω in C_Σ :

$$C'_\Sigma := \left\{ (X, \eta) : \frac{dX}{dt} \equiv 0, \frac{d\eta}{dt} \equiv 0 \right\} \cong \mathbb{R} \oplus \mathbb{R}^*.$$

It is a complement since the two subspaces are disjoint any pair (X, η) in C_Σ can be written as

$$(X, \eta) = \left(0, \eta - \int_1^0 \eta dt \right) + \left(X, \int_0^1 \eta dt \right)$$

with

$$\left(0, \eta - \int_1^0 \eta dt \right) \in C_\Sigma^\omega, \left(X, \int_0^1 \eta dt \right) \in C'_\Sigma.$$

Now, if $(X, \eta) \in C_\Sigma^C$ and (X', η') in C'_Σ , we get that

$$\omega((X, \eta), (X', \eta')) = \int_0^1 - (0 \cdot X'(t) + \eta' \cdot X(t)) dt = \eta' \int_0^1 X(t) dt = 0,$$

therefore C_Σ^C and C'_Σ are symplectically orthogonal, hence C_Σ is a split coisotropic subspace, as we wanted. \square

5.4. The evolution relations

5.4.1. The relation L_Σ^1

Now, consider the disk D_1 and the following subspace L_Σ^1 of C_Σ :

$$L_\Sigma^1 := \left\{ (X, \eta) : \frac{dX}{dt} \equiv 0, \int_0^1 \eta(t) dt = 0 \right\}.$$

PROPOSITION 5.4. — L_Σ^1 is a split Lagrangian subspace of V and L_Σ^1 and C_Σ intersect neatly

Proof. — To check that L_Σ^1 is isotropic, if (X, η) and (X', η') are in L_Σ^1 , then

$$\omega((X, \eta), (X', \eta')) = X \int_0^1 \eta'(t) dt - X' \int_0^1 \eta(t) dt = 0 - 0 = 0.$$

Consider the following complement to L_Σ^1 in V :

$$(L_\Sigma^1)^C := \left\{ (X, \eta) : \frac{d\eta}{dt} \equiv 0, \int_0^1 X(t) dt = 0 \right\}.$$

It is isotropic since, if (X, η) and (X', η') are in $(L_\Sigma^1)^C$, then

$$\omega((X, \eta), (X', \eta')) = \eta \int_0^1 X'(t) dt - \eta' \int_0^1 X(t) dt = 0 - 0 = 0.$$

The neat intersection condition follows from Proposition 3.9. \square

In this case we obtain that the reduced space \underline{L}_Σ^1 is the zero section \mathbb{R} of $\mathbb{R}^* \oplus \mathbb{R}$, that is clearly split Lagrangian (it is a Lagrangian subspace of a finite dimensional symplectic vector space).

5.4.2. The relation L_Σ^2

In a similar way as before, we consider the disk D_2 for which we can define the following relation $L_\Sigma^2: V \rightrightarrows V$:

$$L_\Sigma^2 := \left\{ (X, \eta, X', \eta') \subset V \oplus V : X = X' = \text{ct}, \int_0^1 (\eta - \eta')(t) dt = 0 \right\},$$

where ct is a constant.

PROPOSITION 5.5. — L_Σ^2 is a split Lagrangian subspace of $V \oplus \bar{V}$, where L_Σ^2 and $C_\Sigma \oplus C_\Sigma$ intersect neatly.

Proof. —

To check that L_Σ^2 is isotropic, if $(X_1, \eta_1, X_1, \eta'_1)$ and $(X_2, \eta_2, X_2, \eta'_2)$ are in L_Σ^2 , then

$$\begin{aligned} & \omega \oplus \bar{\omega} \left((X_1, \eta_1, X_1, \eta'_1), (X_2, \eta_2, X_2, \eta'_2) \right) \\ &= X_1 \int_0^1 \eta_2(t) dt - X_2 \int_0^1 \eta_1(t) dt - X_1 \int_0^1 \eta'_2(t) dt + X_2 \int_0^1 \eta'_1(t) dt \\ &= X_1 \int_0^1 (\eta_2 - \eta'_2)(t) dt - X_2 \int_0^1 (\eta_1 - \eta'_1)(t) dt \\ &= 0 - 0 = 0. \end{aligned}$$

Now we consider the following subspace of $V \oplus \bar{V}$:

$$(L_\Sigma^2)^C := \left\{ (X, \eta, X', \eta') \subset V \oplus V : \int_0^1 (X + X')(t) dt = 0, \eta = -\eta', \eta = ct \right\}.$$

To check that $(L_\Sigma^2)^C$ is a complement to L_Σ^2 , we realize that any element of the space $V \oplus \bar{V}$ can be decomposed as

$$\begin{aligned} (X, \eta, X', \eta') = & \left(\frac{\int_0^1 (X + X')(t) dt}{2}, \eta + \frac{\int_0^1 (\eta - \eta')(t) dt}{2}, \frac{\int_0^1 (X + X')(t) dt}{2}, \eta' \right. \\ & \left. + \frac{\int_0^1 (\eta' - \eta)(t) dt}{2} \right) + \left(X - \frac{\int_0^1 (X + X')(t) dt}{2}, \frac{\int_0^1 (\eta' - \eta)(t) dt}{2}, \right. \\ & \left. X' - \frac{\int_0^1 (X + X')(t) dt}{2}, \frac{\int_0^1 (\eta - \eta')(t) dt}{2} \right). \end{aligned}$$

The first summand belongs to L_Σ^2 and the second one belongs to $(L_\Sigma^2)^C$ and it is a direct check that the summands are disjoint. To check that $(L_\Sigma^2)^C$ is isotropic, we check that, for two elements $(X_1, \eta_1, X'_1, -\eta'_1)$ and $(X_2, \eta_2, X'_2, -\eta'_2)$

$$\begin{aligned} & \omega \oplus \bar{\omega} \left((X_1, \eta_1, X'_1, -\eta'_1), (X_2, \eta_2, X'_2, -\eta'_2) \right) \\ &= \eta_2 \int_0^1 X_1(t) dt - \eta_1 \int_0^1 X_2(t) dt + \eta_2 \int_0^1 X'_1(t) dt - \eta_1 \int_0^1 X'_2(t) dt \\ &= \eta_2 \int_0^1 (X_1 + X'_1)(t) dt - \eta_1 \int_0^1 (X_2 + X'_2)(t) dt \\ &= 0 - 0 = 0. \end{aligned}$$

□

We can verify that the reduced relation \underline{L}_2 is the diagonal relation $\Delta_{\mathbb{R}^* \oplus \mathbb{R}} : \mathbb{R}^* \oplus \mathbb{R} \rightarrow \mathbb{R}^* \oplus \mathbb{R}$ that is split Lagrangian.

5.4.3. The relation L_Σ^3

For L_Σ^3 we consider the concatenation (composition) of paths the X - coordinate, namely, if $X(1) = X'(0)$

$$X * X'(t) = \begin{cases} X(2t) & \text{if } 0 \leq t \leq \frac{1}{2} \\ X'(2t - 1) & \text{if } \frac{1}{2} \leq t \leq 1 \end{cases}$$

Note that in general, $*$ is not a well defined operation on V , since the concatenation of two smooth paths is not in general smooth and since we have not defined the operation for the η -coordinate. However, as proven in [CF01], given two composable paths X and X' such that (X, η) and (X', η') in C_Σ , there are pairs $(\tilde{X}, \tilde{\eta})$ and $(\tilde{X}', \tilde{\eta}')$ in C_Σ such that

- (1) $((X, \eta), (\tilde{X}, \tilde{\eta}))$ and $(X', \eta'), (\tilde{X}', \tilde{\eta}')$ are elements of L_Σ^2 .
- (2) The paths $\tilde{\eta}$ and $\tilde{\eta}'$ are composable, furthermore, $\tilde{\eta}(0) = \tilde{\eta}(1) = \tilde{\eta}'(0) = \tilde{\eta}'(1) = 0$.
- (3) $(\tilde{X}, \tilde{\eta}) * (\tilde{X}', \tilde{\eta}') := (\tilde{X} * \tilde{X}', \tilde{\eta} * \tilde{\eta}') \in C_\Sigma$.

Therefore, for the next evolution relation, we can assume without loss of generality that the path concatenation is smooth. We consider the following relation

$$L_\Sigma^3 := \left\{ (X, \eta, X', \eta', X'', \eta'') \in C_\Sigma \oplus C_\Sigma \oplus C_\Sigma : ((\tilde{X} * \tilde{X}', \tilde{\eta} * \tilde{\eta}'), (X'', \eta'')) \in L_\Sigma^2 \right\},$$

that can be described by the following constraints:

$$\begin{aligned} X &= X' = X'' = ct, \\ \int_0^1 \eta''(t) dt &= \int_0^1 (\eta + \eta')(t) dt. \end{aligned}$$

PROPOSITION 5.6. — L_Σ^3 is a split Lagrangian subspace of $V \oplus V \oplus \bar{V}$ intersecting neatly $C_\Sigma \oplus C_\Sigma \oplus C_\Sigma$.

Proof. — To check that L_Σ^3 is isotropic, if we consider to elements $(X_1, \eta_1, X_1, \eta'_1, X_1, \eta''_1)$ and $(X_2, \eta_2, X_2, \eta'_2, X_2, \eta''_2)$ in L_Σ^3 , then

$$\begin{aligned} &\omega \oplus \omega \oplus \bar{\omega} \left((X_1, \eta_1, X_1, \eta'_1, X_1, \eta''_1), (X_2, \eta_2, X_2, \eta'_2, X_2, \eta''_2) \right) \\ &= X_1 \int_0^1 \eta_2(t) dt - X_2 \int_0^1 \eta_1(t) dt + X_1 \int_0^1 \eta'_2(t) dt - X_2 \int_0^1 \eta'_1(t) dt \\ &\quad - X_1 \int_0^1 \eta''_2(t) dt + X_2 \int_0^1 \eta''_1(t) dt \\ &= X_1 \int_0^1 (\eta_2 + \eta'_2 - \eta''_2)(t) dt - X_2 \int_0^1 (\eta_1 + \eta'_1 - \eta''_1)(t) dt \\ &= 0 - 0 = 0. \end{aligned}$$

Now, let us consider the following complement $(L_\Sigma^3)^C$ of L_Σ^3 in $V \oplus V \oplus \bar{V}$:

$$(L_\Sigma^3)^C := \left\{ \begin{aligned} (X, \eta, X', \eta', X'', \eta'') \in V \oplus V \oplus \bar{V} : \\ \eta = \eta' = -\eta'' = ct, \\ \int_0^1 (X + X' + X'')(t) dt = 0. \end{aligned} \right\}.$$

To check that it is a complement, we consider the following decomposition of any element $(X, \eta, X', \eta', X'', \eta'')$ in $V \oplus V \oplus V$, where $\sigma_3(X) := \frac{1}{3} \int_0^1 (X + X' + X'')(t) dt$ and $\sigma_3(\eta) := \frac{1}{3} \int_0^1 (\eta + \eta' + \eta'')(t) dt$:

$$\begin{aligned} (X, \eta, X', \eta', X'', \eta'') &= \left(\sigma_3(X), \eta - \sigma_3(\eta), \sigma_3(X), \eta' - \sigma_3(\eta), \sigma_3(X), \eta'' + \sigma_3(\eta) \right) \\ &\quad + \left(X - \sigma_3(X), \sigma_3(\eta), X' - \sigma_3(X), \sigma_3(\eta), X'' - \sigma_3(X), -\sigma_3(\eta) \right). \end{aligned}$$

It can be checked that the summands are disjoint.

To check that $(L_\Sigma^3)^C$ is isotropic, if we consider to elements $(X_1, \eta_1, X'_1, \eta_1, X''_1, -\eta_1)$ and $(X_2, \eta_2, X'_2, \eta_2, X''_2, -\eta_2)$ in L_Σ^3 , then

$$\begin{aligned} \omega \oplus \omega \oplus \bar{\omega} &\left((X_1, \eta_1, X'_1, \eta_1, X''_1, -\eta_1), (X_2, \eta_2, X'_2, \eta_2, X''_2, -\eta_2) \right) \\ &= \eta_2 \int_0^1 X_1(t) dt - \eta_1 \int_0^1 X_2(t) dt + \eta_2 \int_0^1 X'_1(t) dt - \eta_1 \int_0^1 X'_2(t) dt \\ &\quad - \eta_1 \int_0^1 X''_2(t) dt + \eta_2 \int_0^1 X''_1(t) dt \\ &= \eta_2 \int_0^1 (X_1 + X'_1 + X''_1)(t) dt - \eta_1 \int_0^1 (X_2 + X'_2 + X''_2)(t) dt \\ &= 0 - 0 = 0. \end{aligned}$$

□

The reduced relation \underline{L}_3 can be described as

$$\begin{aligned} \underline{L}_3 &= \left\{ \left((\mu_1, x), (\mu_2, x), (\mu_1 + \mu_2, x) \right), x \in \mathbb{R}, \mu_i \in \mathbb{R}^* \right\} \\ &= (\text{graph})(+_x): T^*\mathbb{R} \oplus T^*\mathbb{R} \rightarrow T^*\mathbb{R}, \end{aligned}$$

where $+_x$ denotes the fiber addition in the cotangent bundle of \mathbb{R} . It can be checked directly that this is a split canonical relation.

5.5. The symplectic case

We will consider now the case where

$$M = \mathbb{R}^2 \quad \text{and} \quad \Pi = \Omega^{-1} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}.$$

In this case, the space of solutions of Euler–Lagrange equations is given by

$$\begin{aligned} C_\Sigma &= \left\{ (X, \eta) : \frac{dX}{dt} = \Omega^{-1}\eta \right\} \\ &= \left\{ \left(X, \Omega \frac{dX}{dt} \right), X \in C^\infty([0, 1], \mathbb{R}^2) \right\} \\ &\cong C^\infty([0, 1], \mathbb{R}^2). \end{aligned}$$

PROPOSITION 5.7. — C_Σ is split coisotropic.

Proof. — We compute the symplectic orthogonal subspace C_Σ^ω :

$$C_\Sigma^\omega = \left\{ \left((X, \eta) \in V : X \in C_0^\infty([0, 1], \mathbb{R}^2), \eta = \Omega \frac{dX}{dt} \right) \right\} \subset C_\Sigma,$$

where

$$C_0^\infty([0, 1], \mathbb{R}^2) := \left\{ X \in C^\infty([0, 1], \mathbb{R}^2) : X(0) = X(1) = 0 \right\}.$$

The symplectic reduction is

$$\underline{C}_\Sigma = C_\Sigma / (C_\Sigma)^\perp = \mathbb{R}^2 \oplus \mathbb{R}^2.$$

Now consider the following complement $(C_\Sigma)^C$ of C_Σ in V :

$$(C_\Sigma)^C := \left\{ (X, 0) : \int_0^1 X(t) dt = 0 \right\}.$$

It is isotropic since the η -component of $(C_\Sigma)^C$ is always zero. To check that it is a complement, we consider the following disjoint decomposition of (X, η) in V : If $\widetilde{X} := X(t) - \omega^{-1} \int_0^t \eta(t) dt$, then

$$(X, \eta) = \left(\widetilde{X} - \int_0^1 \widetilde{X}(t) dt, 0 \right) + \left(\Omega^{-1} \left(\int_0^t \eta(t) dt + \int_0^1 \eta(t) dt \right), \eta \right)$$

Now consider the following complement $(C_\Sigma)'$ of $(C_\Sigma)^\omega$ in C_Σ :

$$(C_\Sigma)' := \left\{ (X_0 + X_1 t, \Omega X_1) : X_0, X_1 \in \mathbb{R}^2 \right\}.$$

It is a complement since the following is a disjoint decomposition of elements $(X, \Omega \frac{dX}{dt})$ in C_Σ :

$$\begin{aligned} \left(X, \Omega \frac{dX}{dt} \right) &= \left(X - (X(0) - (X(1) - X(0))t), \Omega \left(\frac{dX}{dt} - (X(1) - X(0)) \right) \right) \\ &\quad + \left(X(0) - (X(1) - X(0))t, \Omega(X(1) - X(0)) \right), \end{aligned}$$

where the first summand is in $(C_\Sigma)^\omega$ and the second summand is in $(C_\Sigma)'$.

Now, we check that $(C_\Sigma)^C$ and $(C_\Sigma)'$ are symplectically orthogonal:

$$\omega \left((X, 0), (X'_0 + X'_1 t, \Omega X'_1) \right) = \Omega X'_1 \int_0^1 X(t) dt = 0.$$

□

The following is the relation L_Σ^1 for this case:

$$L_\Sigma^1 = \left\{ \left(X, \Omega \frac{dX}{dt} \right) : X(0) = X(1) \right\}.$$

PROPOSITION 5.8. — L_Σ^1 is split Lagrangian intersecting neatly C_Σ .

Proof. — L_Σ^1 is isotropic since

$$\omega \left(\left(X, \Omega \frac{dX}{dt} \right), \left(X', \Omega \frac{dX'}{dt} \right) \right) = X \Omega X' \Big|_0^1 = 0.$$

Its isotropic complement $(L_\Sigma^1)^C$ is given by

$$(L_\Sigma^1)^C := \left\{ (X, \text{ct}) : \int_0^1 X(t) dt = 0 \right\}.$$

It is clearly isotropic. We verify that it is in fact a complement. If we denote

$$\tilde{\eta}(t) := \int_0^t \left(\eta(t) dt - \int_0^1 \eta(t) \right) dt$$

and

$$\tilde{X} = \int_0^1 X(t) - \omega^{-1} \tilde{\eta}(t) dt$$

(X, η) can be decomposed as

$$(X, \eta) = \left(\omega^{-1} \tilde{\eta} + \tilde{X}, \eta - \int_0^1 \eta(t) dt \right) + \left(X - \omega^{-1} \tilde{\eta} - \tilde{X}, \int_0^1 \eta(t) dt \right),$$

in which the first component belongs to L_Σ^1 and the second one to $(L_\Sigma^1)^C$. It is disjoint since

$$\begin{aligned} \eta - \int_0^1 \eta(t) dt = \int_0^1 \eta(t) dt &\iff \eta \equiv 0; \\ \omega^{-1} \tilde{\eta} + \tilde{X} = X - \omega^{-1} \tilde{\eta} - \tilde{X} &\iff X = 2 \int_0^1 X(t) dt \iff X \equiv 0. \end{aligned}$$

□

The following is the description of L_Σ^2 in the symplectic case:

$$L_2 = \left\{ \left(X, \Omega \frac{dX}{dt} \right), \left(X', \Omega \frac{dX'}{dt} \right) : X(0) = X'(0), X(1) = X'(1) \right\}.$$

PROPOSITION 5.9. — L_Σ^2 is a split Lagrangian subspace intersecting neatly $C_\Sigma \oplus C_\Sigma$

Proof. — L_Σ^2 is isotropic since

$$\begin{aligned} \omega \oplus \bar{\omega} \left(X_1, \Omega \frac{dX_1}{dt}, X'_1, \Omega \frac{dX'_1}{dt}, X_2, \Omega \frac{dX_2}{dt}, X'_2, \Omega \frac{dX'_2}{dt} \right) \\ = \int_0^1 X_1(t) \Omega \frac{dX_2}{dt} - \int_0^1 X_2(t) \Omega \frac{dX_1}{dt} - \int_0^1 X'_1(t) \Omega \frac{dX'_2}{dt} + \int_0^1 X'_2(t) \Omega \frac{dX'_1}{dt} \\ = X_1(1) \Omega X_2(1) - X_1(0) \Omega X_2(0) - X'_1(1) \Omega X'_2(1) + X'_1(0) \Omega X'_2(0) = 0. \end{aligned}$$

Now, we consider the following isotropic complement $(L_\Sigma^2)^C$ of L_Σ^2 in $V \oplus \bar{V}$:

$$(L_\Sigma^2)^C := \left\{ (X, \text{ct}, X', \text{ct}') : \int_0^1 X(t) dt = \int_0^1 X'(t) dt = 0 \right\}.$$

It is clearly isotropic; in order to check that it is a complement, we consider the following decomposition of $V \oplus \underline{V}$: If (X, η, X', η') is an element of $V \oplus \bar{V}$ and if we set

$$\begin{aligned} \tilde{\eta} &:= \eta - \int_0^1 \eta(t) dt, \\ \tilde{\eta}' &:= \eta' - \int_0^1 \eta'(t) dt, \\ \tilde{X} &:= \int_0^1 \left(X - \Omega^{-1} \int_0^t \tilde{\eta} \right) (t) dt, \\ \tilde{X}' &:= \int_0^1 \left(X' - \Omega^{-1} \int_0^t \tilde{\eta}' \right) (t) dt, \end{aligned}$$

then

$$\begin{aligned} &(X, \eta, X', \eta') \\ &= \left(\Omega^{-1} \int_0^t \tilde{\eta}(t) dt + \tilde{X}, \eta - \int_0^1 \eta(t) dt, \Omega^{-1} \int_0^t \tilde{\eta}'(t) dt + \tilde{X}', \eta' - \int_0^1 \eta'(t) dt \right) \\ &+ \left(X - \Omega^{-1} \int_0^t \tilde{\eta}(t) dt - \tilde{X}, \int_0^1 \eta(t) dt, X' - \Omega^{-1} \int_0^t \tilde{\eta}'(t) dt - \tilde{X}', \int_0^1 \eta'(t) dt \right). \end{aligned}$$

It is a disjoint decomposition since

$$\begin{aligned} \eta - \int_0^1 \eta(t) dt = \int_0^1 \eta(t) dt &\iff \eta \equiv 0 \iff \tilde{\eta} \equiv 0 \iff \tilde{X} = \int_0^1 X(dt), \\ X - \Omega^{-1} \int_0^t \tilde{\eta}(t) - \tilde{X} = \Omega^{-1} \int_0^t \tilde{\eta}(t) dt + \tilde{X} &\iff X = 2 \int_0^1 X(t) dt \iff X \equiv 0. \end{aligned}$$

□

The following is a description of L_Σ^3 in the symplectic case:

$$\begin{aligned} L_\Sigma^3 &= \left\{ X, \Omega \left(\frac{dX}{dt} \right), X', \Omega \left(\frac{dX'}{dt} \right), X'', \Omega \left(\frac{dX''}{dt} \right) \in C_\Sigma \oplus C_\Sigma \oplus C_\Sigma : X(1) \right. \\ &= \left. X'(0), X''(0) = X(0), X''(1) = X'(1) \right\}. \end{aligned}$$

PROPOSITION 5.10. — L_Σ^3 is a split Lagrangian subspace intersecting neatly $C_\Sigma \oplus C_\Sigma \oplus C_\Sigma$.

Proof. — L_Σ^3 is isotropic since

$$\begin{aligned} &\omega \oplus \omega \oplus \bar{\omega} \left(X_1, \Omega \left(\frac{dX_1}{dt} \right), X'_1, \Omega \left(\frac{dX'_1}{dt} \right), X''_1, \Omega \left(\frac{dX''_1}{dt} \right) \right), \\ &\left(X_2, \Omega \left(\frac{dX_2}{dt} \right), X'_2, \Omega \left(\frac{dX'_2}{dt} \right), X''_2, \Omega \left(\frac{dX''_2}{dt} \right) \right) X_1(1) \Omega X_2(1) - X_1(0) \Omega X_2(0) \\ &+ X'_1(1) \Omega X'_2(1) - X'_1(0) \Omega X'_2(0) - X''_1(1) \Omega X''_2(1) - X''_1(0) \Omega X''_2(0) = 0. \end{aligned}$$

Similarly as the previous case, the following is an isotropic complement $(L_\Sigma^3)^C$ of L_Σ^3 in $V \oplus V \oplus \bar{V}$:

$$(L_\Sigma^3)^C := \left\{ (ct, \eta, ct', \eta', ct'', \eta'') \in V \oplus V \oplus \bar{V} : \int_0^1 \eta(t) dt = \int_0^1 \eta'(t) dt = \int_0^1 \eta''(t) dt = 0 \right\}. \quad \square$$

5.6. The constant case

The case where $M = \mathbb{R}^n$ and Π is a constant Poisson bivector is a combination of the previous two cases. More precisely, if Π is constant, we can decompose Π as a direct sum $\Omega \oplus 0$, M as a direct sum $M_\Omega \oplus M_0$ and V as $V_\Omega \oplus V_0$. This decomposition induces a decomposition of the Euler–Lagrange space

$$C_\Sigma = C_\Omega \oplus C_0.$$

It is easy to check that the direct sum of split coisotropic subspaces and split Lagrangians subspaces is again split, therefore the following proposition holds:

PROPOSITION 5.11. — *If Π is a constant Poisson bivector on M , then C_Σ is a split coisotropic subspace of V , and similarly the evolutions relations L_1, L_2 and L_3 intersecting neatly $C_\Sigma, C_\Sigma \oplus C_\Sigma$ and $C_\Sigma \oplus C_\Sigma \oplus C_\Sigma$ respectively.*

5.7. The general case

We will prove that the above considerations for C_Σ and the relations $L_\Sigma^1, L_\Sigma^2, L_\Sigma^3$ can be naturally extended for any Poisson bivector field Π on $M = \mathbb{R}^n$. In [Cat14], the linearized version of the Euler–Lagrange equations is modified via a change of coordinates, and the resulting Euler–Lagrange space is identified with the solutions (λ, φ) for the equation $\frac{d\lambda}{dt} = -\tilde{P}^\sharp \varphi$ where \tilde{P}^\sharp is constant. The change of coordinates leaves invariant the weak symplectic structure of V and hence the splitting condition for coisotropic and Lagrangian subspaces is preserved. In order to show this, let us briefly recall the change of coordinates from [Cat14]. If we fix an element (X, η) in C_Σ and if we choose a linear connection ∇ on M , it induces a connection ∇^* in X^*TM , described in local coordinates by the following exterior covariant derivative ∂ :

$$\begin{aligned} \partial : \Gamma(X^*TM) &\rightarrow \Gamma(T^*[0, 1] \otimes X^*TM) : \\ \sigma^i &\mapsto \partial \sigma^i := d\sigma^i + \Gamma_{rs}^i dx^r \sigma^s, \end{aligned}$$

where Γ_{rs}^i are the Christoffel symbols for ∇ . We can now deform this connection using the Poisson bivector Π , namely, if we denote

$$A := \nabla^* \Pi^\sharp(X)\eta$$

described locally by

$$(\nabla^* \Pi)_k^{ij} = \partial_k \Pi^{ij} + \Gamma_{kr}^i \Pi^{rj} + \Gamma_{kr}^j \Pi^{ir},$$

we can construct the covariant derivative

$$D := \partial + A.$$

[Cat14, Proposition 2.1] shows that the space $T_{(X,\eta)}C_\Sigma$ can be written as follows:

$$(5.5) \quad T_{(X,\eta)}C_\Sigma = \left\{ (\xi, e) \in \Gamma(X^*TM) \oplus \Gamma(T^*[0, 1] \otimes X^*T^*M) : D\xi = -\Pi^\sharp(X)e \right\}.$$

Furthermore, if U denotes the parallel transport of D , the following change of coordinates:

$$(5.6) \quad \xi \mapsto \lambda \quad := U\xi \quad \in \Omega^0([0, 1], T_{X(0)}M)$$

$$(5.7) \quad e \mapsto \varphi \quad := (U^t)^{-1}e \quad \in \Omega^1([0, 1], T_{X(0)}^*M)$$

$$(5.8) \quad \Pi^\sharp \mapsto P^\sharp \quad := U\Pi^\sharp U^t \quad \in \Omega^0([0, 1], \text{Hom}(T_{X(0)}^*M, T_{X(0)}M)),$$

maps the Equation (5.5) to

$$(5.9) \quad \frac{d\lambda}{dt} = -P^\sharp\varphi.$$

[Cat14, Lemma 2.2] states that if Π is a Poisson bivector, then P is skew symmetric and constant. This implies that the space $C_\Sigma(\lambda, \varphi)$ of solutions of Equation (5.9) can be identified with the Euler Lagrange space $C_\Sigma \subset V(\lambda, \varphi)$ for the constant Poisson structure P^\sharp , thus it is split coisotropic by Proposition 5.11. By using the fact that the space $V(\lambda, \varphi)$ is weak symplectomorphic to $V(X, \eta)$ using the changes of coordinates stated above, the following result holds:

THEOREM 5.12. — *Let $M = \mathbb{R}^n$ and Π be a Poisson bivector on M . Then the linearized Euler–Lagrange space $T_{(X,\eta)}C_\Sigma$ is a split coisotropic subspace of $V(X, \eta)$, for all $(X, \eta) \in C_\Sigma$.*

In [CC15] it is shown that the evolution relations L_Σ^1, L_Σ^2 and L_Σ^3 can be written only in terms of C_Σ and its symplectic orthogonal C_Σ^ω . A similar argument as before proves that

THEOREM 5.13. — *The tangent spaces of the evolution relations TL_Σ^1, TL_Σ^2 and TL_Σ^3 described by*

$$TL_\Sigma^1 = \left\{ (\delta X, \delta \eta) \in TC_\Sigma : \exists \tilde{X} = \text{ct}, \tilde{\eta} \in \ker \Pi^\sharp : (\delta X - \delta \tilde{X}, \delta \eta - \delta \tilde{\eta}) \in (TC_\Sigma)^\omega \right\}$$

$$TL_\Sigma^2 = \left\{ ((\delta X, \delta \eta), (\delta \tilde{X}, \delta \tilde{\eta})) \in TC_\Sigma \oplus TC_\Sigma : (\delta X - \delta \tilde{X}, \delta \eta - \delta \tilde{\eta}) \in (TC_\Sigma)^\omega \right\}$$

$$TL_\Sigma^3 = \left\{ \left((\delta X, \delta \eta), (\delta \tilde{X}, \delta \tilde{\eta}), (\delta \tilde{\tilde{X}}, \delta \tilde{\tilde{\eta}}) \right) \in TC_\Sigma \oplus TC_\Sigma \oplus TC_\Sigma \right. \\ \left. : \left((\delta X, \delta \eta) * (\delta \tilde{X}, \delta \tilde{\eta}), (\delta \tilde{\tilde{X}}, \delta \tilde{\tilde{\eta}}) \right) \in TL_\Sigma^2 \right\},$$

are split Lagrangian subspaces and they intersect neatly with $TC_\Sigma, TC_\Sigma \oplus TC_\Sigma$ and $TC_\Sigma \oplus TC_\Sigma \oplus TC_\Sigma$ respectively, for any Poisson bivector Π .

Remark 5.14. — A possible further applications in field theory of the formalism of split Lagrangian and coisotropic spaces is to consider the cases in which the symplectic structure is strong. In order to achieve that, one has to restrict to the case in which the spaces of fields are Sobolev spaces, and we can describe them as reflexive spaces. A natural example is quantum mechanics and scalar field theory on Riemannian manifolds, in which the space of states is chosen to be l_2 functions, rather

than C^∞ -functions. This viewpoint imposes some conditions over the solutions of the Euler–Lagrange equations, possibly allowing weak solutions to the $EL(M)$ -spaces. In several cases, the spaces of weak and strong solutions coincide, but there are several analytical issues to be considered while understanding the symplectic formulation of weak solutions of PDE.

Remark 5.15. — (Split canonical relations and symplectic groupoids). In [CC15, Con13], the evolution relations for the Poisson sigma model are used to construct an integration of the Poisson manifold M . These relations are the analogues of the graphs of the unit, identity and multiplication maps of a conventional symplectic groupoid $G \rightrightarrows M$ integrating the Lie algebroid T^*M . Theorem 5.13 shows that the relational symplectic groupoid associated to any Poisson manifold M via the Poisson sigma model is split, in the sense that its associated canonical relations L^1, L^2 and L^3 are split.

Remark 5.16. — By using smooth sections of the cotangent bundle of the path space, we consider the Poisson sigma model in terms of weak symplectic Fréchet spaces. However, in the path Sobolev space $L_2^1([0, 1], \mathbb{R})$ of L_2 -differentiable paths, the canonical symplectic form on the cotangent bundle is strong symplectic. The price to pay here is that the Euler–Lagrange space $EL(M)$ must be redefined to allow weak solutions of the O.D.E. given by Equation (5.4).

BIBLIOGRAPHY

- [Arn67] Vladimir I. Arnol'd, *Characteristic class entering in quantization conditions*, Funkts. Anal. Prilozh. **1** (1967), no. 1, 1–14. ↑156
- [Cat14] Alberto S. Cattaneo, *Coisotropic submanifolds and dual pairs*, Lett. Math. Phys. **104** (2014), no. 3, 243–270. ↑173, 182, 183
- [CC15] Alberto S. Cattaneo and Ivan Contreras, *Relational symplectic groupoids*, Lett. Math. Phys. **105** (2015), no. 5, 723–767. ↑157, 171, 172, 183, 184
- [CF00] Alberto S. Cattaneo and Giovanni Felder, *A path integral approach to the Kontsevich quantization formula*, Commun. Math. Phys. **212** (2000), no. 3, 591–611. ↑172
- [CF01] ———, *Poisson sigma models and symplectic groupoids*, Quantization of Singular Symplectic Quotients (Nicolaas P. Landsman, Pflaum Markus J., and Martin Schlichenmaier, eds.), Progress in Mathematics, vol. 198, Springer, 2001, pp. 61–93. ↑171, 172, 177
- [CM14] Alberto S. Cattaneo and Pavel Mněv, *Wave relations*, Commun. Math. Phys. **332** (2014), no. 3, 1083–1111. ↑171
- [CMR12] Alberto S. Cattaneo, Pavel Mněv, and Nikolai Reshetikhin, *Classical and quantum Lagrangian field theories with boundary*, <https://arxiv.org/abs/1207.0239>, 2012. ↑156, 171
- [Con13] Ivan Contreras, *Relational symplectic groupoids and Poisson sigma models with boundary*, Ph.D. thesis, Zürich University, Switzerland, 2013. ↑156, 157, 166, 171, 172, 184
- [EM99] William N. Everitt and Lawrence Markus, *Boundary value problems and symplectic algebra for ordinary and quasi-differential operators*, Mathematical Surveys and Monographs, vol. 61, American Mathematical Society, 1999. ↑169
- [Got82] Mark J. Gotay, *On coisotropic imbeddings of presymplectic manifolds*, Proc. Am. Math. Soc. **84** (1982), 111–114. ↑157

- [Kon03] Maxim Lvovich Kontsevich, *Deformation quantization of Poisson manifolds*, Lett. Math. Phys. **66** (2003), no. 3, 157–216. ↑172
- [KS82] Nigel J. Kalton and R. Charles Swanson, *A symplectic reflexive Banach space with no Lagrangian subspaces*, Trans. Am. Math. Soc. **273** (1982), no. 1, 385–392. ↑160, 170
- [LW15] Jonathan Lorand and Alan D. Weinstein, *Decomposition of (co)isotropic relations*, <https://arxiv.org/abs/1509.04035>, 2015. ↑165
- [Rud91] Walter Rudin, *Functional Analysis*, 2nd ed., International Series in Pure and Applied Mathematics, McGraw-Hill, 1991. ↑158
- [Wei71] Alan D. Weinstein, *Symplectic manifolds and their Lagrangian submanifolds*, Adv. Math. **6** (1971), 329–346. ↑156
- [Wei10] ———, *Symplectic Categories*, Port. Math. (N.S.) **67** (2010), no. 2, 261–278. ↑156, 161

Manuscript received on 28th November 2018,
revised on 3rd March 2020,
accepted on 21st April 2020.

Recommended by Editor A. Weinstein.

Published under license CC BY 4.0.



This journal is a member of Centre Mersenne.



Alberto CATTANEO
Institut für Mathematik,
Universität Zürich, Winterthurerstrasse 190,
CH-8057 Zürich, (Switzerland)
cattaneo@math.uzh.ch

Ivan CONTRERAS
Department of Mathematics,
Amherst College, 31 Quadrangle Drive,
Amherst, MA 01002, (USA)
icontreraspalacios@amherst.edu