

ANNALES  
HENRI LEBESGUE

---

SIMON RICHE

---

GEORDIE WILLIAMSON

---

# A SIMPLE CHARACTER FORMULA

## UNE FORMULE DE CARACTÈRES SIMPLES

---

*Dedicated to Jens Carsten Jantzen,  
on the occasion of his 70<sup>th</sup> birthday.*

**ABSTRACT.** — In this paper we prove a character formula expressing the classes of simple representations in the principal block of a simply-connected semisimple algebraic group  $G$  in terms of baby Verma modules, under the assumption that the characteristic of the base field is bigger than  $2h - 1$ , where  $h$  is the Coxeter number of  $G$ . This provides a replacement for Lusztig's conjecture, valid under a reasonable assumption on the characteristic.

**RÉSUMÉ.** — Dans cet article nous démontrons une formule de caractères exprimant les classes des représentations simples dans le bloc principal d'un groupe algébrique semi-simple  $G$  en termes des modules de Verma bébé, sous l'hypothèse que la caractéristique du corps de base est supérieure à  $2h - 1$ , où  $h$  est le nombre de Coxeter de  $G$ . Ceci fournit un remplacement de la conjecture de Lusztig, valable sous une hypothèse raisonnable sur la caractéristique.

## 1. Introduction

### 1.1. Simple modules for reductive groups

Let  $G$  denote a connected reductive algebraic group over an algebraically closed field  $\mathbb{k}$  of characteristic  $p > 0$ , with simply connected derived subgroup. We fix a

---

*Keywords:* reductive algebraic groups, characters,  $p$ -canonical basis.

2020 *Mathematics Subject Classification:* 20G05.

DOI: <https://doi.org/10.5802/ahl.79>

(\*) This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 677147).

maximal torus and Borel subgroup  $T \subset B \subset G$ . Then for every dominant weight  $\lambda$  we have a Weyl module  $\Delta(\lambda)$  and its simple quotient  $\mathbb{L}(\lambda)$ , both of highest weight  $\lambda$ . We obtain in this way a classification of the simple algebraic  $G$ -modules. A central problem in the field is to compute the characters of these simple modules.

Steinberg’s tensor product theorem reduces this question to the case of  $p$ -restricted highest weights. For a  $p$ -restricted dominant weight  $\lambda$  it is known that  $\mathbb{L}(\lambda)$  stays simple upon restriction to  $G_1$ , the first Frobenius kernel of  $G$ . Moreover, all simple  $G_1$ -modules occur in this way. Thus, understanding the simple  $G$ -modules is equivalent to understanding the simple  $G_1$ -modules.

Instead of working with  $G_1$ -modules, it is technically more convenient to work with  $G_1T$ -modules. Simple  $G_1T$ -modules stay simple upon restriction to  $G_1$ , and thus we can instead try to answer our question in terms of  $G_1T$ -modules. Via Brauer–Humphreys reciprocity, this question can be rephrased in terms of indecomposable projective  $G_1T$ -modules (see Section 1.2 below for details).

In 1980 Lusztig [Lus80a] proposed a conjecture for these characters if  $p$  is not too small (in the guise of “Jantzen’s generic decomposition patterns”). His conjecture is in terms of the canonical basis in the periodic module for the affine Hecke algebra. This formula is known to hold for large  $p$ , see [AJS94, Fie12, KL93a, KL93b, KL94a, KL94b, KT95, KT96, Lus94]; however it is also known to fail for “medium sized”  $p$ , see [Wil17]. In fact, at this point it is not known precisely when this formula holds.

Our goal in this paper is to define the  $p$ -canonical basis in the periodic module and prove that the  $p$ -analogue of Lusztig’s conjecture is true, as long as  $p \geq 2h - 1$  where  $h$  is the Coxeter number of  $G$ . Thus we obtain a character formula for simple  $G$ -modules in terms of  $p$ -Kazhdan–Lusztig polynomials. Our proof builds on a character formula for tilting  $G$ -modules, proved in a joint work with P. Achar and S. Makisumi [AMRW19].<sup>(1)</sup>

At present,  $p$ -canonical bases are very difficult to compute; for this reason our formula is certainly not the final answer to this problem. However it gives a good conceptual understanding of where the difficulties lie, and provides a way to compute characters or multiplicities which is much more efficient than classical techniques in representation theory. For instance, unpublished intensive efforts of Jantzen were not sufficient to completely answer the question of describing simple characters in types  $\mathbf{B}_3$ ,  $\mathbf{C}_3$  and  $\mathbf{A}_4$ . Preliminary results of Jensen–Scheinmann indicate that our formula will allow one to solve these cases, and maybe the case of some bigger groups with the help of a computer. (For example, Jensen and Scheinmann obtain a missing multiplicity in Jantzen’s work for  $\mathbf{A}_4$  in a few lines using our results.) We also believe these results will help answer the important question of when exactly Lusztig’s character formula holds.

*Remark 1.1.* —

- (1) A conjecture of Donkin [Don93] would imply that our formula is valid for  $p > h$ ; see Remark 1.6 below for more details.
- (2) It has been known for a long time that, in theory, knowing the characters of tilting modules is enough to determine the characters of simple modules

---

<sup>(1)</sup>One year after the first version of the present paper was made available, a different proof of this formula was obtained by the authors in [RW20].

(see e.g. [RW18, Section 1.8] for details, and [Sob20] for more recent advances on this question). However, obtaining a *concrete* character formula for simples out of a given character formula for tilting modules is a different story, which is the main topic of the present paper (taking as input the character formula from [AMRW19]).

## 1.2. Representation theory of $G_1T$

We continue with the notation of Section 1.1, and let  $\mathbf{X} := X^*(T)$  be the character lattice of  $T$ . Let also  $B^+$  be the Borel subgroup of  $G$  opposite to  $B$  with respect to  $T$ .

If as above  $G_1T$ , resp.  $B_1^+T$ , denotes the preimage of the Frobenius twist  $\dot{T}$  of  $T$  under the Frobenius morphism of  $G$ , resp.  $B^+$ , then for each  $\lambda \in \mathbf{X}$  we have a  $G_1T$ -module  $\hat{Z}(\lambda)$  (called the *baby Verma module* attached to  $\lambda$ ) obtained by coinducing to  $G_1T$  the 1-dimensional representation of  $B_1^+T$  defined by  $\lambda$  (see Section 5.2 below for details). The module  $\hat{Z}(\lambda)$  has a unique simple quotient  $\hat{\mathbb{L}}(\lambda)$ , and the assignment  $\lambda \mapsto \hat{\mathbb{L}}(\lambda)$  induces a bijection between  $\mathbf{X}$  and the set of isomorphism classes of simple  $G_1T$ -modules. For any  $\lambda, \mu \in \mathbf{X}$  we have

$$\hat{\mathbb{L}}(\lambda + p\mu) \cong \hat{\mathbb{L}}(\lambda) \otimes \mathbb{k}_{\dot{T}}(\mu)$$

(where we identify the weight lattice of  $\dot{T}$  with  $\mathbf{X}$  in such a way that the pullback under the Frobenius morphism  $T \rightarrow \dot{T}$  corresponds to  $\nu \mapsto p\nu$ , and  $\mathbb{k}_{\dot{T}}(\mu)$  is viewed as a  $G_1T$ -module via the Frobenius morphism  $G_1T \rightarrow \dot{T}$ ), and for  $\lambda$  dominant and  $p$ -restricted  $\hat{\mathbb{L}}(\lambda)$  is the restriction to  $G_1T$  of the simple  $G$ -module  $\mathbb{L}(\lambda)$  considered in Section 1.1. In this way, understanding the simple  $G$ -modules, the simple  $G_1$ -modules or the simple  $G_1T$  are equivalent problems.

## 1.3. Characters of $G_1T$ -modules and alcoves

Assume now that  $p \geq h$ .

Let  $\Delta \subset \mathbf{X}$  be the root system of  $(G, T)$ . Let  $W_f$  and  $W = W_f \ltimes \mathbb{Z}\Delta$  denote the finite and affine Weyl groups, and denote their subsets of simple reflections (determined by  $B$ ) by  $S_f$  and  $S$  respectively. The affine Weyl group acts naturally on  $\mathbf{X} \otimes_{\mathbb{Z}} \mathbb{R}$ , giving rise to the set of alcoves  $\mathcal{A}$ . Let  $A_{\text{fund}}$  denote the fundamental alcove; then the map  $x \mapsto x(A_{\text{fund}})$  gives a bijection

$$(1.1) \quad W \xrightarrow{\sim} \mathcal{A}.$$

We will also consider the “dot-action” of  $W$  on  $\mathbf{X}$ , defined by

$$(xt_{\lambda}) \cdot_p \mu = x(\mu + p\lambda + \rho) - \rho$$

for  $x \in W_f$ ,  $\lambda \in \mathbb{Z}\Delta$  and  $\mu \in \mathbf{X}$ , where  $\rho$  is the halfsum of the positive roots. Let  $\text{Rep}_0(G_1T)$  denote the principal block of the category of algebraic finite-dimensional  $G_1T$ -modules, i.e. the Serre subcategory generated by the simple modules  $\hat{\mathbb{L}}(\lambda)$  with  $\lambda \in W \cdot_p 0$ . By construction the highest weights of simple and baby Verma modules

belonging to this block are labelled by the affine Weyl group, and hence (via (1.1)) by alcoves. Given an alcove  $A$ , let us denote the corresponding simple and baby Verma modules by  $\widehat{\mathbb{L}}_A$  and  $\widehat{Z}_A$ . We denote the projective cover (equivalently, injective hull) of  $\widehat{\mathbb{L}}_A$  by  $\widehat{Q}_A$ .

Each  $\widehat{Q}_A$  admits a baby Verma flag (that is, a finite filtration whose successive subquotients are isomorphic to baby Verma modules). We write  $(\widehat{Q}_A : \widehat{Z}_B)$  for the number of times the baby Verma module  $\widehat{Z}_B$  occurs in such a filtration. (This number is known to be independent of the chosen filtration.) For an alcove  $A \in \mathcal{A}$ , consider the element

$$q_A := \sum_{B \in \mathcal{A}} (\widehat{Q}_A : \widehat{Z}_B) B \in \mathbb{Z}[\mathcal{A}].$$

By Brauer–Humphreys reciprocity [Hum71], it is known that

$$(\widehat{Q}_A : \widehat{Z}_B) = [\widehat{Z}_B : \widehat{\mathbb{L}}_A]$$

for any pair of alcoves  $(A, B)$ . As the characters of the baby Verma modules are known (and easy!), knowledge of the elements  $q_A$  for all alcoves  $A$  therefore implies knowledge of the characters of the simple  $G_1T$ -modules in the principal block, and hence of all simple  $G_1T$ -modules by Jantzen’s translation principle.

#### 1.4. Statement

Let  $\mathcal{H}$  denote the Hecke algebra of the Coxeter system  $(W, S)$  (an algebra over the ring  $\mathbb{Z}[v^{\pm 1}]$ , with standard basis  $(H_w : w \in W)$ ), and let  $\mathcal{P}$  denote its periodic (right) module. (We follow the notational conventions of [Soe97].) As a  $\mathbb{Z}[v^{\pm 1}]$ -module,  $\mathcal{P}$  is free with basis given by alcoves:

$$\mathcal{P} = \bigoplus_{A \in \mathcal{A}} \mathbb{Z}[v^{\pm 1}] A.$$

In [Lus80a], Lusztig has defined a canonical basis for  $\mathcal{P}$ ; following the conventions of [Soe97, Theorem 4.3] we will denote this family of elements by  $\{\underline{P}_A : A \in \mathcal{A}\}$ . (Note that this terminology might be misleading:  $\{\underline{P}_A : A \in \mathcal{A}\}$  is not a  $\mathbb{Z}[v^{\pm 1}]$ -basis of  $\mathcal{P}$ , but of a certain submodule.)

Lusztig conjectured (see the last three paragraphs of the introduction to [Lus80a]) that the canonical basis determines the characters of indecomposable projective modules, as follows:

$$q_A = (\underline{P}_{\hat{A}})_{v \mapsto 1}.$$

(Here, given  $R = \sum p_A A \in \mathcal{P}$ ,  $R_{v \mapsto 1} = \sum p_A(1)A \in \mathbb{Z}[\mathcal{A}]$  denotes its specialisation at  $v = 1$ , and  $A \mapsto \hat{A}$  is a simple operation on alcoves, whose definition is recalled in Section 2.) See also [Fie10, Section 3] for the relation with Lusztig’s original conjecture [Lus80b] for characters of the modules  $\mathbb{L}(\lambda)$ .

In this paper, we define the  $p$ -canonical basis  $\{^p\underline{P}_A : A \in \mathcal{A}\}$  in the periodic module  $\mathcal{P}$ , and we prove that it can be used to compute the elements  $q_A$ , as follows.

**THEOREM 1.2.** — *Assume  $p \geq 2h - 1$ . Then for any alcove  $A$  we have*

$$q_A = (^p\underline{P}_{\hat{A}})_{v \mapsto 1}.$$

### 1.5. Spherical and antispherical modules

Theorem 1.2 will be obtained as a consequence of a relation between the  $p$ -canonical bases in two other  $\mathcal{H}$ -modules, namely the (twisted) spherical and the antispherical modules. We will denote by  $(\underline{H}_w : w \in W)$  the Kazhdan–Lusztig basis of  $\mathcal{H}$ , see [Soe97, Theorem 2.1], and by  $({}^p\underline{H}_w : w \in W)$  its  $p$ -canonical basis (see [JW17, RW18]).

The antispherical module  $\mathcal{M}^{\text{asph}}$  is defined as

$$\mathcal{M}^{\text{asph}} = \text{sgn} \otimes_{\mathcal{H}_f} \mathcal{H}$$

where  $\mathcal{H}_f$  is the Hecke algebra of  $(W_f, S_f)$  (which is a subalgebra in  $\mathcal{H}$  in a natural way), and  $\text{sgn}$  is the right  $\mathcal{H}_f$ -module defined as  $\mathbb{Z}[v^{\pm 1}]$  with  $H_s$  acting as multiplication by  $-v$  for  $s \in S_f$ . This module has a standard basis  $(N_w : w \in {}^fW)$  parametrized by the subset  ${}^fW \subset W$  consisting of elements  $w$  which are minimal in  $W_fw$ , where  $N_w := 1 \otimes H_w$ . It also has a Kazhdan–Lusztig basis  $(\underline{N}_w : w \in {}^fW)$  and a  $p$ -canonical basis  $({}^p\underline{N}_w : w \in {}^fW)$ , where

$$\underline{N}_w = 1 \otimes \underline{H}_w, \quad {}^p\underline{N}_w = 1 \otimes {}^p\underline{H}_w.$$

The subset  $\mathcal{A}^+$  of  $\mathcal{A}$  corresponding to  ${}^fW$  under the bijection (1.1) consists of the alcoves contained in the dominant Weyl chamber  $\mathcal{C}^+$ ; hence we will rather parametrize these bases by  $\mathcal{A}^+$  and denote them  $(N_A : A \in \mathcal{A}^+)$ ,  $(\underline{N}_A : A \in \mathcal{A}^+)$  and  $({}^p\underline{N}_A : A \in \mathcal{A}^+)$ .

On the other hand, for  $\lambda \in \mathbf{X}$  we denote by  $\bar{\lambda}$  the unique element in  $\overline{A_{\text{fund}}} \cap (W \cdot \lambda)$ , and let  $W_{\bar{\lambda}} \subset W$  denote its isotropy group. This is a standard parabolic subgroup of  $W$  isomorphic to  $W_f$ ; we denote by  $S_{\bar{\lambda}} := S \cap W_{\bar{\lambda}}$  its subset of simple reflections, and by  $\mathcal{H}_{\bar{\lambda}} \subset \mathcal{H}$  the associated Hecke algebra. Let  $\text{triv}_{\bar{\lambda}}$  denote the “trivial” right  $\mathcal{H}_{\bar{\lambda}}$ -module (defined as  $\mathbb{Z}[v^{\pm 1}]$ , with  $H_s$  acting as multiplication by  $v^{-1}$  for  $s \in S_{\bar{\lambda}}$ ), and set

$$\mathcal{M}_{\bar{\lambda}}^{\text{sph}} := \text{triv}_{\bar{\lambda}} \otimes_{\mathcal{H}_{\bar{\lambda}}} \mathcal{H}.$$

Then  $\mathcal{M}_{\bar{\lambda}}^{\text{sph}}$  has a standard basis

$$(M_w^{\bar{\lambda}} := 1 \otimes H_w : w \in \bar{\lambda}W)$$

parametrized by the subset  $\bar{\lambda}W \subset W$  consisting of elements  $w$  which are minimal in  $W_{\bar{\lambda}}w$ .

If  $w_{\bar{\lambda}}$  is the longest element in  $W_{\bar{\lambda}}$ , then the map  $1 \otimes h \mapsto \underline{H}_{w_{\bar{\lambda}}} \cdot h$  provides an embedding

$$(1.2) \quad \zeta_{\bar{\lambda}} : \mathcal{M}_{\bar{\lambda}}^{\text{sph}} \hookrightarrow \mathcal{H}.$$

For any  $w \in \bar{\lambda}W$ , the element  $\underline{H}_{w_{\bar{\lambda}}w}$ , resp.  ${}^p\underline{H}_{w_{\bar{\lambda}}w}$ , belongs to the image of  $\zeta_{\bar{\lambda}}$ , and if we denote by  $\underline{M}_w^{\bar{\lambda}}$ , resp.  ${}^p\underline{M}_w^{\bar{\lambda}}$ , its preimage in  $\mathcal{M}_{\bar{\lambda}}^{\text{sph}}$ , then

$$(\underline{M}_w^{\bar{\lambda}} : w \in \bar{\lambda}W) \quad \text{and} \quad ({}^p\underline{M}_w^{\bar{\lambda}} : w \in \bar{\lambda}W)$$

are bases of  $\mathcal{M}_{\bar{\lambda}}^{\text{sph}}$ , called the Kazhdan–Lusztig and the  $p$ -canonical basis respectively.

Recall that the set  $\mathcal{A}$  also admits a natural *right* action of  $W$ ; through the identification (1.1) this action corresponds to the right multiplication of  $W$  on itself, and will be denoted  $(A, x) \mapsto A \cdot x$ . Then the assignment  $w \mapsto (\lambda + A_{\text{fund}}) \cdot w$  identifies  ${}^{\bar{\lambda}}W$  with the subset  $\mathcal{A}_{\bar{\lambda}}^+ \subset \mathcal{A}$  consisting of alcoves contained in  $\lambda + \mathcal{C}^+$ . In this way the bases of  $\mathcal{M}_{\bar{\lambda}}^{\text{sph}}$  considered above can be labelled by  $\mathcal{A}_{\bar{\lambda}}^+$ , and will be denoted  $(M_A^{\bar{\lambda}} : A \in \mathcal{A}_{\bar{\lambda}}^+)$  and  $({}^p M_A^{\bar{\lambda}} : A \in \mathcal{A}_{\bar{\lambda}}^+)$ . (Note that this labelling depends on  $\lambda$ , not only on  $\bar{\lambda}$ .)

*Remark 1.3.* — The bases  $(\underline{N}_w : w \in {}^f W)$  and  $(\underline{M}_w^{\bar{\lambda}} : w \in {}^{\bar{\lambda}}W)$  coincide with the bases considered in [Soe97, Theorem 3.1] (for the parabolic subgroups  $W_{\mathbf{f}}$  and  $W_{\bar{\lambda}}$  respectively), see [Soe97, Proof of Proposition 3.4]. In case  $\lambda = \bar{\lambda} = 0$ , we will write  $\mathcal{M}^{\text{sph}}$  for  $\mathcal{M}_0^{\text{sph}}$ ,  $M_w$  for  $M_w^0$ ,  $\underline{M}_w$  for  $\underline{M}_w^0$ , and  ${}^p \underline{M}_w$  for  ${}^p \underline{M}_w^0$ .

### 1.6. Outline of the proof of Theorem 1.2

Let  $\lambda \in \mathbf{X}$  be such that  $\lambda + x(A_{\text{fund}})$  belongs to  $\mathcal{C}^+$  for all  $x \in W_{\mathbf{f}}$ . It is easily seen that the assignment  $1 \otimes h \mapsto \underline{N}_{\lambda + A_{\text{fund}}} \cdot h$  induces an  $\mathcal{H}$ -module morphism

$$\varphi_{\lambda} : \mathcal{M}_{\bar{\lambda}}^{\text{sph}} \rightarrow \mathcal{M}^{\text{asph}}.$$

Now we assume (for simplicity) that  $G$  is semisimple, and specialize to the case  $\lambda = \rho$ . The key step in our approach to Theorem 1.2 is the following claim.

**THEOREM 1.4.** — *Assume that  $p$  is good for  $G$ . Then for any  $A \in \mathcal{A}_{\rho}^+$  we have*

$$\varphi_{\rho}({}^p \underline{M}_A^{\rho}) = {}^p \underline{N}_A.$$

*Remark 1.5.* —

- (1) In the body of the paper, we find it more convenient to work with the *extended* affine Hecke algebra and its spherical/antispherical modules. This allows us to remove some of the twistings above.
- (2) Our proof of Theorem 1.4 also applies to the Kazhdan–Lusztig bases, and shows that for any  $A \in \mathcal{A}_{\rho}^+$  we have  $\varphi_{\rho}(\underline{M}_A^{\rho}) = \underline{N}_A$ . This fact could have been stated many years ago (since it does not involve the  $p$ -canonical bases in any way), but seems to be new. A direct combinatorial proof of this formula (explained to us by Wolfgang Soergel) is provided in Section 6.

Theorem 1.4 is obtained as a “combinatorial trace” of a statement of categorical nature. Namely, the modules  $\mathcal{M}_{\bar{\rho}}^{\text{sph}}$  and  $\mathcal{M}^{\text{asph}}$  can be “categorified” via some categories of parity complexes (in the sense of [JMW14]) on the affine flag variety of the Langlands dual group  $G^{\vee}$ :  $\mathcal{M}_{\bar{\rho}}^{\text{sph}}$  corresponds to (twisted) *spherical* parity complexes, while  $\mathcal{M}^{\text{asph}}$  corresponds to *Iwahori–Whittaker*<sup>(2)</sup> parity complexes. The morphism  $\varphi_{\rho}$  can also be categorified by a functor  $\Phi_{\rho}$ , given by convolution with a

<sup>(2)</sup>This construction is a “finite-dimensional” and geometric counterpart for the classical (and very useful!) Whittaker constructions in representation theory of  $p$ -adic groups; see in particular [BBM04, BGM<sup>+</sup>19, BY13].

certain object. We then prove that this functor is full (but *not* faithful); in particular, as both categories involved are Krull-Schmidt, it must send indecomposable objects to indecomposable objects, and Theorem 1.4 follows. This fullness result is deduced from a similar result in the case where the affine flag variety is replaced by the affine Grassmannian of  $G^\vee$  (in which case the corresponding functor is even an equivalence of categories) proved recently by the first author with R. Bezrukavnikov, D. Gaitsgory, I. Mirković and L. Rider, see [BGM<sup>+</sup>19].


Once Theorem 1.4 is proved, we deduce Theorem 1.2 from the fact that the  $p$ -canonical basis of  $\mathcal{M}^{\text{asph}}$  encodes the characters of indecomposable tilting  $G$ -modules (as proved in joint work with P. Achar and S. Makisumi [AMRW19]) and a result of Jantzen [Jan80] (in the interpretation of Donkin [Don93]) saying that if the highest weight of an indecomposable tilting module is of the form  $(p-1)\rho + \mu$  with  $\mu$  dominant and  $p$ -restricted, then this tilting module is indecomposable as a  $G_1T$ -module. It is well known (and easy to check) that these modules are projective over  $G_1T$ ; this therefore provides a useful relation between indecomposable tilting modules for  $G$  and indecomposable projective modules for  $G_1T$ , which allows us to deduce Theorem 1.2 from Theorem 1.4 and the results of [AMRW19].

*Remark 1.6.* — A conjecture by Donkin states that Jantzen’s result recalled above should be true in any characteristic, which would imply that Theorem 1.2 holds as soon as  $p > h$ . Very recent work of Bendel–Nakano–Pillen–Sobaje [BNPS20] shows that this conjecture is not true in full generality. It is currently not known whether the condition that  $p > h$  is sufficient to ensure that this conjecture holds (which would be sufficient for our purposes).

### 1.7. Computational complexity

Let us return briefly to our earlier claim that our formula is potentially useful in practice. In all known algorithms to compute a  $p$ -canonical basis element  ${}^p\overline{H}_w$ ,  ${}^p\overline{N}_w$ ,  ${}^p\overline{M}_w$  etc. the computational complexity is exponential in the length of  $w$ . For a character formula involving these data to be computable in practice, it must therefore only involve elements whose length is small (or at least as small as possible). In the following, we compare the lengths involved in the computation based directly on the approach of Andersen [And98] (see also [Sob20]) and our formula.


We use  $G = \text{Sp}_4$  as a running example.<sup>(3)</sup> Here are the first few dominant alcoves:



<sup>(3)</sup>This choice is made only so that we can draw pictures. The case of  $G = \text{Sp}_4$  can easily be settled by classical techniques, e.g. the Jantzen sum formula.

By Steinberg's tensor product theorem, it is enough to know the characters of the simple modules corresponding to the shaded alcoves.

As we explained above, Brauer–Humphreys reciprocity combined with a result of Donkin (assuming  $p \geq 2h - 2$ ) allows us to rephrase this question in terms of the characters of indecomposable tilting modules indexed by the following shaded alcoves:




The main theorem of [AMRW19] (see also [RW20]) asserts that the character of the indecomposable tilting module attached to an alcove  $A \in \mathcal{A}^+$  is calculated by the element  ${}^pN_w$  for  $w \in W$  such that  $w(A_{\text{fund}}) = A$ .

This looks innocent enough in this example, however for a general group the smallest and largest alcoves in this set (marked  $A$  and  $D$  above) correspond to elements of  ${}^fW$  of lengths

$$(1.3) \quad 2\langle \rho^\vee, \rho \rangle = \sum_{\alpha \in \Delta^+} \text{ht}(\alpha) \quad \text{and} \quad 4\langle \rho^\vee, \rho \rangle - \ell(w_f) = 2 \left( \sum_{\alpha \in \Delta^+} \text{ht}(\alpha) \right) - \ell(w_f).$$

(Here  $\Delta^+ \subset \Delta$  denotes the subset of positive roots,  $\text{ht}$  denotes height,  $\rho^\vee$  denotes the halfsum of the positive coroots and  $w_f$  denotes the longest element of  $W_f$ . This formula is easily deduced from standard formulas for the length function in affine Weyl groups, see [IM65, Proposition 1.23].) These numbers grow fast in the rank of the group, and soon exceed current computational capability.


The main technical result of this paper (Theorem 1.4) says that the span of the  $p$ -canonical basis elements indexed by the following shaded alcoves



in the anti-spherical module  $\mathcal{M}^{\text{asph}}$  yield a submodule isomorphic to the (twisted) spherical module  $\mathcal{M}_\rho^{\text{sph}}$ , and moreover that this inclusion preserves the  $p$ -canonical



basis. In other words, for the tilting characters that concern us above, it is enough to calculate the  $p$ -canonical basis elements for the following shaded alcoves in the (twisted) spherical module:



This is computationally a much simpler prospect, because the elements involved are shorter; the elements we have to consider range from length 0 to

$$(1.4) \quad 2\langle \rho^\vee, \rho \rangle - \ell(w_f) = \left( \sum_{\alpha \in \Delta^+} \text{ht}(\alpha) \right) - \ell(w_f).$$

(That is, we have improved (1.3) by a constant factor of  $2\langle \rho^\vee, \rho \rangle$ .)

### Acknowledgements

Part of the work on this paper was done while the first author was a member of the Freiburg Institute for Advanced Studies, as part of the Research Focus “Cohomology in Algebraic Geometry and Representation Theory” led by A. Huber–Klawitter, S. Kebekus and W. Soergel. We thank H. H. Andersen, G. Lusztig and the referees for their helpful suggestions. A special thanks to W. Soergel for providing the proof explained in Section 6.

## 2. The periodic module

### 2.1. Extended affine Weyl group

As in Section 1.2 we consider a connected reductive algebraic group  $G$  with simply-connected derived subgroup over an algebraically closed field  $\mathbb{k}$  of characteristic  $p > 0$ , with a fixed choice of maximal torus and Borel subgroup  $T \subset B \subset G$ . We set  $\mathbf{X} := X^*(T)$ , and denote by  $\Delta \subset \mathbf{X}$  the root system of  $(G, T)$ , by  $\Delta^+ \subset \Delta$  the positive system consisting of the opposites of the  $T$ -weights in the Lie algebra of  $B$ , by  $\Sigma \subset \Delta$  the corresponding subset of simple roots, by  $W_f$  and  $W = W_f \ltimes \mathbb{Z}\Delta$  the Weyl group and the affine Weyl group, and finally by  $S_f$  and  $S$  their subsets of simple roots, see e.g. [Jan03, Chapter 6]. (Here, “f” stands for “finite.”)

By a result of Iwahori–Matsumoto [IM65], the length function associated with the Coxeter system  $(W, S)$  satisfies

$$(2.1) \quad \ell(w \cdot t_\lambda) = \sum_{\substack{\alpha \in \Delta^+ \\ w(\alpha) \in \Delta^+}} |\langle \lambda, \alpha^\vee \rangle| + \sum_{\substack{\alpha \in \Delta^+ \\ w(\alpha) \in -\Delta^+}} |1 + \langle \lambda, \alpha^\vee \rangle|$$

for  $w \in W_f$  and  $\lambda \in \mathbb{Z}\Delta$ . (Here, to avoid confusion, the image of  $\lambda \in \mathbb{Z}\Delta$  in  $W$  is denoted  $t_\lambda$ .) We will also consider the *extended* affine Weyl group

$$W_{\text{ext}} := W_f \ltimes \mathbf{X}.$$

The formula (2.1) defines a function  $\ell : W_{\text{ext}} \rightarrow \mathbb{Z}$ , and we set  $\Omega := \{w \in W_{\text{ext}} \mid \ell(w) = 0\}$ . Then it is known that  $\Omega$  is a subgroup of  $W_{\text{ext}}$ , and that multiplication induces a group isomorphism

$$\Omega \ltimes W \xrightarrow{\sim} W_{\text{ext}}.$$

More precisely, conjugation by any element of  $\Omega$  defines a Coxeter group automorphism of  $W$ ; in other words it stabilizes  $S$  (hence preserves lengths). It is known also that the composition

$$(2.2) \quad \Omega \hookrightarrow W_{\text{ext}} = W \ltimes \mathbf{X} \rightarrow \mathbf{X} \rightarrow \mathbf{X}/\mathbb{Z}\Delta$$

is a group isomorphism. (In particular,  $\Omega$  is abelian.)

We consider the action of  $W$  and  $W_{\text{ext}}$  on  $V := \mathbb{R} \otimes_{\mathbb{Z}} \mathbf{X}$  as in [Soe97, Section 4], and denote by  $\mathcal{A}$  the set of alcoves in  $V$  (i.e. the connected components of the complement of the union of the reflection hyperplanes associated with the action of  $W$ ) and by  $A_{\text{fund}} \in \mathcal{A}$  the fundamental alcove, defined as

$$A_{\text{fund}} = \left\{ v \in V \mid \forall \alpha \in \Delta^+, 0 < \langle v, \alpha^\vee \rangle < 1 \right\}.$$

The action of  $W_{\text{ext}}$  on  $V$  preserves alcoves, hence induces an action on  $\mathcal{A}$ . Moreover the assignment  $w \mapsto wA_{\text{fund}}$  induces a bijection  $W \xrightarrow{\sim} \mathcal{A}$ , see (1.1). If we denote by  $\mathcal{A}^+ \subset \mathcal{A}$  the subset of alcoves contained in the dominant chamber (denoted  $\mathcal{C}$  in [Soe97]), then this bijection restricts to a bijection

$$(2.3) \quad {}^fW \xrightarrow{\sim} \mathcal{A}^+$$

where  ${}^fW$  is as in Section 1.5.

*Remark 2.1.* — The subset  $\Omega \subset W_{\text{ext}}$  can be characterized as consisting of the elements  $w \in W_{\text{ext}}$  such that  $w(A_{\text{fund}}) = A_{\text{fund}}$ . For any  $\lambda \in \mathbf{X}$ , the subset  $\lambda + A_{\text{fund}}$  is an alcove, so that there exists  $x_\lambda \in W$  such that  $x_\lambda(A_{\text{fund}}) = \lambda + A_{\text{fund}}$ , see (1.1). Then  $\omega_\lambda = (x_\lambda)^{-1} \cdot t_\lambda$  belongs to  $\Omega$ , and has image  $\lambda + \mathbb{Z}\Delta$  under (2.2). This procedure realizes the isomorphism inverse to (2.2). It also allows us to associate to any  $\lambda \in \mathbf{X}$  a Coxeter group automorphism  $\tau_\lambda$  of  $W$ , given by conjugation by  $\omega_\lambda$  in  $W_{\text{ext}}$ . With this notation, the subset  $S_{\bar{\lambda}}$  of Section 1.5 is given by  $S_{\bar{\lambda}} = \tau_\lambda(S_f)$ .

## 2.2. More about alcoves

Using again the identification (1.1) we may transport the Bruhat order on  $W$  to obtain a partial order on  $\mathcal{A}$ , which we also denote by  $\leq$ . We define the *generic order* on  $\mathcal{A}$  by

$$A \preceq B \text{ if } A + m\gamma \leq B + m\gamma \text{ for all strictly dominant } \gamma \text{ and } m \gg 0.$$

Equivalently, the generic order is uniquely determined by the fact that it agrees with the Bruhat order on  $\mathcal{A}^+$  and is invariant under translation by  $\mathbf{X}$ .

For any  $\mu \in \mathbf{X}$  we will consider the following subsets of  $V$ :

$$\begin{aligned}\check{\Pi}_\mu &:= \{v \in V \mid \langle \mu, \alpha^\vee \rangle - 1 < \langle v, \alpha^\vee \rangle \leq \langle \mu, \alpha^\vee \rangle \text{ for all } \alpha \in \Sigma\}, \\ \hat{\Pi}_\mu &:= \{v \in V \mid \langle \mu, \alpha^\vee \rangle \leq \langle v, \alpha^\vee \rangle < \langle \mu, \alpha^\vee \rangle + 1 \text{ for all } \alpha \in \Sigma\}.\end{aligned}$$

Note that for  $\mu, \nu \in \mathbf{X}$  we have

$$(2.4) \quad \check{\Pi}_\mu = \check{\Pi}_\nu \Leftrightarrow \hat{\Pi}_\mu = \hat{\Pi}_\nu \Leftrightarrow (\forall \alpha \in \Delta, \langle \mu - \nu, \alpha^\vee \rangle = 0).$$

The set  $\hat{\Pi}_0$  is often called the “fundamental box.” (It is denoted  $\Pi$  in [Soe97, Section 4].) The notation is intended to suggest that  $\check{\Pi}_\mu$  (resp.  $\hat{\Pi}_\mu$ ) is “the fundamental box below (resp. above)  $\mu$ .”

Given any alcove  $A \in \mathcal{A}$  there exists  $\mu \in \mathbf{X}$  such that  $A \subset \check{\Pi}_\mu$ . The stabiliser of  $\mu$  in  $W$  is then  $t_\mu W_f t_{-\mu}$ . We define

$$\hat{A} := (t_\mu w_f t_{-\mu})(A),$$

where  $w_f$  is the longest element in  $W_f$ . From (2.4) we see that  $\hat{A}$  does not depend on the choice of  $\mu$ , that  $\hat{A} \subset \hat{\Pi}_\mu$ , and that  $A \mapsto \hat{A}$  is a bijection  $\mathcal{A} \xrightarrow{\sim} \mathcal{A}$ . We denote its inverse by  $A \mapsto \check{A}$ . (These operations agree with the maps denoted similarly in [Soe97], see [Soe97, Comments before Lemma 4.21].)

### 2.3. The periodic module and its canonical basis

The *periodic module*  $\mathcal{P}$  is the right  $\mathcal{H}$ -module (where, as in Section 1,  $\mathcal{H}$  is the Hecke algebra of the Coxeter system  $(W, S)$ ) defined as follows. As a  $\mathbb{Z}[v^{\pm 1}]$ -module,  $\mathcal{P}$  is free with basis the set of alcoves:

$$\mathcal{P} := \bigoplus_{A \in \mathcal{A}} \mathbb{Z}[v^{\pm 1}]A.$$

The structure of a right  $\mathcal{H}$ -module on  $\mathcal{P}$  is characterized by the following formulas for  $s \in S$ :

$$(2.5) \quad A \cdot \underline{H}_s = \begin{cases} As + vA & \text{if } A \preccurlyeq As; \\ As + v^{-1}A & \text{if } As \preccurlyeq A, \end{cases}$$

see [Soe97, Lemma 4.1]. (Here,  $\underline{H}_s = H_s + v$ ).

The action of  $\mathbf{X}$  on  $\mathcal{A}$  by translations extends to an action of  $\mathbf{X}$  on  $\mathcal{P}$ : namely, given  $R = \sum p_A A \in \mathcal{P}$ , set  $R + \mu := \sum p_A (\mu + A)$ . This action does not commute with the  $\mathcal{H}$ -action; it rather satisfies the following relation for all  $h \in \mathcal{H}$ :

$$(2.6) \quad (R \cdot h) + \mu = (R + \mu) \cdot \tau_\mu(h),$$

where we still denote by  $\tau_\mu$  the automorphism of  $\mathcal{H}$  defined by  $\tau_\mu(H_w) = H_{\tau_\mu(w)}$ .

As in Section 1.4 we denote by  $\{\underline{P}_A : A \in \mathcal{A}\}$  the canonical basis of  $\mathcal{P}$ . Then for any  $\mu \in \mathbf{X}$  we have

$$(2.7) \quad \underline{P}_{A+\mu} = \underline{P}_A + \mu$$

(see [Soe97, Comments before Proposition 4.18]) and

$$(2.8) \quad \underline{P}_{A_{\text{fund}}+\mu} = \sum_{x \in W_{\text{f}}} v^{\ell(x)} \cdot (x(A_{\text{fund}}) + \mu)$$

(see [Soe97, Proof of Proposition 4.16]).

We now recall a crucial observation of Lusztig. Recall the embedding  $\zeta_{\bar{\mu}} : \mathcal{M}_{\bar{\mu}}^{\text{sph}} \rightarrow \mathcal{H}$  from (1.2). By [Lus80a, Theorem 5.2] we then have the following formula, where we set  $\pi_{\text{f}} := v^{-\ell(w_{\text{f}})} \sum_{x \in W_{\text{f}}} v^{\ell(x)}$ .

**LEMMA 2.2.** — *Let  $A \in \mathcal{A}$ , let  $\mu \in \mathbf{X}$  be such that  $A \subset \hat{\Pi}_{\mu}$ , and let  $w \in W$  be the unique element such that  $(\mu + A_{\text{fund}}) \cdot w = A$ . Then we have*

$$\underline{P}_A = \frac{1}{\pi_{\text{f}}} \underline{P}_{A_{\text{fund}}+\mu} \cdot \zeta_{\bar{\mu}}(\underline{M}_{\bar{w}}^{\bar{\mu}}).$$

*Remark 2.3.* — The formula in Lemma 2.2 is compatible with (2.7) in view of (2.6).

## 2.4. The $p$ -canonical basis of the periodic module

The usual procedure to define a  $p$ -canonical basis of an  $\mathcal{H}$ -module (see [AR19, JW17, RW18]) is to start with a categorification of this module in terms of a  $\mathbb{C}$ -linear category (in practice, either via some diagrammatic category or some category of parity complexes) such that the classes of indecomposable objects correspond to the Kazhdan–Lusztig basis, and then to replace (in some appropriate way) the coefficients  $\mathbb{C}$  by a field of characteristic  $p$ . In the case of the periodic module, the known categorifications involve semi-infinite geometry, and are beyond the authors' present understanding of the subject. So we will use a different strategy to define this basis: we will start with the formula of Lemma 2.2, and replace there the canonical basis of  $\mathcal{M}_{\bar{\mu}}^{\text{sph}}$  by the  $p$ -canonical version. We expect that any reasonable categorification of  $\mathcal{P}$  with characteristic- $p$  coefficients should provide the same basis as the one constructed here.

Namely, for  $A \in \mathcal{A}$ , we choose  $\mu \in \mathbf{X}$  such that  $A \subset \hat{\Pi}_{\mu}$ , and let  $w \in W$  be the unique element such that  $(\mu + A_{\text{fund}}) \cdot w = A$ . Then we set

$${}^p\underline{P}_A := \frac{1}{\pi_{\text{f}}} \underline{P}_{A_{\text{fund}}+\mu} \cdot \zeta_{\bar{\mu}}({}^p\underline{M}_{\bar{w}}^{\bar{\mu}}).$$

The following properties are easy to check (using in particular (2.6) and the fact that  $\tau_{\mu} = \tau_{\nu}$  if  $\hat{\Pi}_{\mu} = \hat{\Pi}_{\nu}$ , as follows from (2.4)):

- (1) for any  $h \in \mathcal{M}_{\bar{\mu}}^{\text{sph}}$  the element  $\underline{P}_{A_{\text{fund}}+\mu} \cdot \zeta_{\bar{\mu}}(h)$  belongs to  $\pi_{\text{f}} \cdot \mathcal{P}$ , so that  ${}^p\underline{P}_A$  belongs to  $\mathcal{P}$ ;
- (2) the element  ${}^p\underline{P}_A$  does not depend on the choice of  $\mu$ ;
- (3) for any  $\nu \in \mathbf{X}$  we have

$$(2.9) \quad {}^p\underline{P}_{A+\nu} = {}^p\underline{P}_A + \nu.$$

It can also be shown (although this is less obvious, and will not be proved here) that for any alcove  $A$  we have

$${}^p P_A \in \sum_{B \in \mathcal{A}} \mathbb{Z}_{\geq 0} [v^{\pm 1}] \cdot P_B.$$

### 3. The extended affine Hecke algebra and its spherical and antispherical modules

#### 3.1. The spherical and antispherical modules

We continue with the notation of Section 2, and fix a weight  $\varsigma \in \mathbf{X}$  such that  $\langle \varsigma, \alpha^\vee \rangle = 1$  for any  $\alpha \in \Sigma$ . (Such a weight exists thanks to our assumption on the derived subgroup of  $G$ . However, it might not be unique.)

As mentioned in Remark 1.5, to avoid difficulties related to the twists  $\tau_\lambda$ , it will be more convenient to work with the Hecke algebra  $\mathcal{H}_{\text{ext}}$  associated with the “quasi-Coxeter” group  $W_{\text{ext}}$  (see Section 2.1), i.e. the  $\mathbb{Z}[v^{\pm 1}]$ -algebra with a “standard” basis consisting of elements  $(H_w : w \in W_{\text{ext}})$ , with multiplication characterized by the following relations:

- (1)  $(H_s + v) \cdot (H_s - v^{-1}) = 0$  for  $s \in S$ ;
- (2)  $H_x \cdot H_y = H_{xy}$  if  $x, y \in W_{\text{ext}}$  and  $\ell(xy) = \ell(x) + \ell(y)$ .

The algebra  $\mathcal{H}_{\text{ext}}$  contains  $\mathcal{H}$  as a subalgebra (spanned by the elements  $H_w$  with  $w \in W$ ). Inducing from  $\mathcal{H}_f$  to  $\mathcal{H}_{\text{ext}}$  the modules considered in Section 1.5, we obtain the right  $\mathcal{H}_{\text{ext}}$ -modules

$$\mathcal{M}_{\text{ext}}^{\text{sph}} := \text{triv}_0 \otimes_{\mathcal{H}_f} \mathcal{H}_{\text{ext}} \quad \text{and} \quad \mathcal{M}_{\text{ext}}^{\text{asph}} := \text{sgn} \otimes_{\mathcal{H}_f} \mathcal{H}_{\text{ext}},$$

which are called the *spherical* and *antispherical* module respectively.

We denote by  ${}^f W_{\text{ext}} \subset W_{\text{ext}}$  the subset consisting of elements  $w$  which are of minimal length in the coset  $W_f w$  (in other words, of the form  $w\omega$  with  $w \in {}^f W$  and  $\omega \in \Omega$ ). Then for  $w \in {}^f W_{\text{ext}}$  we set

$$M_w := 1 \otimes H_w \in \mathcal{M}_{\text{ext}}^{\text{sph}}, \quad N_w := 1 \otimes H_w \in \mathcal{M}_{\text{ext}}^{\text{asph}}.$$

The collections  $(M_w : w \in {}^f W_{\text{ext}})$  and  $(N_w : w \in {}^f W_{\text{ext}})$  are  $\mathbb{Z}[v^{\pm 1}]$ -bases of  $\mathcal{M}^{\text{sph}}$  and  $\mathcal{M}^{\text{asph}}$  respectively (called again the *standard* bases). Of course there are natural embeddings  $\mathcal{M}^{\text{sph}} \hookrightarrow \mathcal{M}_{\text{ext}}^{\text{sph}}$  and  $\mathcal{M}^{\text{asph}} \hookrightarrow \mathcal{M}_{\text{ext}}^{\text{asph}}$ , such that the elements denoted  $M_w$ , resp.  $N_w$ , in Section 1.5 (see Remark 1.3) correspond to the elements denoted similarly here.

*Remark 3.1.* — Let  $\lambda \in \mathbf{X}$ , and consider the associated element  $\omega_\lambda \in \Omega$ , see Remark 2.1. Then the map  $h \mapsto H_{\omega_\lambda^{-1}} \cdot h$  induces an embedding of right  $\mathcal{H}$ -modules  $\mathcal{M}_\lambda^{\text{sph}} \hookrightarrow \mathcal{M}_{\text{ext}}^{\text{sph}}$ . In this way, one can consider  $\mathcal{M}_{\text{ext}}^{\text{sph}}$  as the result of “gluing together” all the modules  $\mathcal{M}_\lambda^{\text{sph}}$  from the introduction.

### 3.2. Kazhdan–Lusztig and $p$ -canonical bases

Via the natural embeddings  $\mathcal{H} \hookrightarrow \mathcal{H}_{\text{ext}}$ ,  $\mathcal{M}^{\text{sph}} \hookrightarrow \mathcal{M}_{\text{ext}}^{\text{sph}}$  and  $\mathcal{M}^{\text{asph}} \hookrightarrow \mathcal{M}_{\text{ext}}^{\text{asph}}$ , the Kazhdan–Lusztig and  $p$ -canonical bases of  $\mathcal{H}$ ,  $\mathcal{M}^{\text{sph}}$  and  $\mathcal{M}^{\text{asph}}$  define families of elements in  $\mathcal{H}_{\text{ext}}$ ,  $\mathcal{M}_{\text{ext}}^{\text{sph}}$  and  $\mathcal{M}_{\text{ext}}^{\text{asph}}$ . We complete these families into bases by setting, for  $w \in W$  and  $\omega \in \Omega$ ,

$$\begin{aligned} \underline{H}_{w\omega} &:= \underline{H}_w \cdot H_\omega, & {}^p\underline{H}_{w\omega} &:= {}^p\underline{H}_w \cdot H_\omega, \\ \underline{M}_{w\omega} &:= \underline{M}_w \cdot H_\omega, & {}^p\underline{M}_{w\omega} &:= {}^p\underline{M}_w \cdot H_\omega, \\ \underline{N}_{w\omega} &:= \underline{N}_w \cdot H_\omega, & {}^p\underline{N}_{w\omega} &:= {}^p\underline{N}_w \cdot H_\omega. \end{aligned}$$

(It can be easily checked that we also have  $\underline{H}_{\omega w} = H_\omega \underline{H}_w$  and  ${}^p\underline{H}_{\omega w} = H_\omega {}^p\underline{H}_w$  for any  $\omega \in \Omega$  and  $w \in W$ .)

Let us recall the following well-known property of the Kazhdan–Lusztig basis.

LEMMA 3.2. — *Let  $w \in W_{\text{ext}}$  and  $s \in S$ . If  $\ell(ws) < \ell(w)$ , then*

$$\underline{H}_w \cdot \underline{H}_s = (v + v^{-1}) \cdot \underline{H}_w.$$

As in the setting of Section 1.5 we have an  $\mathcal{H}_{\text{ext}}$ -module morphism

$$\xi : \mathcal{H}_{\text{ext}} \rightarrow \mathcal{M}_{\text{ext}}^{\text{asph}}$$

defined by  $\xi(h) = N_{\text{id}} \cdot h$ . This morphism is clearly surjective; moreover, for  $w \in W_{\text{ext}}$  we have

$$\xi(\underline{H}_w) = \begin{cases} \underline{N}_w & \text{if } w \in {}^fW_{\text{ext}}; \\ 0 & \text{otherwise,} \end{cases} \quad \xi({}^p\underline{H}_w) = \begin{cases} {}^p\underline{N}_w & \text{if } w \in {}^fW_{\text{ext}}; \\ 0 & \text{otherwise} \end{cases}$$

(see [RW18] for details).

Now, let  $w_f$  be the longest element in  $W_f$ . Consider the endomorphism of  $\mathcal{H}_{\text{ext}}$  (as a right  $\mathcal{H}_{\text{ext}}$ -module) sending  $h$  to  $\underline{H}_{w_f} \cdot h$ . It follows from Lemma 3.2 that this morphism factors through a morphism

$$(3.1) \quad \zeta : \mathcal{M}_{\text{ext}}^{\text{sph}} \rightarrow \mathcal{H}_{\text{ext}}.$$

This morphism is injective, and satisfies

$$(3.2) \quad \zeta(\underline{M}_w) = \underline{H}_{w_f w}, \quad \zeta({}^p\underline{M}_w) = {}^p\underline{H}_{w_f w}$$

for any  $w \in {}^fW_{\text{ext}}$ . In particular, for  $\omega \in \Omega$  we have

$$(3.3) \quad \zeta(M_\omega) = \underline{H}_{w_f} \cdot H_\omega = \sum_{z \in W_f} v^{\ell(w_f) - \ell(z)} H_{z\omega}$$

(where the second equality uses [Soe97, Proposition 2.9]).

*Remark 3.3.* — Recall that for any  $w \in W_{\text{ext}}$ , there exists  $K_w \in \mathbb{Z}_{\geq 0}$  such that  ${}^p\underline{H}_w = \underline{H}_w$  for any prime number  $p$  such that  $p \geq K_w$ . (However, determining  $K_w$  is a very difficult task.) A similar claim holds for the  $p$ -canonical bases in  $\mathcal{M}_{\text{ext}}^{\text{asph}}$  and  $\mathcal{M}_{\text{ext}}^{\text{sph}}$ .

### 3.3. Statement

We can now state a version of Theorem 1.4 in terms of the *extended* affine Hecke algebra (and for reductive groups which are not necessarily semisimple).

The statement will involve the element considered in the following lemma. (Here the second equality follows from [Soe97, Lemma 5.7]. We will give a (geometric) proof of both equalities in Section 4.2 below.)

LEMMA 3.4. — We have

$$(3.4) \quad {}^p\underline{N}_{t_\varsigma} = \underline{N}_{t_\varsigma} = \sum_{z \in W_f} v^{\ell(z)} N_{t_\varsigma \cdot z}.$$

Moreover, for any  $s \in S_f$  we have

$$\underline{N}_{t_\varsigma} \cdot \underline{H}_s = (v + v^{-1}) \cdot \underline{N}_{t_\varsigma}.$$

Note for later use that, if  $\omega \in \Omega$ , multiplying (3.4) on the right by  $H_\omega$  we obtain that

$$(3.5) \quad {}^p\underline{N}_{t_\varsigma \omega} = \underline{N}_{t_\varsigma \omega} = \sum_{z \in W_f} v^{\ell(z)} N_{t_\varsigma z \omega}.$$

Lemma 3.4 shows that the map  $\mathcal{H}_{\text{ext}} \rightarrow \mathcal{M}_{\text{ext}}^{\text{asph}}$  defined by  $h \mapsto \underline{N}_{t_\varsigma} \cdot h$  factors through a morphism of right  $\mathcal{H}_{\text{ext}}$ -modules

$$(3.6) \quad \varphi : \mathcal{M}_{\text{ext}}^{\text{sph}} \rightarrow \mathcal{M}_{\text{ext}}^{\text{asph}}.$$

The main technical result of the paper is the following.

THEOREM 3.5. — Assume that  $p$  is good for  $G$ . Then for any  $w \in {}^fW_{\text{ext}}$  we have

$$\varphi({}^p\underline{M}_w) = {}^p\underline{N}_{t_\varsigma \cdot w}.$$

Remark 3.6. —

- (1) Theorem 3.5 implies in particular that  $\varphi$  is injective. (Of course, this can also be seen more directly.)
- (2) To deduce Theorem 1.4 from Theorem 3.5, one simply observes that  $\varphi$  restricts to a morphism of right  $\mathcal{H}$ -modules from the submodule of  $\mathcal{M}_{\text{ext}}^{\text{sph}}$  generated by  $N_{\omega_\varsigma^{-1}}$  to  $\mathcal{M}^{\text{asph}}$ . Now the latter submodule identifies with  $\mathcal{M}_{\varsigma}^{\text{sph}}$  (see Remark 3.1), so that Theorem 1.4 becomes the special case of Theorem 3.5 when  $w \in \omega_\varsigma^{-1}W$ .

## 4. Proof of Theorem 3.5

### 4.1. Categorification and $p$ -canonical bases

The proof of Theorem 3.5 will use the geometric description of the  $p$ -canonical bases in terms of parity complexes, which we now recall. For this we need to choose a field  $\mathbb{K}$  of coefficients for the parity complexes, which should be of characteristic  $p$  but might differ from  $\mathbb{k}$ . In fact, for technical reasons we will take for  $\mathbb{K}$  a finite field.

We also choose a prime number  $\ell \neq p$ , and assume that  $\mathbb{K}$  contains a nontrivial  $\ell^{\text{th}}$  root of unity.

We now fix an algebraically closed field  $\mathbb{F}$  of characteristic  $\ell$ . Let  $G^\vee$  be the connected reductive algebraic group over  $\mathbb{F}$  which is Langlands dual to  $G$ . By definition, this group comes with a maximal torus  $T^\vee \subset G^\vee$  whose cocharacter lattice is  $\mathbf{X}$ . We will denote by  $B^\vee \subset G^\vee$  the Borel subgroup containing  $T^\vee$  whose  $T^\vee$ -weights are the negative coroots of  $(G, T)$ . We set  $\mathcal{K} := \mathbb{F}((z))$ ,  $\mathcal{O} := \mathbb{F}[[z]]$ , and denote by  $G_{\mathcal{K}}^\vee$ , resp.  $G_{\mathcal{O}}^\vee$ , the ind-group scheme, resp. group scheme, over  $\mathbb{F}$  which represents the functor  $R \mapsto G^\vee(R((z)))$ , resp.  $R \mapsto G^\vee(R[[z]])$ . We also denote by  $I^\vee \subset G_{\mathcal{K}}^\vee$  the Iwahori subgroup, i.e. the inverse image of  $B^\vee$  under the evaluation morphism  $G_{\mathcal{O}}^\vee \rightarrow G^\vee$ . We then consider the affine flag variety

$$\mathcal{Fl} := G_{\mathcal{K}}^\vee / I^\vee.$$

Following [JMW14] we can consider the category  $\text{Parity}_{I^\vee}(\mathcal{Fl}, \mathbb{K})$  of  $I^\vee$ -equivariant parity (étale)  $\mathbb{K}$ -complexes on  $\mathcal{Fl}$ . The  $I^\vee$ -orbits on  $\mathcal{Fl}$  are parametrized in a natural way by  $W_{\text{ext}}$ ; we will denote by  $\mathcal{Fl}_w$  the orbit corresponding to  $w$  (so that  $\dim(\mathcal{Fl}_w) = \ell(w)$ ). For any  $w \in W_{\text{ext}}$ , there exists a unique indecomposable parity complex  $\mathcal{E}_w$  on  $\mathcal{Fl}$  which is supported on  $\overline{\mathcal{Fl}_w}$  and whose restriction to  $\mathcal{Fl}_w$  is  $\mathbb{K}_{\mathcal{Fl}_w}[\ell(w)]$ . Then the assignment  $(w, n) \mapsto \mathcal{E}_w[n]$  defines a bijection between  $W_{\text{ext}} \times \mathbb{Z}$  and the set of isomorphism classes of indecomposable objects in  $\text{Parity}_{I^\vee}(\mathcal{Fl}, \mathbb{K})$ .

The usual convolution construction endows  $\text{Parity}_{I^\vee}(\mathcal{Fl}, \mathbb{K})$  with the structure of a monoidal category. (The fact that a convolution of parity complexes is parity is proved in [JMW14, Section 4.1].) In particular, the split Grothendieck group

$$[\text{Parity}_{I^\vee}(\mathcal{Fl}, \mathbb{K})]$$

has a natural product; we will in fact view this ring as a  $\mathbb{Z}[v^{\pm 1}]$ -algebra, where  $v$  acts via the automorphism induced by the cohomological shift [1]. It is well known (see [JMW14, JW17, Spr82]) that there exists a unique  $\mathbb{Z}[v^{\pm 1}]$ -algebra isomorphism

$$(4.1) \quad \mathcal{H}_{\text{ext}} \xrightarrow{\sim} [\text{Parity}_{I^\vee}(\mathcal{Fl}, \mathbb{K})]$$

sending  $\underline{H}_s$  to  $[\mathcal{E}_s]$  for any  $s \in S$  and  $H_\omega$  to  $[\mathcal{E}_\omega]$  for any  $\omega \in \Omega$ . Then for  $w \in W_{\text{ext}}$ , the element  ${}^p\underline{H}_w$  is the inverse image of  $[\mathcal{E}_w]$  under (4.1), see e.g. [RW18, Part 3]. Recall also that the  $p$ -Kazhdan–Lusztig polynomials are the elements  $({}^ph_{y,w})_{y,w \in W_{\text{ext}}}$  of  $\mathbb{Z}[v^{\pm 1}]$  such that

$${}^p\underline{H}_w = \sum_{y \in W_{\text{ext}}} {}^ph_{y,w} \cdot H_y.$$

In order to categorify the module  $\mathcal{M}_{\text{ext}}^{\text{asph}}$ , we consider the category of “Iwahori–Whittaker” parity complexes  $\text{Parity}_{I^\vee, +}(\mathcal{Fl}, \mathbb{K})$  on  $\mathcal{Fl}$ . These objects are defined using the action of the unipotent radical  $I_{\mathbf{u}}^{\vee, +}$  of the Iwahori subgroup  $I^{\vee, +}$  associated with the Borel subgroup  $B^{\vee, +}$  of  $G^\vee$  which is opposite to  $B^\vee$  with respect to  $T^\vee$ ; see [RW18, Section 11] for details. (Here we use our assumption on  $\ell^{\text{th}}$  roots of unity in  $\mathbb{K}$ .) The  $I_{\mathbf{u}}^{\vee, +}$ -orbits on  $\mathcal{Fl}$  are parametrized in a natural way by  $W_{\text{ext}}$ ; but only those corresponding to elements in  ${}^fW_{\text{ext}}$  support nonzero Iwahori–Whittaker local systems. Therefore the isomorphism classes of indecomposable objects in  $\text{Parity}_{I^\vee, +}(\mathcal{Fl}, \mathbb{K})$  are naturally in bijection with  ${}^fW_{\text{ext}} \times \mathbb{Z}$ ; we will denote by  $\mathcal{E}_w^{TW}$  the object associated with  $(w, 0)$ .



The convolution construction defines a right action of the monoidal category  $\mathbf{Parity}_{I^\vee}(\mathcal{F}l, \mathbb{K})$  on the category  $\mathbf{Parity}_{I\mathcal{W}}(\mathcal{F}l, \mathbb{K})$ , and there exists a unique isomorphism of right  $\mathcal{H}_{\text{ext}}$ -modules

$$(4.2) \quad \mathcal{M}_{\text{ext}}^{\text{asph}} \xrightarrow{\sim} [\mathbf{Parity}_{I\mathcal{W}}(\mathcal{F}l, \mathbb{K})]$$

sending  $N_{\text{id}}$  to  $\mathcal{E}_{\text{id}}^{I\mathcal{W}}$ . Then for  $w \in {}^fW_{\text{ext}}$ ,  ${}^pN_w$  is the inverse image of  $[\mathcal{E}_w^{I\mathcal{W}}]$  under this isomorphism, see [RW18, Section 11].

Finally, we explain the categorification of  $\mathcal{M}_{\text{ext}}^{\text{sph}}$ . We consider the “opposite affine Grassmannian”

$$\mathcal{G}r^{\text{op}} := G_{\mathcal{O}}^\vee \backslash G_{\mathcal{H}}^\vee.$$

This variety admits an action of  $I^\vee$  induced by right multiplication on  $G_{\mathcal{H}}^\vee$ , and we can consider the corresponding category of parity complexes  $\mathbf{Parity}_{I^\vee}(\mathcal{G}r^{\text{op}}, \mathbb{K})$ . The  $I^\vee$ -orbits on  $\mathcal{G}r^{\text{op}}$  are parametrized by  ${}^fW_{\text{ext}}$ ; therefore the indecomposable objects in  $\mathbf{Parity}_{I^\vee}(\mathcal{G}r^{\text{op}}, \mathbb{K})$  are parametrized in a natural way by  ${}^fW_{\text{ext}} \times \mathbb{Z}$ . The object associated with  $(w, 0)$  (for  $w \in {}^fW_{\text{ext}}$ ) will be denoted  $\mathcal{F}_w$ .

Again, the convolution construction defines a right action of the monoidal category  $\mathbf{Parity}_{I^\vee}(\mathcal{F}l, \mathbb{K})$  on the category  $\mathbf{Parity}_{I^\vee}(\mathcal{G}r^{\text{op}}, \mathbb{K})$ , and there exists a unique isomorphism of right  $\mathcal{H}_{\text{ext}}$ -modules

$$\mathcal{M}_{\text{ext}}^{\text{sph}} \xrightarrow{\sim} [\mathbf{Parity}_{I^\vee}(\mathcal{G}r^{\text{op}}, \mathbb{K})]$$

sending  $M_{\text{id}}$  to  $\mathcal{F}_{\text{id}}$ . Using [ACR18, Lemma A.5] and the construction of the  $p$ -canonical basis in  $\mathcal{M}_{\text{ext}}^{\text{sph}}$ , one can check that, for  $w \in {}^fW_{\text{ext}}$ ,  ${}^pM_w$  is the inverse image of  $[\mathcal{F}_w]$  under this isomorphism.

## 4.2. Parity complexes on affine Grassmannians

From now on we assume that  $p$  is good for  $G$ .

Consider the “usual” affine Grassmannian

$$\mathcal{G}r := G_{\mathcal{H}}^\vee / G_{\mathcal{O}}^\vee,$$

and the  $G_{\mathcal{O}}^\vee$ -equivariant constructible derived category  $D_{G_{\mathcal{O}}^\vee}^b(\mathcal{G}r, \mathbb{K})$ . This category possesses a natural perverse t-structure, whose heart will be denoted  $\mathbf{Perv}_{G_{\mathcal{O}}^\vee}(\mathcal{G}r, \mathbb{K})$ .

Under our assumptions that  $p$  is good for  $G$  (equivalently, for  $G^\vee$ ) and that  $G$  has a simply-connected derived subgroup (so that the quotient of  $X^*(T^\vee)$  by the coroot lattice of  $G$  is torsion-free), it is known that the equivariant cohomology  $\mathbf{H}_{G_{\mathcal{O}}^\vee}^\bullet(\text{pt}; \mathbb{K}) = \mathbf{H}_{G_{\mathcal{O}}^\vee}^\bullet(\text{pt}; \mathbb{K})$  vanishes in odd degrees; see e.g. [JMW14, Section 2.6] or [MR18, Section 3.2] for references. Therefore, the theory developed in [JMW14] applies in this context, and we will denote by  $\mathbf{Parity}_{G_{\mathcal{O}}^\vee}(\mathcal{G}r, \mathbb{K})$  the corresponding category of parity complexes.

For  $\lambda \in \mathbf{X}$ , we set  $L_\lambda := z^\lambda \cdot G_{\mathcal{O}}^\vee \in \mathcal{G}r$ . Then the assignment  $\lambda \mapsto G_{\mathcal{O}}^\vee \cdot L_\lambda$  induces a bijection between  $\mathbf{X}^+$  and the set of  $G_{\mathcal{O}}^\vee$ -orbits on  $\mathcal{G}r$ . Therefore, the isomorphism classes of indecomposable objects in  $\mathbf{Parity}_{G_{\mathcal{O}}^\vee}(\mathcal{G}r, \mathbb{K})$  are parametrized in a natural way by  $\mathbf{X}^+ \times \mathbb{Z}$ ; for  $\lambda \in \mathbf{X}^+$  we will denote by  $\mathcal{E}_\lambda^{\text{sph}}$  the object associated with  $(\lambda, 0)$ .

The following result is proved in [JMW16] under some technical assumptions, and in [MR18, Corollary 1.6] in the present generality. (This claim is known to be false if we remove the assumption that  $p$  is good for  $G$ , see [JMW16].)

**THEOREM 4.1.** — *For any  $\lambda \in \mathbf{X}^+$ , the object  $\mathcal{E}_\lambda^{\text{sph}}$  is perverse.*

*Remark 4.2.* — From the combinatorial point of view, this theorem says that if  $w \in W_{\text{ext}}$  is maximal in  $W_{\text{f}} w W_{\text{f}}$ , then  ${}^p H_w$  belongs to  $\bigoplus_y \mathbb{Z} \cdot H_y$ .

The other result which we will need is the main result of [BGM<sup>+</sup>19]. Here we consider the Iwahori–Whittaker derived category of sheaves on  $\mathcal{G}r$ , denoted  $D_{\text{IW}}^b(\mathcal{G}r, \mathbb{K})$ . This category is endowed with the perverse t-structure, whose heart will be denoted  $\text{Perv}_{\text{IW}}(\mathcal{G}r, \mathbb{K})$ . This abelian category admits a natural structure of highest weight category (in the sense considered e.g. in [Ric, Section 7]), and moreover the realization functor provides an equivalence of triangulated categories

$$D^b \text{Perv}_{\text{IW}}(\mathcal{G}r, \mathbb{K}) \xrightarrow{\sim} D_{\text{IW}}^b(\mathcal{G}r, \mathbb{K}).$$

The  $I_{\mathbf{u}}^{\vee,+}$ -orbits on  $\mathcal{G}r$  are parametrized by  $\mathbf{X}$ , and those which support a nonzero Iwahori–Whittaker local system are the ones parametrized by elements in  $\varsigma + \mathbf{X}^+$  (i.e. by strictly dominant weights). In particular, no orbit in the boundary of the orbit associated with  $\varsigma$  supports such a local system; therefore the corresponding standard perverse sheaf is simple, and isomorphic to the associated costandard perverse sheaf (see [BGM<sup>+</sup>19, Equation (3.2)]). Hence this object is also parity, and will be denoted  $\mathcal{F}_\varsigma^{\text{IW}}$ .

The following result is proved in [BGM<sup>+</sup>19]. (The first claim holds without any assumption on  $p$ ; however for the second assertion we need the restriction that  $p$  is good.) Here we denote by  $\star^{G_\varsigma^\vee}$  the natural convolution bifunctor

$$D_{\text{IW}}^b(\mathcal{G}r, \mathbb{K}) \times D_{G_\varsigma^\vee}^b(\mathcal{G}r, \mathbb{K}) \rightarrow D_{\text{IW}}^b(\mathcal{G}r, \mathbb{K})$$

(see [BGM<sup>+</sup>19] for details).

**THEOREM 4.3.** — *The functor*

$$\Psi : D_{G_\varsigma^\vee}^b(\mathcal{G}r, \mathbb{K}) \rightarrow D_{\text{IW}}^b(\mathcal{G}r, \mathbb{K})$$

*defined by  $\Psi(\mathcal{F}) = \mathcal{F}_\varsigma^{\text{IW}} \star^{G_\varsigma^\vee} \mathcal{F}$  is t-exact for the perverse t-structures, and restricts to an equivalence of categories  $\text{Perv}_{G_\varsigma^\vee}(\mathcal{G}r, \mathbb{K}) \xrightarrow{\sim} \text{Perv}_{\text{IW}}(\mathcal{G}r, \mathbb{K})$ . Moreover, for any  $\lambda \in \mathbf{X}$ , the object  $\Psi(\mathcal{E}_\lambda^{\text{sph}})$  is a tilting perverse sheaf.*

The consequence of Theorems 4.1 and 4.3 that we will use below is the following.

**COROLLARY 4.4.** — *For any  $\mathcal{G}, \mathcal{G}'$  in  $\text{Parity}_{G_\varsigma^\vee}(\mathcal{G}r, \mathbb{K})$ , the morphism*

$$\text{Hom}_{D_{G_\varsigma^\vee}^b(\mathcal{G}r, \mathbb{K})}(\mathcal{G}, \mathcal{G}') \rightarrow \text{Hom}_{D_{\text{IW}}^b(\mathcal{G}r, \mathbb{K})}(\Psi(\mathcal{G}), \Psi(\mathcal{G}'))$$

*induced by  $\Psi$  is surjective.*

*Proof.* — Any object of  $\text{Parity}_{G_\varsigma^\vee}(\mathcal{G}r, \mathbb{K})$  is a direct sum of cohomological shifts of objects of the form  $\mathcal{E}_\lambda^{\text{sph}}$  (with  $\lambda \in \mathbf{X}^+$ ); therefore to prove the corollary it suffices

to prove that for any  $\lambda, \mu \in \mathbf{X}^+$  and  $n \in \mathbb{Z}$  the functor  $\Psi$  induces a surjection

$$(4.3) \quad \mathrm{Hom}_{D_{G_{\mathcal{O}}^{\vee}}^b(\mathcal{Gr},)}(\mathcal{E}_{\lambda}^{\mathrm{sph}}, \mathcal{E}_{\mu}^{\mathrm{sph}}[n]) \rightarrow \mathrm{Hom}_{D_{\mathcal{IW}}^b(\mathcal{Gr}, \mathbb{K})}(\Psi(\mathcal{E}_{\lambda}^{\mathrm{sph}}), \Psi(\mathcal{E}_{\mu}^{\mathrm{sph}})[n]).$$

Now, by Theorem 4.3 the objects  $\Psi(\mathcal{E}_{\lambda}^{\mathrm{sph}})$  and  $\Psi(\mathcal{E}_{\mu}^{\mathrm{sph}})$  are tilting perverse sheaves; therefore the right-hand side vanishes unless  $n = 0$ . And if  $n = 0$ , since  $\mathcal{E}_{\lambda}^{\mathrm{sph}}$  and  $\mathcal{E}_{\mu}^{\mathrm{sph}}$  are perverse, and since  $\Psi$  restricts to an equivalence on perverse sheaves, the map (4.3) is an isomorphism in this case.  $\square$

We can now give the proof of Lemma 3.4.

*Proof of Lemma 3.4.* — One can easily check using (2.1) (and the fact that  $\ell(x) = \ell(x^{-1})$  for any  $x \in W_{\mathrm{ext}}$ ) that  $t_{\varsigma}$  is of maximal length in  $t_{\varsigma} \cdot W_{\mathrm{f}}$ . Therefore, the  $I_{\mathrm{u}}^{\vee, +}$ -orbit in  $\mathcal{Fl}$  associated with  $t_{\varsigma}$  is the inverse image under the projection  $\pi : \mathcal{Fl} \rightarrow \mathcal{Gr}$  of the orbit of  $L_{\varsigma}$ . Using [ACR18, Lemma A.5] we deduce that

$$\mathcal{E}_{t_{\varsigma}}^{\mathcal{IW}} = \pi^*(\mathcal{F}_{\varsigma}^{\mathcal{IW}})[\ell(w_{\mathrm{f}})].$$

It follows that  ${}^pN_{t_{\varsigma}} = \sum_{z \in W_{\mathrm{f}}} v^{\ell(z)} N_{t_{\varsigma} \cdot z}$ , and that for any  $s \in S_{\mathrm{f}}$  we have  ${}^pN_{t_{\varsigma}} \cdot \underline{H}_s = (v + v^{-1}) \cdot {}^pN_{t_{\varsigma}}$ . The claims about  $\underline{N}_{t_{\varsigma}}$  follow, taking  $p \gg 0$  (see Remark 3.3).  $\square$

### 4.3. Fullness

To prove Theorem 3.5 we will consider a categorification of  $\varphi$ . For this, we work with the  $G_{\mathcal{O}}^{\vee}$ -equivariant derived category  $D_{G_{\mathcal{O}}^{\vee}}^b(\mathcal{Fl}, \mathbb{K})$ . This category admits a right action of the  $I^{\vee}$ -equivariant derived category  $D_{I^{\vee}}^b(\mathcal{Fl}, \mathbb{K})$  (by convolution, as usual), and it is clear that there exists a canonical equivalence of triangulated categories

$$\iota : D_{I^{\vee}}^b(\mathcal{Gr}^{\mathrm{op}}, \mathbb{K}) \xrightarrow{\sim} D_{G_{\mathcal{O}}^{\vee}}^b(\mathcal{Fl}, \mathbb{K})$$

sending  $\mathcal{F}_{\mathrm{id}}$  to  $\mathbb{K}_{G_{\mathcal{O}}^{\vee}/I^{\vee}}[\ell(w_{\mathrm{f}})]$  (where  $\mathcal{Gr}^{\mathrm{op}}$  is as in Section 4.1) and commuting with the right actions of  $D_{I^{\vee}}^b(\mathcal{Fl}, \mathbb{K})$  on both sides. Moreover, the theory of parity complexes from [JMW14] applies in  $D_{G_{\mathcal{O}}^{\vee}}^b(\mathcal{Fl}, \mathbb{K})$  also, and  $\iota$  restricts to an equivalence of categories

$$\iota_{\mathrm{Par}} : \mathrm{Parity}_{I^{\vee}}(\mathcal{Gr}^{\mathrm{op}}, \mathbb{K}) \xrightarrow{\sim} \mathrm{Parity}_{G_{\mathcal{O}}^{\vee}}(\mathcal{Fl}, \mathbb{K}),$$

where in the right-hand side  $\mathrm{Parity}_{G_{\mathcal{O}}^{\vee}}(\mathcal{Fl}, \mathbb{K})$  means the full subcategory of parity complexes in  $D_{G_{\mathcal{O}}^{\vee}}^b(\mathcal{Fl}, \mathbb{K})$ . In particular, the indecomposable objects in the category  $\mathrm{Parity}_{G_{\mathcal{O}}^{\vee}}(\mathcal{Fl}, \mathbb{K})$  are the objects  $\iota(\mathcal{F}_w)[n]$  for  $w \in {}^{\mathrm{f}}W_{\mathrm{ext}}$  and  $n \in \mathbb{Z}$ , and we have a canonical isomorphism

$$\mathcal{M}_{\mathrm{ext}}^{\mathrm{sph}} \xrightarrow{\sim} [\mathrm{Parity}_{G_{\mathcal{O}}^{\vee}}(\mathcal{Fl}, \mathbb{K})]$$

sending  ${}^pM_w$  to  $[\iota(\mathcal{F}_w)]$  for any  $w \in {}^{\mathrm{f}}W_{\mathrm{ext}}$ .

We now consider the functor

$$\Phi : D_{G_{\mathcal{O}}^{\vee}}^b(\mathcal{Fl}, \mathbb{K}) \rightarrow D_{\mathcal{IW}}^b(\mathcal{Fl}, \mathbb{K})$$

defined by

$$\Phi(\mathcal{F}) = \mathcal{F}_{\varsigma}^{\mathcal{IW}} \star_{G_{\mathcal{O}}^{\vee}} \mathcal{F},$$

where  $\star^{G_\theta^\vee}$  now denotes the natural convolution bifunctor

$$D_{\mathcal{IW}}^b(\mathcal{G}r, \mathbb{K}) \times D_{G_\theta^\vee}^b(\mathcal{F}l, \mathbb{K}) \rightarrow D_{\mathcal{IW}}^b(\mathcal{F}l, \mathbb{K}).$$

LEMMA 4.5. — *The functor  $\Phi$  sends parity complexes to parity complexes. Moreover, the map on split Grothendieck groups induced by the restriction*

$$\Phi_{\text{Par}} : \text{Parity}_{G_\theta^\vee}(\mathcal{F}l, \mathbb{K}) \rightarrow \text{Parity}_{\mathcal{IW}}(\mathcal{F}l, \mathbb{K})$$

is  $\varphi$ .

*Proof.* — The proof of the first claim is similar to that of [BGM<sup>+</sup>19, Lemma 4.14]. For the second claim, we observe that the map induced by  $\Phi_{\text{Par}}$  is clearly a morphism of right  $\mathcal{H}_{\text{ext}}$ -modules. Since  $\mathcal{M}_{\text{ext}}^{\text{sph}}$  is a cyclic module, this reduces the proof to checking that the image of  $[\mathbb{K}_{G_\theta^\vee/I^\vee}[\ell(w_f)]]$  corresponds to  $\varphi(M_e) = \underline{N}_{t_\zeta}$  under (4.2). However we have

$$\Phi\left(\mathbb{K}_{G_\theta^\vee/I^\vee}[\ell(w_f)]\right) = \pi^{*\iota}\left(\mathcal{F}_\zeta^{\mathcal{IW}}\right)[\ell(w_f)],$$

where  $\pi : \mathcal{F}l \rightarrow \mathcal{G}r$  is the projection. As observed in the proof of Lemma 3.4, the right-hand side is  $\mathcal{E}_{t_\zeta}^{\mathcal{IW}}$ , whose class in the Grothendieck group corresponds to  $\underline{N}_{t_\zeta}$  by definition. The claim follows, using the first equality in Lemma 3.4.  $\square$

The key step in our proof of Theorem 3.5 is the following claim.

PROPOSITION 4.6. — *The functor  $\Phi_{\text{Par}}$  from Lemma 4.5 is full.*

*Proof.* — It is easily seen that any object in  $\text{Parity}_{G_\theta^\vee}(\mathcal{F}l, \mathbb{K})$  is a direct sum of direct summands of objects of the form  $\mathbb{K}_{G_\theta^\vee/I^\vee} \star^{I^\vee} \mathcal{E}$  with  $\mathcal{E}$  in  $\text{Parity}_{I^\vee}(\mathcal{F}l, \mathbb{K})$ , where  $\star^{I^\vee}$  is the natural convolution bifunctor

$$\text{Parity}_{G_\theta^\vee}(\mathcal{F}l, \mathbb{K}) \times \text{Parity}_{I^\vee}(\mathcal{F}l, \mathbb{K}) \rightarrow \text{Parity}_{G_\theta^\vee}(\mathcal{F}l, \mathbb{K}).$$

Now any functor of the form  $(-) \star^{I^\vee} \mathcal{E}$  (with  $\mathcal{E}$  in  $\text{Parity}_{I^\vee}(\mathcal{F}l, \mathbb{K})$ ) admits a left adjoint of the form  $(-) \star^{I^\vee} \mathcal{E}'$  with  $\mathcal{E}'$  in  $\text{Parity}_{I^\vee}(\mathcal{F}l, \mathbb{K})$ , hence this remark reduces the proof of fullness of  $\Phi_{\text{Par}}$  to proving that for any  $\mathcal{F}$  in  $\text{Parity}_{G_\theta^\vee}(\mathcal{F}l, \mathbb{K})$  the map

$$\text{Hom}_{D_{G_\theta^\vee}^b(\mathcal{F}l, \mathbb{K})}^\bullet(\mathcal{G}, \mathbb{K}_{G_\theta^\vee/I^\vee}) \rightarrow \text{Hom}_{D_{\mathcal{IW}}^b(\mathcal{F}l, \mathbb{K})}^\bullet(\Phi(\mathcal{G}), \Phi(\mathbb{K}_{G_\theta^\vee/I^\vee}))$$

induced by  $\Phi$  is surjective. If  $\pi$  is as in the proof of Lemma 3.4 (or of Lemma 4.5), then we have

$$\mathbb{K}_{G_\theta^\vee/I^\vee} = \pi^* \mathcal{E}_0^{\text{sph}}, \quad \Phi(\mathbb{K}_{G_\theta^\vee/I^\vee}) = \pi^* \left( \mathcal{F}_\zeta^{\mathcal{IW}} \right).$$

Since  $\pi^* \cong \pi^![-2\ell(w_f)]$ , using adjunction we deduce isomorphisms

$$\text{Hom}_{D_{G_\theta^\vee}^b(\mathcal{F}l, \mathbb{K})}^\bullet(\mathcal{G}, \mathbb{K}_{G_\theta^\vee/I^\vee}) \cong \text{Hom}_{D_{G_\theta^\vee}^b(\mathcal{G}r, \mathbb{K})}^\bullet(-2\ell(w_f), \pi_!(\mathcal{G}), \mathcal{E}_0^{\text{sph}}),$$

$$\text{Hom}_{D_{\mathcal{IW}}^b(\mathcal{F}l, \mathbb{K})}^\bullet(\Phi(\mathcal{G}), \Phi(\mathbb{K}_{G_\theta^\vee/I^\vee})) \cong \text{Hom}_{D_{\mathcal{IW}}^b(\mathcal{G}r, \mathbb{K})}^\bullet(-2\ell(w_f), \pi_!\Phi(\mathcal{G}), \mathcal{F}_\zeta^{\mathcal{IW}}).$$

Now we have  $\pi_!\Phi(\mathcal{G}) \cong \Psi(\pi_!\mathcal{G})$ , where  $\Psi$  is as in Theorem 4.3; hence we are reduced to proving that the morphism

$$\text{Hom}_{D_{G_\theta^\vee}^b(\mathcal{G}r, \mathbb{K})}^\bullet(\pi_!(\mathcal{G}), \mathcal{E}_0^{\text{sph}}) \rightarrow \text{Hom}_{D_{\mathcal{IW}}^b(\mathcal{G}r, \mathbb{K})}^\bullet(\Psi(\pi_!\mathcal{G}), \mathcal{F}_\zeta^{\mathcal{IW}})$$

induced by  $\Psi$  is surjective. However,  $\pi_! \mathcal{G}$  is parity, so that the claim follows from Corollary 4.4.  $\square$

#### 4.4. Proof of Theorem 3.5

Since the functor  $\Phi$  is a full functor between Krull–Schmidt categories, it must send indecomposable objects to indecomposable objects. Indeed, this follows from the observation that any quotient of a local ring is local. Using support considerations it is not difficult to deduce that for any  $w \in {}^f W_{\text{ext}}$  we have

$$\Phi(\iota(\mathcal{F}_w)) \cong \mathcal{E}_{t_\varsigma w}^{TW}.$$

Passing to classes in the split Grothendieck group we deduce the formula of Theorem 3.5.

### 5. Application: a character formula for simple $G$ -modules

In this section we return to the setting of Sections 1–3; in particular,  $G$  is a connected reductive algebraic group with simply connected derived subgroup over an algebraically closed field  $\mathbb{k}$  of characteristic  $p$ . We will assume that  $p > h$ , where  $h$  is the Coxeter number of  $G$ . (In particular, this condition implies that  $p$  is good for  $G$ , so that Theorem 3.5 is applicable.)

#### 5.1. The tilting character formula

We set

$$\mathcal{M}_{\text{ext}}^{\text{asph}} := \mathbb{Z} \otimes_{\mathbb{Z}[v^{\pm 1}]} \mathcal{M}_{\text{ext}}^{\text{asph}},$$

where  $\mathbb{Z}$  is considered as a  $\mathbb{Z}[v^{\pm 1}]$ -module via  $v \mapsto 1$ . This  $\mathbb{Z}$ -module is a right module over

$$\mathbb{Z} \otimes_{\mathbb{Z}[v^{\pm 1}]} \mathcal{H}_{\text{ext}} = \mathbb{Z}[W_{\text{ext}}].$$

We will denote by  $\text{Rep}(G)$  the abelian category of finite-dimensional algebraic  $G$ -modules. The simple objects in this category are labelled in a natural way by the subset  $\mathbf{X}^+ \subset \mathbf{X}$  of dominant weights; as in Section 1.1 we will denote by  $\mathbb{L}(\lambda)$  the simple  $G$ -module of highest weight  $\lambda \in \mathbf{X}^+$ .

We consider the dilated and shifted action of  $W$  on  $\mathbf{X}$  defined by

$$w \cdot_p \lambda = w(\lambda + \varsigma) - \varsigma, \quad t_\mu \cdot_p \lambda = \lambda + p\mu$$

for  $w \in W_f$  and  $\lambda, \mu \in \mathbf{X}$ . (It is a classical fact that this action does not depend on the choice of  $\varsigma$ .) We then denote by  $\text{Rep}_\varnothing(G)$  the “extended principal block” of  $\text{Rep}(G)$ , i.e. the Serre subcategory generated by the simple objects  $\mathbb{L}(w \cdot_p 0)$  with  $w \in {}^f W_{\text{ext}}$ . (Here, under our assumptions, for  $w \in W_{\text{ext}}$  we have  $w \cdot_p 0 \in \mathbf{X}^+$  iff  $w \in {}^f W_{\text{ext}}$ .)

For  $\lambda \in \mathbf{X}^+$  we also denote by  $\Delta(\lambda)$ ,  $\nabla(\lambda)$  and  $\mathbb{T}(\lambda)$  the Weyl, induced, and indecomposable tilting  $G$ -modules of highest weight  $\lambda$  (see [RW18, Section 3.1]). If  $\lambda = w \cdot_p 0$  for some  $w \in {}^f W_{\text{ext}}$ , then these objects belong to  $\text{Rep}_\varnothing(G)$ .

In the following lemma,  $T_\mu^\nu$  is the translation functor from the  $\mu$ -block to the  $\nu$ -block of  $\text{Rep}(G)$ , see [Jan03, Chapter II.7]. See also Remark 2.1 for the definition of  $\omega_\varsigma$ .

LEMMA 5.1. — *For any  $\omega \in \Omega$  we have*

$$\mathbb{T}(p\varsigma + \omega \cdot_p 0) \cong T_{\omega_\varsigma \cdot_p (-\varsigma)}^{\omega_\varsigma \omega \cdot_p 0}(\mathbb{L}((p-1)\varsigma)).$$

Moreover, for any  $\lambda \in \mathbf{X}^+$  we have

$$(\mathbb{T}(p\varsigma + \omega \cdot_p 0) : \nabla(\lambda)) = \begin{cases} 1 & \text{if } \lambda = t_\varsigma x \omega \cdot_p 0 \text{ for some } x \in W_f; \\ 0 & \text{otherwise.} \end{cases}$$

*Proof.* — We have  $p\varsigma + \omega \cdot_p 0 = t_\varsigma \omega_\varsigma^{-1} \cdot_p (\omega_\varsigma \omega \cdot_p 0)$ , and  $(p-1)\varsigma = t_\varsigma \omega_\varsigma^{-1} \cdot_p (\omega_\varsigma \cdot_p (-\varsigma))$ . Moreover,  $p\varsigma + \omega \cdot_p 0$  is maximal among the elements of the form  $t_\varsigma \omega_\varsigma^{-1} \cdot_p (\omega_\varsigma x \omega \cdot_p 0)$  with  $x \in W_f$ . Hence by [Jan03, Proposition E.11] we have

$$\mathbb{T}(p\varsigma + \omega \cdot_p 0) \cong T_{\omega_\varsigma \cdot_p (-\varsigma)}^{\omega_\varsigma \omega \cdot_p 0}(\mathbb{T}((p-1)\varsigma)).$$

Now by [Jan03, Remark in Section II.3.19], the Steinberg module  $\mathbb{L}((p-1)\varsigma)$  is isomorphic to  $\Delta((p-1)\varsigma)$  and to  $\nabla((p-1)\varsigma)$ , hence is tilting. It is also clearly indecomposable, so that  $\mathbb{T}((p-1)\varsigma) = \mathbb{L}((p-1)\varsigma)$ . The first claim follows.

The second claim follows from the first one (and the fact that  $\mathbb{L}((p-1)\varsigma) = \nabla((p-1)\varsigma)$ ) in view of [Jan03, Proposition II.7.13].  $\square$

For any  $s \in S$  we choose a weight  $\mu_s$  as in [RW18, Section 3.1] (i.e. a “generic” weight on the  $s$ -wall of the fundamental alcove for the dilated and shifted action) and consider the exact selfadjoint endofunctor

$$\Theta_s := \bigoplus_{\omega \in \Omega} T_{\omega \cdot_p \mu_s}^{\omega \cdot_p 0} T_{\omega \cdot_p 0}^{\omega \cdot_p \mu_s}$$

of  $\text{Rep}_\varnothing(G)$ . (Here the sum might be infinite but, for each object  $V$  of  $\text{Rep}_\varnothing(G)$ , only finitely many of these functors applied to  $V$  do not vanish; so the functor  $\Theta_s$  is well defined.) If we denote by  $[\text{Rep}_\varnothing(G)]$  the Grothendieck group of the abelian category  $\text{Rep}_\varnothing(G)$ , and by  $[M]$  the class of an object  $M$ , then it is well known (see e.g. [RW18, Section 1.2]) that the assignment

$$1 \otimes N_w \mapsto [\Delta(w \cdot_p 0)] = [\nabla(w \cdot_p 0)]$$

induces an isomorphism of abelian groups

$$(5.1) \quad \mathbf{M}_{\text{ext}}^{\text{asph}} \xrightarrow{\sim} [\text{Rep}_\varnothing(G)].$$

Using [Jan03, Propositions II.7.11 and II.7.12] one can check that, under this identification, the endomorphism of the right-hand side induced by  $\Theta_s$  corresponds to the action of  $(1+s)$  on the left-hand side.

The following statement was conjectured in [RW18] (see in particular [RW18, Corollary 1.4.1]) and proved in [AMRW19].

THEOREM 5.2. — *Under the isomorphism (5.1),  $1 \otimes {}^p N_w$  is sent to  $[\mathbb{T}(w \cdot_p 0)]$  for any  $w \in {}^f W_{\text{ext}}$ .*

Remark 5.3. — In [AMRW19, RW18] we work with  $W$  instead of  $W_{\text{ext}}$ ; but the extension is immediate (see e.g. [AR19]).

## 5.2. $G_1T$ -modules

Let  $\dot{G}$  be the Frobenius twist of  $G$ , and let  $\text{Fr} : G \rightarrow \dot{G}$  be the Frobenius morphism. We denote by  $G_1$  the kernel of  $\text{Fr}$  (a normal finite subgroup scheme of  $G$ ) and by  $G_1T$ , resp.  $G_1B^+$ , the inverse image of the Frobenius twist  $\dot{T}$  of  $T$ , resp.  $\dot{B}^+$  of  $B^+$ , under  $\text{Fr}$ . (Here,  $B^+$  is the Borel subgroup of  $G$  opposite to  $B$  with respect to  $T$ .) We will identify the characters of  $\dot{T}$  with  $\mathbf{X}$ , in such a way that the composition of the Frobenius morphism  $T \rightarrow \dot{T}$  with the character  $\lambda$  of  $\dot{T}$  is the character  $p\lambda$  of  $T$ . We use similar conventions for  $\dot{B}^+$ .

We will denote by  $\text{Rep}(G_1T)$  the category of finite-dimensional algebraic  $G_1T$ -modules. As explained in [Jan03, Proposition II.9.6], the simple objects in  $\text{Rep}(G_1T)$  are in a canonical bijection with  $\mathbf{X}$ ; the simple module corresponding to  $\lambda$  will be denoted  $\hat{\mathbb{L}}(\lambda)$ . If  $\lambda$  is dominant and restricted then  $\hat{\mathbb{L}}(\lambda)$  is the restriction of  $\mathbb{L}(\lambda)$  to  $G_1T$ , and if  $\lambda, \mu \in \mathbf{X}$  we have

$$(5.2) \quad \hat{\mathbb{L}}(\lambda + p\mu) \cong \hat{\mathbb{L}}(\lambda) \otimes \mathbb{K}_{\dot{T}}(\mu),$$

where  $\mathbb{K}_{\dot{T}}(\mu)$  is seen as a  $G_1T$ -module via the surjection  $G_1T \rightarrow \dot{T}$ . In particular, these simple objects are completely determined by the objects  $(\mathbb{L}(\lambda) : \lambda \in \mathbf{X}^+)$ .

The category  $\text{Rep}(G_1T)$  also contains the *baby Verma modules*

$$\hat{Z}(\lambda) = \text{Coind}_{B^+}^{G_1B^+}(\mathbb{K}_{B^+}(\lambda))$$

for  $\lambda \in \mathbf{X}$ . With these conventions, we have a surjection of  $G_1T$ -modules  $\hat{Z}(\lambda) \rightarrow \hat{\mathbb{L}}(\lambda)$ , see [Jan03, Proposition 9.6(d)]. We also have canonical isomorphisms

$$(5.3) \quad \hat{Z}(\lambda + p\mu) \cong \hat{Z}(\lambda) \otimes \mathbb{K}_{\dot{T}}(\mu)$$

for  $\lambda, \mu \in \mathbf{X}$ .

The category  $\text{Rep}(G_1T)$  admits a “block” decomposition similar to that of  $\text{Rep}(G)$ ; see e.g. [Jan03, Section II.9.19]. Hence we can consider  $\text{Rep}_{\emptyset}(G_1T)$ , the “extended block” of the weight 0, i.e. the Serre subcategory generated by the simple objects  $\hat{\mathbb{L}}(\lambda)$  with  $\lambda \in W_{\text{ext}} \cdot_p 0$ . If  $\lambda \in W_{\text{ext}} \cdot_p 0$  then  $\hat{Z}(\lambda)$  also belongs to  $\text{Rep}_{\emptyset}(G_1T)$ . It is clear from the Steinberg tensor product formula [Jan03, Proposition II.3.16] that the restriction functor  $\text{Rep}(G) \rightarrow \text{Rep}(G_1T)$  restricts to a functor

$$(5.4) \quad \text{Rep}_{\emptyset}(G) \rightarrow \text{Rep}_{\emptyset}(G_1T).$$

The image of a  $G$ -module  $M$  under this functor will sometimes be denoted  $M|_{G_1T}$ .

As explained in [Jan03, Section II.9.22], the translation functors can be canonically “lifted” to the category  $\text{Rep}(G_1T)$ . In particular, this means that there exists an exact selfadjoint endofunctor of  $\text{Rep}_{\emptyset}(G_1T)$ , which for simplicity will also be denoted  $\Theta_s$ , and such that the following diagram commutes:

$$\begin{array}{ccc} \text{Rep}_{\emptyset}(G) & \xrightarrow{\Theta_s} & \text{Rep}_{\emptyset}(G) \\ (5.4) \downarrow & & \downarrow (5.4) \\ \text{Rep}_{\emptyset}(G_1T) & \xrightarrow{\Theta_s} & \text{Rep}_{\emptyset}(G_1T). \end{array}$$

LEMMA 5.4. — *For any  $w \in W_{\text{ext}}$  and any  $s \in S$ , in the Grothendieck group of  $\text{Rep}_\varnothing(G_1T)$  we have*

$$\left[ \Theta_s \left( \widehat{Z}(w \cdot_p 0) \right) \right] = \left[ \widehat{Z}(w \cdot_p 0) \right] + \left[ \widehat{Z}(ws \cdot_p 0) \right].$$

*Proof.* — This follows from [Jan03, Equations (2) and (3) in Section II.9.22].  $\square$

Remark 5.5. — The formula in Lemma 5.4 suggests that the Grothendieck group  $[\text{Rep}_\varnothing(G_1T)]$  is closely related with the right  $\mathbb{Z}[W]$ -module  $\mathbb{Z} \otimes_{\mathbb{Z}[v^{\pm 1}]} \mathcal{P}$ . However, two important remarks are in order. First, since  $\mathcal{A}$  is in bijection with  $W$  rather than  $W_{\text{ext}}$ , to make this precise we would have to work with the “true” principal block  $\text{Rep}_0(G_1T)$  in  $\text{Rep}(G_1T)$ , i.e. the Serre subcategory generated by simple modules  $\widehat{\mathbb{L}}(w \cdot_p 0)$  with  $w \in W$ . But even then a difficulty would remain, since the classes of baby Verma modules do *not* form a basis of the Grothendieck group  $[\text{Rep}_0(G_1T)]$ . We will not try to address this problem here.

### 5.3. Injective/projective $G_1T$ -modules

For  $\lambda \in \mathbf{X}$ , we will denote by  $\widehat{Q}(\lambda)$  the injective hull of  $\widehat{\mathbb{L}}(\lambda)$  as a  $G_1T$ -module, see [Jan03, Section II.11.3]. As explained in [Jan03, Equation (3) in Section II.11.5],  $\widehat{Q}(\lambda)$  is also the projective cover of  $\widehat{\mathbb{L}}(\lambda)$  in this category. As for simple and baby Verma modules, for  $\lambda, \mu \in \mathbf{X}$  we have

$$(5.5) \quad \widehat{Q}(\lambda + p\mu) \cong \widehat{Q}(\lambda) \otimes_{\mathbb{k}_T}(\mu).$$

If  $\lambda \in W_{\text{ext}} \cdot_p 0$  then  $\widehat{Q}(\lambda)$  belongs to  $\text{Rep}_\varnothing(G_1T)$ . By [Jan03, Proposition II.11.4], this module admits a filtration with subquotients of the form  $\widehat{Z}(\mu)$  with  $\mu \in W_{\text{ext}} \cdot_p 0$ ; moreover the number of occurrences of  $\widehat{Z}(\mu)$  does not depend on the choice of such a filtration, and is equal to the multiplicity  $[\widehat{Z}(\mu) : \widehat{\mathbb{L}}(\lambda)]$ . More generally, any projective object  $\widehat{Q}$  in  $\text{Rep}_\varnothing(G_1T)$  admits a filtration with subquotients of the form  $\widehat{Z}(\mu)$  with  $\mu \in W_{\text{ext}} \cdot_p 0$ , and the number of occurrences of  $\widehat{Z}(\mu)$  does not depend on the choice of filtration; this number will be denoted  $(\widehat{Q} : \widehat{Z}(\mu))$ .

LEMMA 5.6. — *For any  $\omega \in \Omega$ , the projective  $G_1T$ -module  $\widehat{Q}(t_\varsigma w_f \omega \cdot_p 0)$  is the image under (5.4) of  $\mathbb{T}(p\varsigma + \omega \cdot_p 0)$ . Moreover, for any  $\mu$  in  $\mathbf{X}$  we have*

$$\left( \widehat{Q}(t_\varsigma w_f \omega \cdot_p 0) : \widehat{Z}(\mu) \right) = \begin{cases} 1 & \text{if } \mu = t_\varsigma x \omega \cdot_p 0 \text{ for some } x \in W_f; \\ 0 & \text{otherwise.} \end{cases}$$

*Proof.* — The element  $t_\varsigma w_f \omega \cdot_p 0 = (t_\varsigma w_f \omega_\varsigma^{-1}) \cdot_p (\omega_\varsigma \omega \cdot_p 0)$  belongs to the unique alcove which contains  $(p-1)\varsigma = (t_\varsigma w_f \omega_\varsigma^{-1}) \cdot_p (\omega_\varsigma \cdot_p (-\varsigma))$  in its upper closure. Therefore, by [Jan03, Section II.11.10], the indecomposable projective  $G_1T$ -module  $\widehat{Q}(t_\varsigma w_f \omega \cdot_p 0)$  is obtained by translating from  $\omega_\varsigma \cdot_p (-\varsigma)$  to  $\omega_\varsigma \omega \cdot_p 0$  the  $G_1T$ -module  $\widehat{Q}((p-1)\varsigma)$ . Moreover, by [Jan03, Section II.9.16 and equation (1) in Section II.11.9] we have  $\widehat{Q}((p-1)\varsigma) \cong \widehat{\mathbb{L}}((p-1)\varsigma) \cong \widehat{Z}((p-1)\varsigma)$ . Then the description of  $(\widehat{Q}(t_\varsigma w_f \omega \cdot_p 0) : \widehat{Z}(\mu))$  follows from the considerations surrounding Equations (2)–(3) in [Jan03, Section II.9.22].

The first claim follows from the considerations above and Lemma 5.1.  $\square$



The following result is an easy consequence of [Jan80, Corollary 4.5] (see also [Jan03, Section 11.11]); see [Don93] or [Jan03, Section E.9] for details.

**THEOREM 5.7.** — *Assume that  $p \geq 2h - 2$ . Then for any restricted dominant weight  $\lambda$  we have*

$$\widehat{Q}(\lambda) \cong \mathbb{T}(2(p-1)\rho + w_f \lambda)_{|G_1 T}.$$

We will say that an element  $w \in W_{\text{ext}}$  is *restricted* if  $w \cdot_p 0$  is a restricted dominant weight. Note that this condition does not depend on  $p$ , and that restricted elements belong to  ${}^f W_{\text{ext}}$ .

In terms of the orbit  $W_{\text{ext}} \cdot_p 0$ , since  $w_f(\varsigma) = \varsigma - 2\rho$ , Theorem 5.7 implies in particular that (if  $p \geq 2h - 2$ ) for any  $w \in W_{\text{ext}}$  such that  $t_\varsigma w$  is restricted, we have

$$(5.6) \quad \widehat{Q}(t_\varsigma w \cdot_p 0) \cong \mathbb{T}(t_\varsigma w_f w \cdot_p 0)_{|G_1 T}.$$

#### 5.4. Characters of tilting modules as $G_1 T$ -modules

Now we set

$$\mathbf{M}_{\text{ext}}^{\text{sph}} := \mathbb{Z} \otimes_{\mathbb{Z}[v^{\pm 1}]} \mathcal{M}_{\text{ext}}^{\text{sph}},$$

and still denote by  $\varphi : \mathbf{M}_{\text{ext}}^{\text{sph}} \rightarrow \mathbf{M}_{\text{ext}}^{\text{asph}}$  and  $\zeta : \mathbf{M}_{\text{ext}}^{\text{sph}} \rightarrow \mathbb{Z}[W_{\text{ext}}]$  the (injective) morphisms induced by (3.6) and (3.1) respectively. We then consider the maps

$$\begin{array}{ccccc} \mathbf{M}_{\text{ext}}^{\text{sph}} & \xrightarrow{\varphi} & \mathbf{M}_{\text{ext}}^{\text{asph}} & \xrightarrow[(5.1)]{\sim} & [\text{Rep}_\emptyset(G)] \longrightarrow [\text{Rep}_\emptyset(G_1 T)] \\ \zeta \downarrow & & & & \\ & & \mathbb{Z}[W_{\text{ext}}] & & \end{array}$$

where the rightmost arrow is induced by the restriction functor (5.4).

**PROPOSITION 5.8.** — *Let  $M$  be a tilting module in  $\text{Rep}_\emptyset(G)$ , all of whose direct summands are of the form  $\mathbb{T}(t_\varsigma w \cdot_p 0)$  with  $w \in {}^f W_{\text{ext}}$ . Then  $M_{|G_1 T}$  is a projective  $G_1 T$ -module. Moreover, the inverse image  $a$  of  $[M]$  under (5.1) belongs to the image of  $\varphi$ , and the image under  $\zeta$  of the preimage of  $a$  is equal to*

$$\sum_{w \in W_{\text{ext}}} \left( M_{|G_1 T} : \widehat{Z}(w \cdot_p 0 + p\varsigma) \right) \cdot w.$$

*Proof.* — As explained in [Jan03, Lemma E.8], an indecomposable tilting module  $\mathbb{T}(\lambda)$  (with  $\lambda \in \mathbf{X}^+$ ) is projective as a  $G_1 T$ -module iff  $\lambda - (p-1)\varsigma \in \mathbf{X}^+$ . This implies the first claim in the proposition, and also that the  $G$ -modules  $M$  as in the statement are all isomorphic to direct sums of direct summands of modules of the form

$$\Theta_{s_1} \Theta_{s_2} \cdots \Theta_{s_n} (\mathbb{T}(p\varsigma + \omega \cdot_p 0))$$

with  $s_1, \dots, s_n$  in  $S$  and  $\omega \in \Omega$ . This reduces the proof of the proposition to the case of modules of this form. We will prove this case by induction on  $n$ .

First we treat the case  $n = 0$ . By (3.5) and Lemma 5.1 we have

$$[\mathbb{T}(p\varsigma + \omega \cdot_p 0)] = 1 \otimes \underline{N}_{t_\varsigma \omega} = \varphi(1 \otimes M_\omega).$$

(Of course, this equality is also a special case of Theorem 5.2.) Using (3.3), we deduce that the image under  $\zeta$  of the preimage of  $[\mathbb{T}(p\zeta + \omega \cdot_p 0)]$  is

$$\sum_{x \in W_{\mathfrak{f}}} 1 \otimes H_{x\omega}.$$

On the other hand, by Lemma 5.6 we know that  $\mathbb{T}(p\zeta + \omega \cdot_p 0)_{|G_1T} = \widehat{Q}(t_{\zeta}w_{\mathfrak{f}}\omega \cdot_p 0)$ , and we know the multiplicities of baby Verma modules in this projective module. Comparing with the formula above, we deduce the desired claim.

To prove the induction step, we will prove that if the claim is true for a module  $M$ , then it is true also for  $\Theta_s(M)$  for any  $s \in S$ . As explained just after (5.1), if we denote by  $a$  the inverse image of  $[M]$ , then the inverse image of  $[\Theta_s(M)]$  is  $a \cdot (\text{id} + s)$ . Hence if  $a = \varphi(b)$ , then this inverse image is  $\varphi(a \cdot (\text{id} + s))$ . Now by Lemma 5.4, for any  $w \in W_{\text{ext}}$  we have

$$\begin{aligned} \left( \Theta_s \left( M_{|G_1T} \right) : \widehat{Z}(p\zeta + w \cdot_p 0) \right) \\ = \left( M_{|G_1T} : \widehat{Z}(p\zeta + w \cdot_p 0) \right) + \left( M_{|G_1T} : \widehat{Z}(p\zeta + ws \cdot_p 0) \right), \end{aligned}$$

and the desired claim follows.  $\square$

### 5.5. The simple character formula

Our main application of Theorem 3.5 is the following claim.

**THEOREM 5.9.** — *Assume that  $p \geq 2h - 2$ . If  $w \in W_{\text{ext}}$  is such that  $t_{\zeta}w$  belongs to  ${}^{\mathfrak{f}}W_{\text{ext}}$  and is restricted, then for any  $y \in W_{\text{ext}}$  we have*

$$\left( \widehat{Q}(w \cdot_p 0) : \widehat{Z}(y \cdot_p 0) \right) = {}^p h_{y,w}(1).$$

*Proof.* — By (5.6), we have

$$\mathbb{T}(t_{\zeta}w_{\mathfrak{f}}w \cdot_p 0)_{|G_1T} \cong \widehat{Q}(p\zeta + w \cdot_p 0).$$

By Theorem 5.2, the class  $[\mathbb{T}(t_{\zeta}w_{\mathfrak{f}}w \cdot_p 0)]$  in  $[\text{Rep}_0(G)]$  is the image of  $1 \otimes {}^p \underline{N}_{t_{\zeta}w_{\mathfrak{f}}w}$  under (5.1). Now by Theorem 3.5 we have

$$1 \otimes {}^p \underline{N}_{t_{\zeta}w_{\mathfrak{f}}w} = \varphi \left( 1 \otimes {}^p \underline{M}_{w_{\mathfrak{f}}w} \right).$$

Using (3.2) and Proposition 5.8 we deduce that

$$\sum_{x \in W_{\text{ext}}} \left( \widehat{Q}(p\zeta + w \cdot_p 0) : \widehat{Z}(p\zeta + x \cdot_p 0) \right) \cdot x = 1 \otimes {}^p \underline{H}_w.$$

Since  $(\widehat{Q}(p\zeta + w \cdot_p 0) : \widehat{Z}(p\zeta + x \cdot_p 0)) = (\widehat{Q}(w \cdot_p 0) : \widehat{Z}(x \cdot_p 0))$  for any  $x \in W_{\text{ext}}$  (see (5.3) and (5.5)), this implies the desired equality.  $\square$

**Remark 5.10.** —

- (1) Let  $w_{\max} \in W$  be the unique element such that  $w_{\max} \cdot_p 0$  belongs to the (shifted and dilated) alcove of  $-p\zeta$ . The main result of [Fie11] (see in particular [Fie11,

Theorems 7.8 and 8.6]) states that if  ${}^p H_x = \underline{H}_x$  for any  $x \in W$  such that  $x \leq w_{\max}$  in the Bruhat order, then we have

$$[\widehat{Z}(y \cdot_p 0) : \widehat{\mathbb{L}}(w \cdot_p 0)] = h_{y,w}(1)$$

for any  $w, y \in W$  such that  $t_\zeta w$  belongs to  ${}^f W_{\text{ext}}$  and is restricted (and hence Lusztig's conjecture holds). Of course, this claim also follows from Theorem 5.9.

- (2) Once the multiplicities  $(\widehat{Q}(w \cdot_p 0) : \widehat{Z}(y \cdot_p 0))$  are known for any  $w, y$  as in Theorem 5.9, using (5.3) and (5.5) we can deduce these multiplicities for any  $w, y \in W$ . Then, using the reciprocity formula

$$(\widehat{Q}(w \cdot_p 0) : \widehat{Z}(y \cdot_p 0)) = [\widehat{Z}(y \cdot_p 0) : \widehat{\mathbb{L}}(w \cdot_p 0)]$$

(see Section 5.3) one can deduce the multiplicities on the right-hand side of this equality. And this information allows to compute the characters of the modules  $\widehat{\mathbb{L}}(w \cdot_p 0)$  for any  $w \in W_{\text{ext}}$ . In fact, using (5.2) it suffices to do so when  $w$  is restricted. In this case  $\widehat{\mathbb{L}}(w \cdot_p 0)$  is the restriction of a  $G$ -module; hence its weights (and their multiplicities) are stable under  $W_f$ . As a consequence, to determine them it suffices to compute the dominant weights appearing in  $\widehat{\mathbb{L}}(w \cdot_p 0)$  and their multiplicities. Using the determination of the multiplicities  $[\widehat{Z}(y \cdot_p 0) : \widehat{\mathbb{L}}(w \cdot_p 0)]$  and the “triangularity” of these numbers (see [Jan03, Corollary 9.15(a)]) one can write

$$(5.7) \quad [\widehat{\mathbb{L}}(w \cdot_p 0)] = \sum_{x \in W_{\text{ext}}^{(1)}} a_x \cdot [\widehat{Z}(x \cdot_p 0)] + \sum_{y \in W_{\text{ext}}^{(2)}} b_y \cdot [\widehat{\mathbb{L}}(y \cdot_p 0)]$$

with  $W_{\text{ext}}^{(1)}, W_{\text{ext}}^{(2)}$  subsets of  $W_{\text{ext}}$ , in such a way that  $\widehat{\mathbb{L}}(y \cdot_p 0)$  does not admit any dominant weight for  $y \in W_{\text{ext}}^{(2)}$ . The characters of the baby Verma modules are easy to compute, see e.g. [Fie10, Section 3.1]. Hence from (5.7) one can compute the dominant weights appearing in  $\widehat{\mathbb{L}}(w \cdot_p 0)$  and their multiplicities. (See also [Sob20, Section 4] for a different presentation of this procedure.)

- (3) Our assumptions on  $p$  in Theorem 5.9 are that  $p \geq 2h - 2$  and  $p > h$ . It is easily seen that these two conditions are equivalent to the condition that  $p \geq 2h - 1$ .

## 5.6. Proof of Theorem 1.2

We conclude the paper by explaining how Theorem 5.9 implies Theorem 1.2 from the introduction. As in Section 1.4, for  $m$  in  $\mathcal{P}$  we denote by  $[m]_{v \mapsto 1}$  its image in  $\mathbb{Z} \otimes_{\mathbb{Z}[v \pm 1]} \mathcal{P} \cong \mathbb{Z}[\mathcal{A}]$ . Recall that  $W$  acts on  $\mathcal{A}$  on the right, see Section 1.5. This induces in the natural way a structure of right  $\mathbb{Z}[W]$ -module on  $\mathbb{Z}[\mathcal{A}]$ , and it is clear from (2.5) that this action coincides with the one induced by the  $\mathcal{H}$ -action on  $\mathcal{P}$  (via the canonical isomorphism  $\mathbb{Z} \otimes_{\mathbb{Z}[v \pm 1]} \mathcal{H} \cong \mathbb{Z}[W]$ ).

*Proof of Theorem 1.2.* — Since both  $q_A$  and  ${}^p P_{\bar{A}}$  are invariant under the replacement of  $A$  by  $\mu + A$  for  $\mu \in \mathbf{X}$  (see (5.5) and (2.9) respectively), we can assume that

$A \in \check{\Pi}_0$ , so that  $\hat{A} = w_f(A) \in \hat{\Pi}_0$ . Then

$${}^p P_{\hat{A}} = \frac{1}{\pi_f} P_{A_{\text{fund}}} \cdot \zeta({}^p M_w),$$

where  $w \in W$  is the unique element such that  $w_f(A) = A_{\text{fund}} \cdot w = w(A_{\text{fund}})$ , i.e. such that  $A = w_f w(A_{\text{fund}})$ . By construction we have

$$\zeta({}^p M_w) = {}^p H_{w_f w} = \sum_{y \in W} {}^p h_{y, w_f w} \cdot H_y,$$

hence using (2.8) we obtain that

$$\begin{aligned} [{}^p P_{\hat{A}}]_{v \mapsto 1} &= \frac{1}{|W_f|} \left( \sum_{x \in W_f} x(A_{\text{fund}}) \right) \cdot \left( \sum_{y \in W} {}^p h_{y, w_f w}(1) \cdot y \right) \\ &= \frac{1}{|W_f|} \left( \sum_{\substack{x \in W_f \\ y \in W}} {}^p h_{y, w_f w}(1) \cdot xy(A_{\text{fund}}) \right) \\ &= \sum_{z \in W} \frac{1}{|W_f|} \left( \sum_{x \in W_f} {}^p h_{x^{-1}z, w_f w}(1) \right) \cdot z(A_{\text{fund}}). \end{aligned}$$

Now since  $w \in {}^f W$ , the element  $w_f w$  is maximal in  $W_f \cdot w_f w$ , so that the parity complex  $\mathcal{E}_{w_f w}$  (see Section 4.1) is constructible with respect to the stratification by  $G_{\mathcal{O}}^\vee$ -orbits, which implies that  ${}^p h_{x^{-1}z, w_f w}(1) = {}^p h_{z, w_f w}(1)$  for any  $x \in W_f$ . We deduce that

$$[{}^p P_{\hat{A}}]_{v \mapsto 1} = \sum_{z \in W} {}^p h_{z, w_f w}(1) \cdot z(A_{\text{fund}}).$$

Comparing with Theorem 5.9 and the definition of  $q_A$ , we obtain the desired formula.  $\square$

## 6. A combinatorial proof of Theorem 1.4 in the case of Kazhdan–Lusztig bases

In this section we provide an alternative proof of the version of Theorem 1.4 for “standard” Kazhdan–Lusztig bases, see Remark 1.5(2). This proof is based on the results of [Soe97]<sup>(4)</sup> (and is therefore “combinatorial”), and was explained to us by Soergel.

In this section we assume that  $G$  is semisimple (and simply connected). As in Section 1.4 we consider the “periodic module”  $\mathcal{P}$  for  $\mathcal{H}$ . As explained in [Soe97, Section 4] the family  $(P_A : A \in \mathcal{A})$  forms a  $\mathbb{Z}[v^{\pm 1}]$ -basis of a certain  $\mathcal{H}$ -submodule  $\mathcal{P}^\circ \subset \mathcal{P}$ . By [Soe97, Lemma 4.9], the action of  $\mathbf{X}$  on  $\mathcal{P}$  considered in Section 2.3 extends to a  $\mathbb{Z}[v^{\pm 1}]$ -linear action of the extended affine Weyl group  $W_{\text{ext}}$  on  $\mathcal{P}^\circ$ , for

<sup>(4)</sup>Some of the results of [Soe97] on which this proof is based are not originally due to Soergel, but are restatements of results due to Lusztig and to Kato. We refer to [Soe97] for a discussion of the original references.

which the action of  $w \in W_{\text{ext}}$  is denoted  $\langle w \rangle : \mathcal{P}^\circ \rightarrow \mathcal{P}^\circ$ . It follows from (2.6) that this action satisfies the following formula: for  $\omega \in \Omega$  and  $w \in W$  we have

$$\langle \omega w \rangle (P \cdot h) = (\langle \omega w \rangle P) \cdot (H_\omega h H_\omega^{-1})$$

for  $P \in \mathcal{P}^\circ$  and  $h \in \mathcal{H}$ . (Here the element  $H_\omega$  belongs to the larger algebra  $\mathcal{H}_{\text{ext}}$  introduced in Section 3.1; conjugation by this element stabilizes  $\mathcal{H}$ .)

We next consider the map

$$\text{alt} : \mathcal{P}^\circ \rightarrow \mathcal{P}^\circ$$

defined by the formula

$$\text{alt}(P) = \sum_{w \in W_f} (-1)^{\ell(w)} \langle w \rangle P.$$

We will also consider the  $\mathbb{Z}[v^{\pm 1}]$ -linear map

$$\text{res} : \mathcal{P} \rightarrow \mathcal{M}^{\text{asph}}$$

determined by

$$\text{res}(A) = \begin{cases} N_A & \text{if } A \in \mathcal{A}^+; \\ 0 & \text{otherwise.} \end{cases}$$

This morphism is not  $\mathcal{H}$ -linear; but it follows from [Soe97, Proposition 5.2] that the composition  $\text{res} \circ \text{alt}$  is  $\mathcal{H}$ -linear. Moreover, [Soe97, Theorem 5.3(1)] says that for  $A \in \mathcal{A}_\rho^+$  we have

$$(6.1) \quad \text{res} \circ \text{alt}(\underline{P}_A) = \underline{N}_A.$$

On the other hand, let us consider the morphism of right  $\mathcal{H}$ -modules

$$\text{Alt} := \langle t_{-\rho} \rangle \circ \text{alt} \circ \langle t_\rho \rangle : \mathcal{P}^\circ \rightarrow \mathcal{P}^\circ.$$

In [Soe97, Section 6] Soergel introduces the  $\mathbb{Z}[v^{\pm 1}]$ -module  $\hat{\mathcal{P}}$  consisting of certain formal linear combinations  $\sum_A f_A A$ , and the map  $\eta : \hat{\mathcal{P}} \rightarrow \hat{\mathcal{P}}$ . The module  $\hat{\mathcal{P}}$  contains  $\mathcal{P}$  as a submodule in the natural way. We also have a map

$$\text{Res} : \hat{\mathcal{P}} \rightarrow \mathcal{M}^{\text{sph}}$$

sending  $\sum_A f_A A$  to  $\sum_{A \in \mathcal{A}^+} f_A M_A$ . (Here the linear combination  $\sum_{A \in \mathcal{A}^+} f_A M_A$  is finite due to the form of the combinations authorized in  $\hat{\mathcal{P}}$ .) Then [Soe97, Corollary 6.9] states that for any  $A \in \mathcal{A}^+$  we have

$$(6.2) \quad \underline{M}_A = \text{Res} \circ \eta \circ \text{Alt}(\underline{P}_A).$$

By [Soe97, Proposition 6.6], the map  $\text{Res} \circ \eta \circ \text{Alt}$  is  $\mathcal{H}$ -linear; it follows that  $\text{Res} \circ \eta : \text{Alt}(\mathcal{P}^\circ) \rightarrow \mathcal{M}^{\text{sph}}$  is  $\mathcal{H}$ -linear as well. One can check that the elements  $(\text{Alt}(\underline{P}_A) : A \in \mathcal{A}^+)$  form a  $\mathbb{Z}[v^{\pm 1}]$ -basis of the sub- $\mathcal{H}$ -module  $\text{Alt}(\mathcal{P}^\circ) \subset \mathcal{P}^\circ$ ; therefore this formula implies that  $\text{Res} \circ \eta$  induces an isomorphism of  $\mathcal{H}$ -modules from  $\text{Alt}(\mathcal{P}^\circ)$  to  $\mathcal{M}^{\text{sph}}$ .

By (2.7) we have  $\langle t_\rho \rangle(\underline{P}_A) = \underline{P}_{\rho+A}$  for any  $A \in \mathcal{A}$ ; hence the formula (6.2) can be written as

$$\underline{M}_A = \text{Res} \circ \eta \circ \langle t_{-\rho} \rangle \circ \text{alt}(\underline{P}_{\rho+A})$$

for any  $A \in \mathcal{A}^+$ .

Fix now  $A \in \mathcal{A}^+$ , and choose  $h \in \mathcal{H}$  such that  $\underline{M}_A = \underline{M}_{A_{\text{fund}}} \cdot h$ . Then we have

$$\begin{aligned} \text{Res} \circ \eta \circ \langle t_{-\rho} \rangle \circ \text{alt} \left( \underline{P}_{\rho+A} \right) &= \underline{M}_A = \underline{M}_{A_{\text{fund}}} \cdot h \\ &= \left( \text{Res} \circ \eta \circ \langle t_{-\rho} \rangle \circ \text{alt} \left( \underline{P}_{\rho+A_{\text{fund}}} \right) \right) \cdot h \\ &= \text{Res} \circ \eta \circ \langle t_{-\rho} \rangle \circ \text{alt} \left( \underline{P}_{\rho+A_{\text{fund}}} \cdot H_{\omega_\rho} h H_{\omega_\rho}^{-1} \right). \end{aligned}$$

By injectivity of  $\text{Res} \circ \eta$  on  $\text{Alt}(\mathcal{P}^\circ)$  we deduce that

$$\langle t_{-\rho} \rangle \circ \text{alt} \left( \underline{P}_{\rho+A} \right) = \langle t_{-\rho} \rangle \circ \text{alt} \left( \underline{P}_{\rho+A_{\text{fund}}} \cdot H_{\omega_\rho} h H_{\omega_\rho}^{-1} \right),$$

and then that

$$\text{alt} \left( \underline{P}_{\rho+A} \right) = \text{alt} \left( \underline{P}_{\rho+A_{\text{fund}}} \cdot H_{\omega_\rho} h H_{\omega_\rho}^{-1} \right).$$

Applying  $\text{res}$  and using (6.1), we deduce that

$$\begin{aligned} (6.3) \quad \underline{N}_{\rho+A} &= \text{res} \circ \text{alt} \left( \underline{P}_{\rho+A} \right) = \text{res} \circ \text{alt} \left( \underline{P}_{\rho+A_{\text{fund}}} \cdot H_{\omega_\rho} h H_{\omega_\rho}^{-1} \right) \\ &= \left( \text{res} \circ \text{alt} \left( \underline{P}_{\rho+A_{\text{fund}}} \right) \right) \cdot H_{\omega_\rho} h H_{\omega_\rho}^{-1} = \underline{N}_{\rho+A_{\text{fund}}} \cdot H_{\omega_\rho} h H_{\omega_\rho}^{-1}. \end{aligned}$$

Now we have  $S_\rho = \tau_\rho(S_f) = \omega_\rho S_f \omega_\rho$ , see Remark 2.1. Hence there exists an isomorphism of  $\mathbb{Z}[v^{\pm 1}]$ -modules

$$\mathcal{M}^{\text{sph}} \xrightarrow{\sim} \mathcal{M}_{\bar{\rho}}^{\text{sph}}$$

which sends  $M_{\text{id}} \cdot h$  to  $M_{\text{id}}^{\bar{\rho}} \cdot H_{\omega_\rho} h H_{\omega_\rho}^{-1}$  for any  $h \in \mathcal{H}$ . In terms of the parametrization by alcoves, this morphism sends  $M_B$  to  $M_{\rho+B}^\rho$  for any  $B \in \mathcal{A}^+$ . (In fact, if  $B = w(A_{\text{fund}})$  with  $w \in {}^f W$ , then  $M_B = M_{A_{\text{fund}}} \cdot H_w$  is sent to

$$M_{\rho+A_{\text{fund}}}^\rho \cdot H_{\omega_\rho} H_w H_{\omega_\rho}^{-1} = M_{(\rho+A_{\text{fund}}) \cdot \omega_\rho w \omega_\rho^{-1}}^\rho = M_{x_\rho \omega_\rho w \omega_\rho^{-1}(A_{\text{fund}})}^\rho = M_{\rho+B}^\rho)$$

This morphism also commutes with the appropriate Kazhdan–Lusztig involutions, hence sends  $\underline{M}_B$  to  $\underline{M}_{\rho+B}^\rho$  for any  $B \in \mathcal{A}^+$ . It follows that

$$M_{\rho+A_{\text{fund}}}^\rho \cdot H_{\omega_\rho} h H_{\omega_\rho}^{-1} = \underline{M}_{\rho+A}^\rho,$$

which proves that

$$\underline{N}_{\rho+A_{\text{fund}}} \cdot H_{\omega_\rho} h H_{\omega_\rho}^{-1} = \varphi_\rho \left( \underline{M}_{\rho+A}^\rho \right).$$

Comparing with (6.3) we finally obtain that  $\underline{N}_{\rho+A} = \varphi_\rho(\underline{M}_{\rho+A}^\rho)$ , which finishes the proof of Theorem 1.4.

## BIBLIOGRAPHY

- [ACR18] Pramod N. Achar, Nicolas Cooney, and Simon Riche, *The parabolic exotic  $t$ -structure*, Épijournal de Géom. Algèbr., EPIGA **2** (2018), article no. 8 (31 pages). ↑519, 521
- [AJS94] Henning H. Andersen, Jens C. Jantzen, and Wolfgang Soergel, *Representations of quantum groups at a  $p$ -th root of unity and of semisimple groups in characteristic  $p$ : independence of  $p$* , Astérisque, vol. 220, Société Mathématique de France, 1994. ↑504
- [AMRW19] Pramod N. Achar, Shotaro Makisumi, Simon Riche, and Geordie Williamson, *Koszul duality for Kac–Moody groups and characters of tilting modules*, J. Am. Math. Soc. **32** (2019), no. 1, 261–310. ↑504, 505, 509, 510, 524

- [And98] Henning H. Andersen, *Tilting modules for algebraic groups*, Algebraic groups and their representations (Cambridge, 1997), NATO ASI Series. Series C. Mathematical and Physical Sciences, vol. 517, Kluwer Academic Publishers, 1998, pp. 25–42. ↑509
- [AR19] Pramod N. Achar and Simon Riche, *Dualité de Koszul formelle et théorie des représentations modulaire des groupes algébriques réductifs*, SMF 2018 : Congrès de la Société Mathématique de France, Séminaires et Congrès, vol. 33, Société Mathématique de France, 2019, pp. 83–172. ↑514, 524
- [BBM04] Roman Bezrukavnikov, Alexander Braverman, and Ivan Mirković, *Some results about geometric Whittaker model*, Adv. Math. **186** (2004), no. 1, 143–152. ↑508
- [BGM<sup>+</sup>19] Roman Bezrukavnikov, Dennis Gaitsgory, Ivan Mirković, Simon Riche, and Laura Rider, *An Iwahori–Whittaker model for the Satake category*, J. Éc. Polytech., Math. **6** (2019), 707–735. ↑508, 509, 520, 522
- [BNPS20] Christopher P. Bendel, Daniel K. Nakano, Cornelius Pillen, and Paul Sobaje, *Counterexamples to the tilting and  $(p, r)$ -filtration conjectures*, J. Reine Angew. Math. **767** (2020), 193–202. ↑509
- [BY13] Roman Bezrukavnikov and Zhiwei Yun, *On Koszul duality for Kac–Moody groups*, Represent. Theory **17** (2013), 1–98. ↑508
- [Don93] Stephen Donkin, *On tilting modules for algebraic groups*, Math. Z. **212** (1993), no. 1, 39–60. ↑504, 509, 527
- [Fie10] Peter Fiebig, *Lusztig’s conjecture as a moment graph problem*, Bull. Lond. Math. Soc. **42** (2010), no. 6, 957–972. ↑506, 529
- [Fie11] ———, *Sheaves on affine Schubert varieties, modular representations, and Lusztig’s conjecture*, J. Am. Math. Soc. **24** (2011), no. 1, 133–181. ↑528, 529
- [Fie12] ———, *An upper bound on the exceptional characteristics for Lusztig’s character formula*, J. Reine Angew. Math. **673** (2012), 1–31. ↑504
- [Hum71] James E. Humphreys, *Modular representations of classical Lie algebras and semisimple groups*, J. Algebra **19** (1971), 51–79. ↑506
- [IM65] Nagayoshi Iwahori and Hideya Matsumoto, *On some Bruhat decomposition and the structure of the Hecke rings of  $p$ -adic Chevalley groups*, Publ. Math., Inst. Hautes Étud. Sci. **25** (1965), 5–48. ↑510, 511
- [Jan80] Jens C. Jantzen, *Darstellungen halbeinfacher Gruppen und ihrer Frobenius–Kerne*, J. Reine Angew. Math. **317** (1980), 157–199. ↑509, 527
- [Jan03] ———, *Representations of algebraic groups*, second ed., Mathematical Surveys and Monographs, vol. 107, American Mathematical Society, 2003. ↑511, 524, 525, 526, 527, 529
- [JMW14] Daniel Juteau, Carl Mautner, and Geordie Williamson, *Parity sheaves*, J. Am. Math. Soc. **27** (2014), no. 4, 1169–1212. ↑508, 518, 519, 521
- [JMW16] ———, *Parity sheaves and tilting modules*, Ann. Sci. Éc. Norm. Supér. **49** (2016), no. 2, 257–275. ↑520
- [JW17] Lars T. Jensen and Geordie Williamson, *The  $p$ -canonical basis for Hecke algebras*, Categorification and higher representation theory (Anna Beliakova et al., eds.), Contemporary Mathematics, vol. 683, American Mathematical Society, 2017, pp. 333–361. ↑507, 514, 518
- [KL93a] David Kazhdan and George Lusztig, *Tensor structures arising from affine Lie algebras I*, J. Am. Math. Soc. **6** (1993), no. 4, 905–947. ↑504
- [KL93b] ———, *Tensor structures arising from affine Lie algebras II*, J. Am. Math. Soc. **6** (1993), no. 4, 949–1011. ↑504
- [KL94a] ———, *Tensor structures arising from affine Lie algebras III*, J. Am. Math. Soc. **7** (1994), no. 2, 335–381. ↑504

- [KL94b] ———, *Tensor structures arising from affine Lie algebras IV*, J. Am. Math. Soc. **7** (1994), no. 2, 383–453. ↑504
- [KT95] Masaki Kashiwara and Toshiyuki Tanisaki, *Kazhdan–Lusztig conjecture for affine Lie algebras with negative level*, Duke Math. J. **77** (1995), no. 1, 21–62. ↑504
- [KT96] ———, *Kazhdan–Lusztig conjecture for affine Lie algebras with negative level II: Non-integral case*, Duke Math. J. **84** (1996), no. 3, 771–813. ↑504
- [Lus80a] George Lusztig, *Hecke algebras and Jantzen’s generic decomposition patterns*, Adv. Math. **37** (1980), 121–164. ↑504, 506, 514
- [Lus80b] ———, *Some problems in the representation theory of finite Chevalley groups*, The Santa Cruz Conference on Finite Groups, Proceedings of Symposia in Pure Mathematics, vol. 37, American Mathematical Society, 1980, pp. 313–317. ↑506
- [Lus94] ———, *Monodromic systems on affine flag manifolds*, Proc. R. Soc. Lond. **445** (1994), no. 1923, 231–246, Errata in *Proceedings of the Royal Society of London. A. Mathematical, Physical and Engineering Sciences* vol. 450, (1995), 731–732. ↑504
- [MR18] Carl Mautner and Simon Riche, *Exotic tilting sheaves, parity sheaves on affine Grassmannians, and the Mirković–Vilonen conjecture*, J. Eur. Math. Soc. **20** (2018), no. 9, 2259–2332. ↑519, 520
- [Ric] Simon Riche, *Geometric Representation Theory in positive characteristic*, habilitation thesis, available at <https://tel.archives-ouvertes.fr/tel-01431526>. ↑520
- [RW18] Simon Riche and Geordie Williamson, *Tilting modules and the  $p$ -canonical basis*, Astérisque, vol. 397, Société Mathématique de France, 2018. ↑505, 507, 514, 516, 518, 519, 523, 524
- [RW20] ———, *Smith–Treumann theory and the linkage principle*, <https://arxiv.org/abs/2003.08522>, 2020. ↑504, 510
- [Sob20] Paul Sobaje, *On character formulas for simple and tilting modules*, Adv. Math. **369** (2020), article no. 107172. ↑505, 509, 529
- [Soe97] Wolfgang Soergel, *Kazhdan–Lusztig polynomials and a combinatorics for tilting modules*, Represent. Theory **1** (1997), 83–114. ↑506, 507, 508, 512, 513, 514, 516, 517, 530, 531
- [Spr82] Tonny A. Springer, *Quelques applications de la cohomologie d’intersection*, Séminaire N. Bourbaki, Vol. 1981/82, Astérisque, vol. 92–93, Springer, 1982, pp. 249–273. ↑518
- [Wil17] Geordie Williamson, *Schubert calculus and torsion explosion*, J. Am. Math. Soc. **30** (2017), no. 4, 1023–1046. ↑504

Manuscript received on 30th April 2019,  
revised on 11th May 2020,  
accepted on 8th July 2020.

Recommended by Editor C. Sorger.  
Published under license CC BY 4.0.



This journal is a member of Centre Mersenne.



Simon RICHE  
Université Clermont Auvergne,  
CNRS, LMBP,



F-63000 Clermont-Ferrand, (France)  
simon.riche@uca.fr

Geordie WILLIAMSON  
School of Mathematics and Statistics F07,  
University of Sydney  
NSW 2006, (Australia)  
g.williamson@sydney.edu.au