



ANNALES  
HENRI LEBESGUE

---

JULIEN MARCHÉ

---

CHRISTOPHER-LLOYD SIMON

---

# AUTOMORPHISMS OF CHARACTER VARIETIES

## AUTOMORPHISMES DES VARIÉTÉS DE CARACTÈRES

---

**ABSTRACT.** — We show that the algebraic automorphism group of the  $SL_2(\mathbb{C})$  character variety of a closed orientable surface with negative Euler characteristic is a finite extension of its mapping class group. Along the way, we provide a simple characterization of the valuations on the character algebra coming from measured laminations.

**RÉSUMÉ.** — Nous montrons que le groupe d'automorphismes de la variété des  $SL_2(\mathbb{C})$  caractères d'une surface orientable close de caractéristique d'Euler strictement négative est une extension finie de son groupe modulaire. En cours de route, nous donnons une caractérisation simple des valuations de l'algèbre des fonctions sur la variété des caractères qui proviennent des laminations mesurées.

## 1. Introduction

Let  $\Sigma$  be a closed oriented surface of genus  $g \geq 2$  and denote  $\text{Mod}(\Sigma)$  its mapping class group, which by the Dehn–Nielsen–Baer theorem can be identified with  $\text{Out}(\pi_1(\Sigma)) = \text{Aut}(\pi_1(\Sigma))/\text{Inn}(\pi_1(\Sigma))$ .

---

*Keywords:* mapping class group, character variety, measured lamination.

*2020 Mathematics Subject Classification:* 57K20.

*DOI:* <https://doi.org/10.5802/ahl.82>

The meaningful spaces which carry an action of this group often manifest rigidity properties, as explained for instance in [AS16]: whenever the mapping class group acts preserving some structure, it is almost the full automorphism group; celebrated examples are the Teichmüller space with its Kähler structure, the space of measured laminations with its PL structure, or the curve complex with its simplicial structure.

Here we show the same kind of result for the character variety, that is the space  $X(\Sigma)$  defined as the algebraic quotient of  $\text{Hom}(\pi_1(\Sigma), \text{SL}_2(\mathbb{C}))$  by the conjugation action of  $\text{SL}_2(\mathbb{C})$ . Recall that this affine variety is constructed from its algebra of functions

$$\mathbb{C}[X(\Sigma)] = \mathbb{C}[\text{Hom}(\pi_1(\Sigma), \text{SL}_2(\mathbb{C}))]^{\text{SL}_2(\mathbb{C})}.$$

The automorphism group  $\text{Aut}(X(\Sigma))$  of the affine variety  $X(\Sigma)$  is by definition the group of automorphisms of the  $\mathbb{C}$ -algebra  $\mathbb{C}[X(\Sigma)]$ .

The action  $[\varphi] \cdot [\rho] = [\rho \circ \varphi^{-1}]$  of  $\varphi \in \text{Aut}(\pi_1(\Sigma))$  on  $\rho \in \text{Hom}(\pi_1(\Sigma), \text{SL}_2(\mathbb{C}))$  descends to an algebraic action of the mapping class group on the character variety. Moreover, multiplying all representations  $\rho: \pi_1(\Sigma) \rightarrow \text{SL}_2(\mathbb{C})$  by a central one  $\lambda: \pi_1(\Sigma) \rightarrow \{\pm \text{Id}\}$  yields an action of  $H^1(\Sigma, \mathbb{Z}/2\mathbb{Z})$  on  $X(\Sigma)$ . The purpose of this note is to prove the:

**THEOREM.** — *When  $g \geq 3$ , the so defined map  $H^1(\Sigma, \mathbb{Z}/2\mathbb{Z}) \times \text{Mod}(\Sigma) \rightarrow \text{Aut}(X(\Sigma))$  is an isomorphism. When  $g = 2$  it is surjective with kernel the hyperelliptic involution.*

Similar results were obtained in [ÈH74] for the 1-punctured torus and 4-punctured sphere. We thank Serge Cantat for pointing this out and explaining the proof. The computation of the automorphism group of the character variety had been asked on many occasions to the first author by Juan Souto; however this work stemmed from different motivations, including a better understanding of measured laminations in terms of valuations.

Our strategy is to find an action of  $\text{Aut}(X(\Sigma))$  on the space of measured laminations, and show that it preserves the set of simple curves and their disjointness relation, hence the curve complex. For this consider the action of  $\text{Aut}(X(\Sigma))$  on a space  $\mathcal{V}$  of valuations. Those are by definition the functions  $v: \mathbb{C}[X(\Sigma)] \rightarrow \{-\infty\} \cup [0, +\infty)$  which are null on  $\mathbb{C}^*$ , take finite values except for  $v(0) = -\infty$ , and satisfy for all  $f, g$  the relations  $v(fg) = v(f) + v(g)$  and  $v(f + g) \leq \max(v(f), v(g))$ . We endow  $\mathcal{V}$  with the topology given by pointwise convergence, for which the action of  $\text{Aut}(X(\Sigma))$  is continuous.

Although it is established (for instance in [Ota12]) that the space of measured laminations  $\text{ML}(\Sigma)$  embeds continuously in  $\mathcal{V}$ , it is not clear why this subset should be preserved by  $\text{Aut}(X(\Sigma))$ , and this is one of the main steps in the proof. Corollary 3.2 gives a simple and presumably new characterization of measured laminations as the set of “simple valuations” which is analogous to the set of monomial valuations for polynomial algebras. We hope that this description will help understand the topology of the set of all valuations over the character variety, for instance by showing that it retracts by deformation on the space of simple valuations.

Section 2 is quite elementary: we define simple valuations and show that they contain an embedded copy of the space measured laminations. Section 3 is the

technical heart of the note: we show that any valuation is (sharply) dominated by one which is associated to a measured lamination, for this we apply the Morgan–Otal and Skora theorems to some Bass–Serre tree. Then we identify a property of valuations, called untameability, which is obviously preserved by algebraic automorphism and we show that it defines a dense subset in ML to conclude that  $\text{Aut}(X(\Sigma))$  acts on  $\text{ML}(\Sigma)$ . In Section 4, after identifying multicurves with simple and discrete valuations, we explain how to read the number of components and their geometric intersection in terms of their associated valuation rings. We deduce that  $\text{Aut}(X(\Sigma))$  acts on the curve complex, and conclude using Ivanov’s Theorem.

### Acknowledgements

We wish to thank Javier Aramayona, Serge Cantat, Marco Maculan, Mirko Mauri, Athanase Papadopoulos, Juan Souto and Alex Wright for their implication in this work. We thank in particular Christopher Leininger for his help in the proof of Lemma 2.8 and Lemma 3.4. The referee also helped to bring the article to its present form through his criticism.

## 2. Simple valuations and measured laminations

For  $\alpha \in \pi_1(\Sigma)$ , define  $t_\alpha \in \mathbb{C}[X(\Sigma)]$  by the formula  $t_\alpha([\rho]) = \text{Tr } \rho(\alpha)$ . Let us recall a folklore presentation of the character variety, mainly due to Procesi. We refer to [P17] for a full account and to [Che13, Prop. 2.3] for a proof of this specific statement, which holds more generally for any finitely generated group.

**THEOREM** (Algebra presentation). — *The algebra  $\mathbb{C}[X(\Sigma)]$  is generated by the  $t_\alpha$  for  $\alpha \in \pi_1(\Sigma)$  with ideal of relations generated by  $t_1 - 2$  and  $t_\alpha t_\beta - t_{\alpha\beta} - t_{\alpha\beta^{-1}}$  for  $\alpha, \beta \in \pi_1(\Sigma)$ .*

We call *multicurve* on  $\Sigma$  an embedded one dimensional submanifold  $\Gamma \subset \Sigma$  which is a union of curves homotopic to  $\gamma_i \in \pi_1(\Sigma) \setminus \{1\}$ , and set  $t_\Gamma = \prod_{i=1}^n t_{\gamma_i} \in \mathbb{C}[X(\Sigma)]$ . Components of  $\Gamma$  must be simple and disjoint; we allow the empty multicurve for which  $t_\emptyset = 1$ . The previous theorem has the following important consequence [PS00]:

**THEOREM** (Linear basis for the character algebra). — *The family  $(t_\Gamma)$  where  $\Gamma$  ranges over the isotopy classes of multicurves forms a linear basis of  $\mathbb{C}[X(\Sigma)]$ .*

This allows to treat the elements  $t_\Gamma$  as if they were monomials (despite the fact that they are not stable by multiplication), and motivates the next definition.

**DEFINITION 2.1** (Simple valuations). — *A valuation  $v \in \mathcal{V}$  is called simple if for any  $f \in \mathbb{C}[X(\Sigma)] \setminus \{0\}$  decomposed as  $f = \sum m_\Gamma t_\Gamma$  in the basis of multicurves, one has*

$$(\max) \quad v(f) = \max \{v(t_\Gamma) \mid m_\Gamma \neq 0\}.$$

*Remark 2.2.* — It is not clear at this point why the set of simple valuations should be preserved by  $\text{Aut}(X(\Sigma))$ , since we have not uniquely characterized our linear basis.

**PROPOSITION 2.3** (Measured laminations are simple valuations). — *Fix  $\lambda$  a measured lamination and denote by  $i(\cdot, \cdot)$  the intersection pairing between measured laminations. The map  $v_\lambda : t_\alpha \mapsto i(\lambda, \alpha)$  for  $\alpha \in \pi_1(\Sigma)$  extends to a unique simple valuation  $v_\lambda \in \mathcal{V}$ .*

*Proof.* — First treat the case of the measured lamination associated to a simple curve  $\delta$ . We define  $v_\delta$  on the basis of multicurves by the expression  $v_\delta(\prod_{i=1}^n t_{\gamma_i}) = \sum i(\delta, \gamma_i)$  and extend it to  $\mathbb{C}[X(\Sigma)]$  using equation (max). We must verify that both definitions coincide, meaning that given  $\alpha \in \pi_1(\Sigma)$  such that  $t_\alpha = \sum m_\Gamma t_\Gamma$  we have  $i(\delta, \alpha) = \max\{i(\delta, \Gamma) \mid m_\Gamma \neq 0\}$ , but this is the content of D. Thurston’s [Thu09, Lemma 12].

All that remains is to check the formula  $v_\delta(fg) = v_\delta(f) + v_\delta(g)$ . For this, consider the increasing filtration of  $\mathbb{C}[X(\Sigma)]$  defined by  $F_n = \text{Span}\{t_\gamma \mid \gamma \in \pi_1(\Sigma), i(\delta, \gamma) \leq n\}$ . Let  $k = v_\delta(f)$  and  $l = v_\delta(g)$  be such that  $f \in F_k \setminus F_{k-1}$  and  $g \in F_l \setminus F_{l-1}$ . It is equivalent to prove  $v_\delta(fg) = v_\delta(f) + v_\delta(g)$  and that  $fg$  is non zero in  $F_{k+l}/F_{k+l-1}$ . Hence, we are reduced to showing that the graded algebra  $\bigoplus_{n \in \mathbb{N}} F_n/F_{n-1}$  is an integral domain, which is [PS19, Theorem 12] (this can also be derived from the proof of [CM12, Theorem 5.3]).

For the general case, we extend the map  $v_\lambda$  to  $\mathbb{C}[X(\Sigma)]$  in the same way. The fact that it is indeed an extension and defines a simple valuation will follow from the case of simple curves by a limiting procedure. We know from [FLP79, Théorème 1.3] that any measured lamination  $\lambda$  is a limit of weighted simple curves  $m_j \delta_j$  in the sense that for all  $\alpha \in \pi_1(\Sigma)$  one has  $i(\lambda, \alpha) = \lim_{j \rightarrow \infty} m_j i(\delta_j, \alpha)$ . Hence by equation (max) the  $m_j v_{\delta_j}$  converge pointwise to  $v_\lambda$ , which indeed defines a valuation satisfying  $v_\lambda(t_\alpha) = i(\lambda, \alpha)$  for all  $\alpha \in \pi_1(\Sigma)$ . One could also observe that the subset of simple valuations is closed in  $\mathcal{V}$  for the topology of pointwise convergence.  $\square$

As a consequence of this proposition, we can identify a measured lamination  $\lambda$  with the corresponding valuation  $v_\lambda$ . Moreover, the usual topology on the space of measured laminations ML is the topology of weak convergence of lengths: in other terms, it is the topology induced by the embedding of ML in  $\mathcal{V}$ .

**DEFINITION 2.4** (Strict valuations). — *A valuation  $v \in \mathcal{V}$  is strict if for all multicurves  $\Gamma \neq \Delta$ , we have  $v(t_\Gamma) \neq v(t_\Delta)$ .*

**DEFINITION 2.5** (Positive valuations). — *A valuation  $v \in \mathcal{V}$  is positive when  $v(t_\gamma) > 0$  for every simple curve  $\gamma$ .*

*Remark 2.6.* — Let us comment on the relations between simple, strict and positive:

- (i) For simple valuations it is equivalent to be positive on  $t_\gamma$  for all simple  $\gamma$ , on  $t_\alpha$  for all non-trivial  $\alpha \in \pi_1(\Sigma)$ , or on non-constant elements in  $\mathbb{C}[X(\Sigma)]$ .
- (ii) (Strict implies positive). By definition  $v(t_\emptyset) = v(1) = 0$ , it follows that if  $v$  is strict, then for any simple curve  $\gamma$  we must have  $v(t_\gamma) > 0$ .

- (iii) (Strict implies simple). For any valuation  $v \in \mathcal{V}$ , the relation  $v(f + g) \leq \max\{v(f), v(g)\}$  is an equality as soon as  $v(f) \neq v(g)$ . Therefore strict valuations are simple since they automatically satisfy the (max) relation.

Recall that Thurston introduced a natural measure on the space of measured laminations that we use freely in this section. Masur showed that up to scaling, it is the only  $\text{Mod}(\Sigma)$ -invariant measure on ML in its Lebesgue class. Therefore it is proportional to the Borelian measure assigning to every open set  $U$  the limit as  $r \rightarrow \infty$  of the number of multicurves in the dilated set  $r \cdot U$  divided by  $r^{6g-6}$ . This may serve as a definition.

**PROPOSITION 2.7** (Most measured laminations are strict). — *The set of measured laminations  $\lambda$  such that  $v_\lambda$  is strict has full measure in ML.*

*Proof.* — Let us first show that the set of  $\lambda \in \text{ML}$  such that  $v_\lambda$  is positive has full measure. By [Pap86a, Section 3, Lemma 2], for every simple curve  $\gamma$ , the set  $N(\gamma) = \{\lambda \in \text{ML} \mid i(\gamma, \lambda) \neq 0\}$  is the complement of a codimension-1 PL-submanifold, in particular it has measure 0 (see also [Pap86b, Section 4]). The set of positive laminations is the intersection of all  $N(\gamma)$ , so it has full measure.

Inside that set of positive laminations, the complement to the set of strict valuations is the union over the countable set of pairs of distinct non-empty multicurves, of the  $N(\Gamma_1, \Gamma_2) = \{\lambda \in \text{ML} \mid i(\lambda, \Gamma_1) = i(\lambda, \Gamma_2)\}$ , so the result follows from the next lemma. □

**LEMMA 2.8** (Intersection with distinct multicurves seldom coincides). — *For distinct multicurves  $\Gamma_1 \neq \Gamma_2$ , the set  $N(\Gamma_1, \Gamma_2) = \{\lambda \in \text{ML} \mid i(\lambda, \Gamma_1) = i(\lambda, \Gamma_2)\}$  has measure 0 in the set of positive measured laminations.*

*Proof.* — The space ML has a piecewise linear (PL) structure (given by Dehn–Thurston coordinates or train tracks) for which the functions  $f_j(\lambda) = i(\Gamma_j, \lambda)$  are piecewise linear. In particular there exists an atlas  $(U_k, \phi_k)_{k \in \mathbb{N}}$  of ML such that all functions  $f_j \circ \phi_k^{-1}$  are PL. The subset  $W = \{\lambda \in \text{ML} \mid \exists k \in \mathbb{N}, f_1 \circ \phi_k^{-1} \text{ or } f_2 \circ \phi_k^{-1} \text{ is not smooth at } \phi_k(\lambda)\}$  is a PL-subcomplex of positive codimension so its complement has full measure.

Let  $x \in \text{ML} \setminus W$ , choose  $k \in \mathbb{N}$  such that  $x \in U_k$ , and denote by  $V$  the connected component of  $\phi_k(U_k \setminus W)$  containing  $\phi_k(x)$ . Then  $f_1$  and  $f_2$  are linear on  $V$ : it is sufficient to show that these two maps are distinct to conclude that the locus where they coincide has zero measure.

Consider a collection of  $9g - 9$  curves  $\gamma_1, \dots, \gamma_{9g-9}$  such that

$$\lambda = \mu \iff i(\lambda, \gamma_l) = i(\mu, \gamma_l) \text{ for } l = 1, \dots, 9g - 9.$$

Applying a power of a pseudo-Anosov map whose attracting point belongs to  $\phi_k^{-1}(V)$ , we may suppose that the curves  $\gamma_l$  belong to  $\phi_k^{-1}(V)$  (up to a scalar). As we have  $f_1(\phi_k(\gamma_l)) \neq f_2(\phi_k(\gamma_l))$  for some  $l$ , we can conclude that  $f_1$  and  $f_2$  are distinct. □

### 3. Domination of valuations

DEFINITION (Order structure). — We shall work with the partial order structure on  $\mathcal{V}$  called domination and defined by  $v \leq w$  if  $v(f) \leq w(f)$  for all  $f \in \mathbb{C}[X(\Sigma)]$ . A valuation  $u$  is called untameable if  $u \leq v$  implies  $v = Cu$  for some  $C \in \mathbb{R}$ .

The set  $\mathcal{U}$  of untameable valuations is clearly preserved by the action of  $\text{Aut}(X(\Sigma))$ . The main purpose of this subsection is to obtain the following theorem, whose proof consists in a compilation of known results.

THEOREM 3.1 (Domination by measured laminations). — For all  $v \in \mathcal{V}$ , there exists a unique  $\lambda \in \text{ML}$  such that  $v \leq v_\lambda$  and satisfying  $v(t_\alpha) = v_\lambda(t_\alpha)$  for all  $\alpha \in \pi_1(\Sigma)$ .

Recall that  $\mathbb{C}[X(\Sigma)]$  is an integral domain (see [PS19, Theorem 12], again this can also be derived from the proof of [CM12, Theorem 5.3]), so that we can consider its field of fraction  $K = \mathbb{C}(X(\Sigma))$ . We extend  $v$  to  $K$  by setting  $v(f/g) = v(f) - v(g)$ . Notice that our sign convention differs from the standard one in valuation theory: for instance our definition of the valuation ring associated to  $v$  is  $\mathcal{O}_v = \{f \in K \mid v(f) \leq 0\}$ . We first recall the standard construction of the so-called tautological representation.

LEMMA (Tautological representation). — There exists a finite extension  $\widehat{K}$  of  $K$  along with a representation  $\rho : \pi_1(\Sigma) \rightarrow \text{SL}_2(\widehat{K})$  such that  $\text{Tr } \rho(\alpha) = t_\alpha$  for all  $\alpha \in \pi_1(\Sigma)$ .

Proof. — This follows from classical arguments in geometric invariant theory. In a nutshell, consider an algebraic closure  $\overline{K}$  of  $K$ . One may interpret the inclusion  $\mathbb{C}[X(\Sigma)] \rightarrow \overline{K}$  as a  $\overline{K}$ -point of  $X(\Sigma)$ . Since the map  $\text{Hom}(\pi_1(\Sigma), \text{SL}_2(\overline{K})) \rightarrow X(\Sigma)(\overline{K})$  is surjective, we have a representation  $\rho : \pi_1(\Sigma) \rightarrow \text{SL}_2(\overline{K})$  satisfying  $\text{Tr } \rho(\gamma) = t_\gamma \in \overline{K}$ . As  $\pi_1(\Sigma)$  is finitely generated, the coefficients of  $\rho$  may be taken in a finite extension of  $K$ . One may also consult [Mar16, Proposition 3.3] for a down-to-earth proof using K. Saito’s theorem which even shows that a quadratic extension suffices. □

Proof of Theorem 3.1. — There exists (see [Vaq07] for a proof) an archimedean valuation  $\widehat{v} : \widehat{K}^* \rightarrow \mathbb{R}$  extending  $v$ , so we may consider the Bass–Serre real tree  $T$  associated to the pair  $(\widehat{K}, \widehat{v})$ .

We know from [MO93, Corollary III.1.2] that there exists a measured lamination  $\mu$  with associated real tree  $T_\mu$  and an equivariant morphism of  $\mathbb{R}$ -trees  $\Phi : T_\mu \rightarrow T$  which decreases the distance (by Step 1 of Lemma I.1.1). Thus for any  $\alpha \in \pi_1(\Sigma)$ , the translation length of the action of  $\alpha$  on  $T_\mu$ , which equals  $i(\mu, \alpha)$ , is greater than the translation length of  $\alpha$  acting on  $T$  which is  $\max\{0, 2\widehat{v}(t_\alpha)\} = 2v(t_\alpha)$ .

It follows in particular that  $2v(t_\gamma) \leq v_\mu(t_\gamma)$  for every simple curve  $\gamma$ , therefore this holds also over multicurves. For any  $f \in \mathbb{C}[X(\Sigma)]$  expanded as  $f = \sum w_\Gamma t_\Gamma$ , we get

$$v(f) \leq \max \{v(t_\Gamma) \mid w_\Gamma \neq 0\} \leq \frac{1}{2} \max \{v_\mu(t_\Gamma) \mid w_\Gamma \neq 0\} = \frac{1}{2} v_\lambda(f)$$

where the last equality follows by Proposition 2.3. Hence the first part of the theorem holds if one sets  $\lambda = \frac{1}{2}\mu$ .

If we prove that  $\Phi$  is an isometry on its image then we are done. Indeed this implies that the translation lengths of the actions of  $\alpha \in \pi_1(\Sigma)$  on  $T_\mu$  and  $T$  coincide, in other words that  $v(t_\alpha) = \frac{1}{2}v_\lambda(t_\alpha)$ . But by Skora’s theorem [Sko96], the morphism of  $\mathbb{R}$ -trees  $\Phi : T_\mu \rightarrow T$  mentioned above is an isometry on its image if and only if there does not exist any free subgroup  $F_2 \subset \pi_1(\Sigma)$  stabilizing a non-trivial edge in  $\Phi(T_\mu)$ .

Hence suppose by contradiction there exist  $\alpha, \beta \in \pi_1(\Sigma)$  generating a free subgroup which fixes a non-trivial edge in  $T$  of length  $l$ . Following for instance [Ota12, Section 4.2], this implies that up to conjugation, the tautological representation restricted to  $F_2$  has values in

$$G_l = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid ad - bc = 1, v(a) \leq 0, v(b) \leq 0, v(c) \leq -l, v(d) \leq 0 \right\}.$$

If  $\mathcal{M}_v^l$  denotes the ideal of  $\mathcal{O}_v$  defined by the equation  $v \leq -l$  then  $G_l$  consists precisely of those elements in  $\mathrm{SL}_2(\mathcal{O}_v)$  projecting to triangular matrices in  $\mathrm{SL}_2(\mathcal{O}_v/\mathcal{M}_v^l)$ . The commutator  $\rho([\alpha, \beta])$  is then unipotent in this quotient and we get  $\mathrm{Tr} \rho([\alpha, \beta]) = 2 \pmod{\mathcal{M}_v^l}$ . This means that  $v(t_{[\alpha, \beta]} - 2) \leq -l$ , contradicting the fact that  $v$  is non-negative over  $\mathbb{C}[X(\Sigma)]$ .  $\square$

**COROLLARY 3.2** (Simple valuations are measured laminations). — *A valuation is simple if and only if it has the form  $v_\lambda$  for some  $\lambda \in \mathrm{ML}$ . In particular  $\mathrm{ML}$  is closed.*

**COROLLARY 3.3** (Untameable implies Simple). — *We have an inclusion  $\mathcal{U} \subset \mathrm{ML}$ .*

Recall that a lamination is called *maximal* if it is filling (that is, the valuation  $v_\lambda$  is positive) and its complementary regions are triangles; or equivalently, when there are no lamination with a strictly larger support. A lamination is *uniquely ergodic* if it supports a unique transverse measure up to a scalar.

**LEMMA 3.4.** — *Almost all measured laminations are strict, maximal and uniquely ergodic. Such laminations are untameable. In particular  $\mathcal{U}$  is dense in  $\mathrm{ML}$ .*

*Proof.* — Almost all measured laminations are maximal by [LM08, Lemma 2.3], uniquely ergodic by a theorem of Masur [Mas82, Theorem 2], and strict by Lemma 2.7.

If  $\lambda$  is maximal and uniquely ergodic and  $\mu$  is another measured lamination such that  $v_\lambda \leq v_\mu$ , then the support of  $\lambda$  is included in the support of  $\mu$ . By maximality, the support of  $\mu$  is the same as the support of  $\lambda$ : only the transverse measures may differ. By unique ergodicity,  $\mu$  and  $\lambda$  must be proportional.

Let  $\lambda$  be a strict, maximal and uniquely ergodic measured lamination. Now suppose there exists another valuation  $w \in \mathcal{V}$  with  $v \leq w$ . Applying Theorem 3.1, there exists another measured lamination  $\mu$  with  $w \leq v_\mu$  and  $w(t_\alpha) = v_\mu(t_\alpha)$  for all  $\alpha \in \pi_1(\Sigma)$ . From the previous discussion there exists  $C \geq 1$  such that  $\mu = C\lambda$ . But  $w$  and  $C\lambda$  are strict and coincide on simple curves, so that  $w = C\lambda$  and we are done.  $\square$

**COROLLARY 3.5.** — *The action of  $\mathrm{Aut}(X(\Sigma))$  on  $\mathcal{V}$  preserves  $\mathrm{ML}$ .*

*Proof.* — The action of  $\mathrm{Aut}(X(\Sigma))$  on  $\mathcal{V}$  is continuous and preserves the subset  $\mathcal{U} \subset \mathrm{ML}$  of untameable valuations, so it also preserves its closure  $\overline{\mathcal{U}} = \mathrm{ML}$ .  $\square$

#### 4. Discrete valuations and disjointness

We call *discrete* a valuation in  $\mathcal{V}$  whose finite values belong to  $\mathbb{N}$ , and likewise a measured lamination whose associated valuation is discrete. One expects the underlying lamination to be a multicurve, this is almost true:

LEMMA (Discrete measured laminations). — *Discrete measured laminations are precisely the weighted multicurves  $\frac{1}{2}\Gamma$  such that the class of  $\Gamma$  in  $H_1(\Sigma, \mathbb{Z}/2\mathbb{Z})$  vanishes.*

*Proof.* — This lemma can be shown using Dehn's coordinates, see [FLP79, Exposé 6]. Let us give a proof more in the spirit of this note as it follows the lines of reasoning for Theorem 3.1. For this, observe that the Bass–Serre tree associated to a discrete valuation has a simplicial structure.

We extend the discrete valuation  $v$  on  $K$  to a discrete valuation  $\hat{v}$  on  $\widehat{K}$  whose Bass–Serre tree  $T$  is thus simplicial. The Morgan–Otal–Skora theorem reduces to a theorem of Stallings (see [Sha02, Chapter 2] for a nice account) stating that the action of  $\pi_1(\Sigma)$  on  $T$  is dual to a multicurve  $\Gamma$ . This implies  $v(t_\alpha) = \frac{1}{2}i(\Gamma, \alpha)$ , so  $v$  is associated to the measured lamination  $\lambda = \frac{1}{2}\Gamma$ .

Computing modulo  $\mathbb{Z}$  the weighted intersection of  $\frac{1}{2}\Gamma$  with any curve  $\gamma$ , one should get 0: by Poincaré duality this shows that the class of  $\Gamma$  in  $H_1(\Sigma, \mathbb{Z}/2\mathbb{Z})$  vanishes.  $\square$

Of course, an automorphism of  $\mathbb{C}[X(\Sigma)]$  must preserve the set of discrete valuations, and hence also the set of discrete simple valuations by Corollary 3.5. The previous lemma says that such a valuation has the form  $v = \frac{1}{2}v_\Gamma$  for a multicurve  $\Gamma$  which is null in homology modulo 2.

We wish to show that an algebraic automorphism actually preserves the set of discrete valuations coming from simple curves, and the relation of being disjoint. Thus we need an algebraic characterization for the number of components in a multicurve, as well as the disjointness property, in terms of their associated valuations.

For this recall the notion of valuation ring associated to the valuation  $v \in \mathcal{V}$ , defined by  $\mathcal{O}_v^+ = \{f \in \mathbb{C}[X(\Sigma)] \mid v(f) \leq 0\}$ ; its Krull dimension  $\dim \mathcal{O}_v^+$  is preserved by algebraic automorphisms. When  $\lambda = \frac{1}{2}\Gamma$  for a multicurve  $\Gamma$ , we have  $\mathcal{O}_\lambda^+ = \text{Span}\{t_\Delta \mid i(\Gamma, \Delta) = 0\}$ , which corresponds geometrically to the coordinate ring for the image of the restriction map  $X(\Sigma) \rightarrow X(\Sigma \setminus \Gamma)$ .

LEMMA 4.1 (Number of components). — *The number of distinct homotopy classes of simple curves in a multicurve  $\Gamma$  is equal to  $\text{codim } \mathcal{O}_\Gamma^+ = \dim \mathbb{C}[X(\Sigma)] - \dim \mathcal{O}_\Gamma^+$ .*

*Proof.* — Let  $\gamma_1, \dots, \gamma_n$  be the simple curves composing  $\Gamma$ . We let  $U \subset X(\Sigma)$  be the Zariski open subset of  $X(\Sigma)$  containing irreducible representations  $\rho$  satisfying  $\text{Tr } \rho(\gamma_i) \neq \pm 2$  for  $i = 1, \dots, n$ . The restriction map  $X(\Sigma) \rightarrow X(\Sigma \setminus \{\gamma_1, \dots, \gamma_n\})$  restricted to  $U$  and corestricted to its image is a fibration whose fibres are covered by the orbits of the (commuting) Goldman twist flows along  $\gamma_1, \dots, \gamma_n$  on the character variety. Since the fiber has dimension  $n$ , the image has codimension  $n$ , and the result follows.  $\square$

LEMMA 4.2 (Disjointness property). — *For any pair of non-parallel simple curves  $\gamma, \delta$  one has  $\dim \mathcal{O}_\gamma^+ \cap \mathcal{O}_\delta^+ \leq \dim \mathbb{C}[X(\Sigma)] - 2$ , with equality if and only if  $i(\gamma, \delta) = 0$ .*



*Proof.* — If  $\gamma$  and  $\delta$  are disjoint, Lemma 4.1 already gives  $\text{codim } \mathcal{O}_{\gamma \cup \delta}^+ = 2$  and the result follows from the observation that  $\mathcal{O}_{\gamma \cup \delta}^+ = \mathcal{O}_\gamma^+ \cap \mathcal{O}_\delta^+$ .

Suppose now that  $\gamma$  and  $\delta$  intersect minimally and set  $\Sigma' = \Sigma \setminus (\gamma \cup \delta)$ . The ring  $\mathcal{O}_\gamma^+ \cap \mathcal{O}_\delta^+$  is the image of the natural map  $\mathbb{C}[X(\Sigma')] \rightarrow \mathbb{C}[\Sigma]$ . Geometrically, it is the coordinate ring for the image of the restriction map  $X(\Sigma) \rightarrow X(\Sigma')$ , hence it is sufficient to show that  $\dim X(\Sigma') < \dim X(\Sigma) - 2$ . The surface  $\Sigma'$  is a disjoint union of connected surfaces with connected boundary: their fundamental group is either trivial or free with rank  $\geq 2$ . Denoting by  $F$  the number of simply connected components, we have  $\dim X(\Sigma') = -3\chi(\Sigma') + 3F$ .

But  $\chi(\Sigma') = \chi(\Sigma) + i(\gamma, \delta)$  so  $\dim X(\Sigma') = \dim X(\Sigma) - 3i(\gamma, \delta) + 3F$ , and we must show that  $i(\gamma, \delta) > F$ . Since  $\gamma \cup \delta$  is a taut union of simple curves, the polygonal components have at least 4 corners, so  $4F$  is smaller or equal to the total number of corners which is  $4i(\gamma, \delta)$ . But  $F = i(\gamma, \delta)$  implies that  $\Sigma'$  is a union of quadrilaterals, so  $F = \chi(\Sigma') = \chi(\Sigma) + i(\gamma, \delta)$  and thus  $\chi(\Sigma) = 0$ : a contradiction.  $\square$

**COROLLARY 4.3** (Action on the curve complex). — *The group  $\text{Aut}(X(\Sigma))$  preserves the set*

$$\mathcal{C} = \{v_\gamma | \gamma \text{ simple and non-separating}\} \cup \{v_{\gamma/2} | \gamma \text{ simple and separating}\}$$

along with its orthogonality relation  $v_\gamma \perp v_\delta \iff i(\gamma, \delta) = 0$  for  $v_\gamma, v_\delta \in \mathcal{C}$ .

Set  $\text{Mod}'(\Sigma) = \text{Mod}(\Sigma)$  if  $g \geq 3$  and  $\text{Mod}'(\Sigma) = \text{Mod}(\Sigma)/\langle \tau \rangle$  if  $g = 2$  where  $\tau$  denotes the hyperelliptic involution.

**THEOREM 4.4.** — *There is a natural split extension*

$$0 \rightarrow H^1(\Sigma, \mathbb{Z}/2\mathbb{Z}) \rightarrow \text{Aut}(X(\Sigma)) \rightarrow \text{Mod}'(\Sigma) \rightarrow 0.$$

*Proof.* — Let  $\phi$  be an automorphism of  $X(\Sigma)$ . By Corollary 4.3, it acts on the curve complex  $(\mathcal{C}, \perp)$  and Ivanov’s Theorem (see [Iva97] or [Luo00]) implies the existence of a unique element  $\varphi \in \text{Mod}(\Sigma)$  which acts in the same way: this defines the map  $\text{Aut}(X(\Sigma)) \rightarrow \text{Mod}(\Sigma)$ , which is a group morphism. Since  $\text{Mod}(\Sigma)$  naturally acts on  $X(\Sigma)$  by sending  $t_\gamma$  to  $t_{\varphi(\gamma)}$ , there is a natural section of the previous map whose kernel consists of mapping classes which fix all non-oriented simple curves. This reduces to the identity or the hyperelliptic involution in genus 2 (see for instance [FM12]) hence the surjection  $\text{Aut}(X(\Sigma)) \rightarrow \text{Mod}'(\Sigma)$ .

Suppose  $\phi \in \text{Aut}(X(\Sigma))$  acts trivially on  $\mathcal{C}$ . Fix a simple curve  $\gamma$  and consider the possible values of  $\phi(t_\gamma)$ . As  $\phi$  preserves the valuation  $v_\delta$  for every simple curve  $\delta$  we get  $v_\delta(\phi(t_\gamma)) = v_\delta(t_\gamma) = i(\delta, \gamma)$ . Write  $\phi(t_\gamma) = \sum m_\Gamma t_\Gamma$ : for any simple curve  $\delta$  which does not intersect  $\gamma$  one has  $0 = v_\delta(\phi(t_\gamma)) = \max\{i(\delta, \Gamma) \mid m_\Gamma \neq 0\}$ , so every  $\Gamma$  such that  $m_\Gamma \neq 0$  must be a family of parallel copies of  $\gamma$ . Hence  $\phi(t_\gamma)$  is a polynomial in  $t_\gamma$ , so  $\phi$  induces an automorphism of the subalgebra  $\mathbb{C}[t_\gamma]$ , thus  $\phi(t_\gamma) = a_\gamma t_\gamma + b_\gamma$  with  $a_\gamma \neq 0$ .

Lemma 4.5 shows that  $b_\gamma = 0$  for every non trivial simple curve  $\gamma$ , so  $\phi(\gamma) = a_\gamma t_\gamma$  and for any multicurve  $\Gamma$  with components  $\gamma_j$ , we thus have  $\phi(t_\Gamma) = a_\Gamma t_\Gamma$  where  $a_\Gamma = \prod a_{\gamma_j}$ . Lemma 4.6 says that  $a_\Gamma$  does not change when  $\Gamma$  undergoes a saddle-move (we give the definition after Lemma 4.5). But two multicurves are related by a sequence of saddle moves when they belong to the same homology class modulo

2, so  $a$  factors to a map  $a: H_1(\Sigma, \mathbb{Z}/2\mathbb{Z}) \rightarrow \mathbb{C}^*$ , which according to Lemma 4.7 is in fact a morphism. Thus  $a \in H^1(\Sigma, \mathbb{Z}/2\mathbb{Z})$  and we are done.  $\square$

LEMMA 4.5. — *For every non-trivial simple curve  $\gamma$ , we have  $b_\gamma = 0$  and  $a_\gamma = \pm 1$ .*

*Proof.* — Suppose  $\gamma$  is a non-separating simple curve. Then there is another one  $\delta$  which intersects it once and a tubular neighborhood of their union is a  $\pi_1$ -injectively embedded punctured torus. In that case apply  $\phi$  to the relation  $t_\gamma t_\delta = t_{\gamma\delta} + t_{\gamma\delta^{-1}}$  and decompose in the basis of multicurves to get  $b_\gamma = 0$  and then  $a_\gamma a_\delta = a_{\gamma\delta}$ . But the simple curve  $\gamma\delta$  also intersects  $\gamma$  once, so the same reasoning also gives  $a_{\gamma\delta} a_\gamma = a_\delta$ . Combined with the previous relation this implies  $a_\gamma a_\delta a_\gamma = a_\delta$ , hence  $a_\gamma = \pm 1$ .

From now on  $\gamma$  is a separating simple curve. There exists a non-separating simple curve  $\delta$  which intersects it twice with opposite signs. A neighborhood for their union  $\gamma \cup \delta$  is an embedded four holed sphere  $F \subset \Sigma$  with boundary components  $\eta_1, \eta_2, \eta_3, \eta_4$  such that the triple  $\eta_1, \eta_2, \gamma$  bounds a three holed sphere. (This fixes the configuration up to the action of the mapping class group.)

Since  $i(\gamma, \delta) = 2$  the  $\eta_j$  are non-trivial and as they are disjoint from  $\gamma$  and  $\delta$  they must be homotopically distinct from  $\delta$  and  $\gamma$ . Also, as  $\gamma$  is separating,  $\eta_1$  and  $\eta_2$  are both homotopically distinct from each of  $\eta_3$  and  $\eta_4$ .

Using the trace relation on  $t_\gamma t_\delta$  to resolve the intersections of  $\gamma \cup \delta$ , we decompose  $t_\gamma t_\delta = t_{\eta_1} t_{\eta_3} + t_{\eta_2} t_{\eta_4} - t_\theta - t_\zeta$  where  $\zeta, \theta \subset F$  are two simple curves both intersecting twice  $\gamma$  and  $\delta$ , such that each triple  $\eta_1, \eta_3, \theta$  and  $\eta_2, \eta_4, \zeta$  bounds a three-holed sphere.

Considering intersection numbers as before we see that  $\theta, \zeta$  are homotopically distinct from each other and from  $\gamma, \delta, \eta_j$ : among all simple curves into play here, only the boundary components  $\eta_1$  and  $\eta_2$ , or  $\eta_3$  and  $\eta_4$ , may belong the same class. So when we decompose into the basis of multicurves after applying  $\phi$  to the previous relation, we find that  $b_\gamma = 0$  and then  $a_\gamma a_\delta = a_\zeta$ . Since  $\gamma$  is separating and  $\delta$  is not,  $\zeta$  must also be non-separating, and the previous discussion implies  $a_\delta = \pm 1$  and  $a_\zeta = \pm 1$ , so  $a_\gamma = \pm 1$ .  $\square$

In order to introduce the notion of a saddle move, we will think of multicurves as isotopy classes of 1-submanifolds in  $\Sigma$  up to adding or removing a trivial component. Two multicurves are related by a saddle move if they are represented by two 1-manifolds which differ in a disc as in Figure 4.1.




Figure 4.1. *Saddle move*

It is an easy exercise to show that two multicurves are related by a sequence of saddle moves if and only if they belong to the same homology class modulo 2. Let us sketch the argument. Let  $\Gamma_1, \Gamma_2$  be two multicurves homologous modulo 2 in  $\Sigma$  and suppose that they intersect transversely. Any connected component of  $\Sigma \setminus (\Gamma_1 \cup \Gamma_2)$  containing a corner should contain an even number of those, and two consecutive corners are cancelled by a saddle move. By induction, we are reduced to the case where  $\Gamma_1 \cap \Gamma_2 = \emptyset$ . As  $\Gamma_1$  and  $\Gamma_2$  are homologous modulo 2, there exists an embedded

subsurface  $S \subset \Sigma$  with  $\partial S = \Gamma_1 \cup \Gamma_2$ , and decomposing this cobordism into saddles shows how to transform  $\Gamma_1$  into  $\Gamma_2$ .

Set  $t_\epsilon = 2$  for the trace of a trivial component  $\epsilon$  and extend the  $t_\Gamma$  by multiplicativity on all components  $\gamma_j$  to encompass the more general notion of multicurves. This remains consistent with the trace relation. As  $\phi(t_\epsilon) = t_\epsilon$  we have  $a_\epsilon = 1$ , and according to the previous lemma we now have  $a_\Gamma = \prod a_{\gamma_j}$  defined as a function on multicurves with values in  $\pm 1$ .

LEMMA 4.6. — *The function  $a$  so defined on multicurves is invariant by saddle move.*

*Proof.* — Before starting the proof, observe that a saddle move changes the number of components of a 1-submanifold by 0 (type 0) or 1 (type 1).

Consider first a saddle move of type 0: ignoring the components that are not affected, one can assume it transforms a simple curve  $\eta$  into another one  $\theta$ . Basing the fundamental group at the singular point appearing half way during the saddle move, we see that there exists two simple curves  $\gamma, \delta$  intersecting once such that  $\eta = \gamma\delta$  and  $\theta = \gamma\delta^{-1}$ . As in the proof of Lemma 4.5, we deduce that  $a_\eta = a_\gamma a_\delta = a_\gamma a_{\delta^{-1}} = a_\theta$ , which concludes this case.

Now consider the case of a saddle move of type 1. Again, ignoring the components that are not affected, we observe that we must prove the formula  $a_\gamma = a_{\eta_1} a_{\eta_2}$  where  $\gamma, \eta_1, \eta_2$  are three boundary components of an embedded pair of pants  $P$  in  $\Sigma$ . Since  $a_\gamma = \pm 1$  by Lemma 4.5, this can be written more symmetrically as  $a_\gamma a_{\eta_1} a_{\eta_2} = 1$ . This formula obviously holds if one of the three boundary components is homotopically trivial so that we can suppose that  $P$  is incompressible in  $\Sigma$ .

Thus we may consider another embedded 3-holed sphere bounded by  $\gamma, \eta_3, \eta_4$  such that the  $\eta_j$  bound an embedded 4-holed sphere  $F$  as in the proof of the previous lemma. This time, using the trace relation on  $t_\theta t_\delta$  to resolve the intersections of  $\theta \cup \delta$ , we decompose  $t_\theta t_\delta = t_{\eta_1} t_{\eta_2} + t_{\eta_3} t_{\eta_4} - t_\nu - t_\gamma$  where the simple curve  $\nu$  corresponds to the image of  $\theta$  under the half Dehn twist along  $\delta$ . Applying  $\phi$  to it and decomposing into the basis of multicurves we find that  $a_\gamma = a_{\eta_1} a_{\eta_2}$ , and the lemma follows.  $\square$

LEMMA 4.7. — *The map induced on homology  $a: H_1(\Sigma; \mathbb{Z}/2\mathbb{Z}) \rightarrow \mathbb{C}^*$  is a morphism.*

*Proof.* — Let  $\alpha, \beta$  be two classes in  $H_1(\Sigma, \mathbb{Z}/2\mathbb{Z})$ . They can be represented by simple curves  $\gamma, \delta$  which are either disjoint (if  $\alpha \cdot \beta = 0$ ) and then  $\alpha + \beta$  is represented by the disjoint union  $\gamma \cup \delta$ , or else intersect once (if  $\alpha \cdot \beta = 1$ ) and then  $\alpha + \beta$  is represented by the product  $\gamma\delta$ . In any case, we already proved that  $a_{\alpha\beta} = a_\alpha a_\beta$ .  $\square$

## BIBLIOGRAPHY

- [AS16] Javier Aramayona and Juan Souto, *Rigidity phenomena in the mapping class group*, Handbook of Teichmüller theory. Vol. VI, IRMA Lectures in Mathematics and Theoretical Physics, vol. 27, European Mathematical Society, 2016, pp. 131–165.  $\uparrow 592$
- [Che13] Gaëtan Chenevier, *Représentations galoisiennes automorphes et conséquences arithmétiques des conjectures de Langlands et Arthur*, Mémoire de HDR <https://arxiv.org/abs/1307.5170>, 2013.  $\uparrow 593$

- [CM12] Laurent Charles and Julien Marché, *Multicurves and regular functions on the representation variety of a surface in  $SU(2)$* , *Comment. Math. Helv.* **87** (2012), 409–431. ↑594, 596
- [P17] Corrado De Concini and Claudio Procesi, *The invariant theory of matrices*, University Lecture Series, vol. 69, American Mathematical Society, 2017. ↑593
- [ÈH74] M. H. Èl’-Huti, *Cubic surfaces of Markov type*, *Mat. Sb., N. Ser.* **93(135)** (1974), 331–346, 487. ↑592
- [FLP79] A. Fathi, F. Laudenbach, and V. Poénaru, *Comment Thurston compactifie l’espace de Teichmüller*, *Travaux de Thurston sur les surfaces. Seminaire Orsay, Astérisque*, no. 66-67, Société Mathématique de France, 1979, pp. 139–150. ↑594, 598
- [FM12] Benson Farb and Dan Margalit, *A primer in mapping class groups*, Princeton Mathematical Series, vol. 49, Princeton University Press, 2012. ↑599
- [Iva97] Nikolai Ivanov, *Automorphisms of complexes of curves and of Teichmüller spaces*, *Internat. Math. Res. Notices* **14** (1997), 651–666. ↑599
- [LM08] Elon Lindenstrauss and Maryam Mirzakhani, *Ergodic theory of the space of measured laminations*, *Int. Math. Res. Not.* **2008** (2008), no. 4, article no. rnm126. ↑597
- [Luo00] Feng Luo, *Automorphisms of the complex of curves*, *Topology* **39** (2000), no. 2, 283–298. ↑599
- [Mar16] Julien Marché, *Character varieties in  $SL_2$  and skein algebras*, *Topology, Geometry and Algebra of low dimensional manifolds*, RIMS, Kyoto, vol. 1991, Kyoto University, 2016, pp. 27–42. ↑596
- [Mas82] Howard Masur, *Interval exchange transformations and measured foliations*, *Ann. Math.* **115** (1982), no. 1, 169–200. ↑597
- [MO93] John W. Morgan and Jean-Pierre Otal, *Relative growth rates of closed geodesics on a surface under varying hyperbolic structures*, *Comment. Math. Helv.* **68** (1993), no. 2, 171–208. ↑596
- [Ota12] Jean-Pierre Otal, *Compactification of spaces of representations, after Culler, Morgan and Shalen*, *Berkovich Spaces and Applications* (A. Ducros, C. Favre, and J. Nicaise, eds.), *Lecture Notes in Mathematics*, vol. 2119, 2012, pp. 367–413. ↑592, 597
- [Pap86a] Athanase Papadopoulos, *Deux remarques sur la géométrie symplectique de l’espace des feuilletages mesurés sur une surface*, *Ann. Inst. Fourier* **36** (1986), no. 2, 127–141. ↑595
- [Pap86b] ———, *Geometric intersection functions and hamiltonian vector fields on the space of measured laminations*, *Pac. J. Math.* **124** (1986), no. 2, 375–402. ↑595
- [PS00] Józef H. Przytycki and Adam S. Sikora, *On skein algebras and  $Sl_2(\mathbb{C})$ -character varieties*, *Topology* **39** (2000), no. 1, 115–148. ↑593
- [PS19] ———, *Skein algebras of surfaces*, *Trans. Am. Math. Soc.* **371** (2019), no. 2, 1309–1332. ↑594, 596
- [Sha02] Peter B. Shalen, *Representations of 3-manifold groups*, *Handbook of geometric topology*, North-Holland, 2002, pp. 955–1044. ↑598
- [Sko96] Richard K. Skora, *Splittings of surfaces*, *J. Am. Math. Soc.* **9** (1996), no. 2, 605–616. ↑597
- [Thu09] D. Thurston, *Geometric intersection of curves on surfaces*, preprint, <https://dpthurst.pages.iu.edu/DehnCoordinates.pdf>, 2009. ↑594
- [Vaq07] Michel Vaquié, *Extension d’une valuation*, *Trans. Am. Math. Soc.* **359** (2007), no. 7, 3439–3481. ↑596

Manuscript received on 9th October 2019,  
revised on 28th May 2020,  
accepted on 29th July 2020.

Recommended by Editor S. Cantat.  
Published under license CC BY 4.0.



This journal is a member of Centre Mersenne.



Julien MARCHÉ  
Sorbonne Université, IMJ-PRG,  
75252 Paris cedex 05, (France)  
julien.marche@imj-prg.fr

Christopher-Lloyd SIMON  
ENS de Lyon,  
15 parvis René Descartes,  
69342 Lyon Cedex 07, (France)  
christopher-lloyd.simon@ens-lyon.fr